

Fermat Last Theorem was Proved in 1991

Jiang, Chun-xuan

Institute for Basic Research, Palm Harbor, FL34682-1577, USA

P. O. Box 3924, Beijing 100854, China

jiangchunxuan@sohu.com, cxjiang@mail.bcf.net.cn

Abstract: We found out a new method for proving Fermat last theorem (FLT) on the afternoon of October 25, 1991.

We proved FLT at one stroke for all prime exponents $p > 3$, It led to the discovery to calculate $n = 15, 21, 35, 105, \dots$. To this date, no one disprove this proof. Anyone can not deny it, because it is a simple and marvelous proof. It can fit in the margin of Fermat book.

[Jiang, Chun-xuan. **Fermat Last Theorem was Proved in 1991**. *Academ Arena* 2015;7(12):73-77]. (ISSN 1553-992X). <http://www.sciencepub.net/academia>. 8. doi:[10.7537/marsaaj071215.08](https://doi.org/10.7537/marsaaj071215.08).

Keywords: Fermat last theorem (FLT); calculate; cyclotomic real number; cyclotomic field

We found out a new method for proving Fermat last theorem (FLT) on the afternoon of October 25, 1991. We proved FLT at one stroke for all prime exponents $p > 3$, It led to the discovery to calculate $n = 15, 21, 35, 105, \dots$. To this date, no one disprove this proof. Anyone can not deny it, because it is a simple and marvelous proof. It can fit in the margin of Fermat book.

In 1974 we found out Euler formula of the cyclotomic real numbers in the cyclotomic fields [1].

$$\exp\left(\sum_{i=1}^{n-1} t_i J^i\right) = \sum_{i=1}^n S_i J^{i-1}, \quad (1)$$

where J denotes a n -th root of unity, $J^n = 1$, n is an odd number, t_i are the real numbers.

S_i is called the complex hyperbolic functions of order n with $n-1$ variables,

$$S_i = \frac{1}{n} \left[e^A + 2 \sum_{j=1}^{\frac{n-1}{2}} (-1)^{(i-1)j} e^{B_j} \cos\left(\theta_j + (-1)^j \frac{(i-1)j\pi}{n}\right) \right], \quad (2)$$

where

$$A = \sum_{\alpha=1}^{n-1} t_\alpha, \quad B_j = \sum_{\alpha=1}^{n-1} t_\alpha (-1)^{\alpha j} \cos \frac{\alpha j \pi}{n}, \quad \theta_j = (-1)^{j+1} \sum_{\alpha=1}^{n-1} t_\alpha (-1)^{\alpha j} \sin \frac{\alpha j \pi}{n},$$

$$A + 2 \sum_{i=1}^{\frac{n-1}{2}} B_i = 0 \quad (3)$$

Using (1) the cyclotomic theory may extend to totally real number fields. It is called the hypercomplex variable theory [1]. (2) may be written in the matrix form

$$\begin{bmatrix} S_1 \\ S_2 \\ S_3 \\ \dots \\ S_n \end{bmatrix} = \frac{1}{n} \begin{bmatrix} 1 & 1 & 0 & \dots & 0 \\ 1 & -\cos \frac{\pi}{n} & -\sin \frac{\pi}{n} & \dots & -\sin \frac{(n-1)\pi}{2n} \\ 1 & \cos \frac{2\pi}{n} & \sin \frac{2\pi}{n} & \dots & -\sin \frac{(n-1)\pi}{n} \\ \dots & \dots & \dots & \dots & \dots \\ 1 & \cos \frac{(n-1)\pi}{n} & \sin \frac{(n-1)\pi}{n} & \dots & -\sin \frac{(n-1)^2\pi}{2n} \end{bmatrix} \begin{bmatrix} e^A \\ 2e^{B_1} \cos \theta_1 \\ 2e^{B_1} \sin \theta_1 \\ \dots \\ 2 \exp(B_{\frac{n-1}{2}}) \sin(\theta_{\frac{n-1}{2}}) \end{bmatrix} \tag{4}$$

where $(n-1)/2$ is an even number.

From (4) we may obtain its inverse transformation

$$\begin{bmatrix} e^A \\ e^{B_1} \cos \theta_1 \\ e^{B_1} \sin \theta_1 \\ \dots \\ \exp(B_{\frac{n-1}{2}}) \sin(\theta_{\frac{n-1}{2}}) \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 & \dots & 1 \\ 1 & -\cos \frac{\pi}{n} & \cos \frac{2\pi}{n} & \dots & \cos \frac{(n-1)\pi}{n} \\ 0 & -\sin \frac{\pi}{n} & \sin \frac{2\pi}{n} & \dots & \sin \frac{(n-1)\pi}{n} \\ \dots & \dots & \dots & \dots & \dots \\ 0 & -\sin \frac{(n-1)\pi}{2n} & -\sin \frac{(n-1)\pi}{n} & \dots & -\sin \frac{(n-1)^2\pi}{2n} \end{bmatrix} \begin{bmatrix} S_1 \\ S_2 \\ S_3 \\ \dots \\ S_n \end{bmatrix} \tag{5}$$

From (5) we have

$$e^A = \sum_{i=1}^n S_i, e^{B_j} \cos \theta_j = S_1 + \sum_{i=1}^{n-1} S_{1+i} (-1)^{ij} \cos \frac{ij\pi}{n},$$

$$e^{B_j} \sin \theta_j = (-1)^{j+1} \sum_{i=1}^{n-1} S_{1+i} (-1)^{ij} \sin \frac{ij\pi}{n}. \tag{6}$$

In (3) and (6) t_i and S_i have the same formulas such that every factor of n has a Fermat equation. Assume $S_1 \neq 0, S_2 \neq 0, S_i = 0$ where $i = 3, 4, \dots, n, S_i = 0$ are $n-2$ indeterminate equations with $n-1$ variables. From (6) we have

$$e^A = S_1 + S_2, e^{2B_j} = S_1^2 + S_2^2 + 2S_1S_2(-1)^j \cos \frac{j\pi}{n}. \tag{7}$$

From (3) and (7) we may obtain the Fermat equation

$$\exp \left(A + 2 \sum_{j=1}^{\frac{n-1}{2}} B_j \right) = (S_1 + S_2) \prod_{j=1}^{\frac{n-1}{2}} (S_1^2 + S_2^2 + 2S_1S_2(-1)^j \cos \frac{j\pi}{n}) = S_1^n + S_2^n = 1. \tag{8}$$

Theorem. Fermat last theorem has no rational solutions with $S_1S_2 \neq 0$ for all odd exponents.

Proof. The proof of FLT is difficult when n is an odd prime. We consider n is a composite number.

Let $n = \prod n_i$, where n_i ranges over all odd number. From (3) we have

$$\exp\left(A + 2 \sum_{j=1}^{\frac{f-1}{2}} \frac{B_n}{f^j}\right) = \left[\exp\left(\sum_{\alpha=1}^{\frac{n-1}{f}} t_{f\alpha}\right)\right]^f \quad (9)$$

From (7) we have

$$\exp\left(A + 2 \sum_{j=1}^{\frac{f-1}{2}} \frac{B_n}{f^j}\right) = S_1^f + S_2^f \quad (10)$$

where f is a factor of n . From (9) and (10) we may obtain Fermat equation

$$\exp\left(A + 2 \sum_{j=1}^{\frac{f-1}{2}} \frac{B_n}{f^j}\right) = S_1^f + S_2^f = \left[\exp\left(\sum_{\alpha=1}^{\frac{n-1}{f}} t_{f\alpha}\right)\right]^f \quad (11)$$

Every factor of n has a Fermat equation. From (11) we have

$$f = 1, B_n = B_0 = 0, \quad e^A = S_1 + S_2 = \exp\left(\sum_{\alpha=1}^{n-1} t_{\alpha}\right) \quad (12)$$

$$f = n, t_n = t_0 = 0, \quad \exp\left(A + 2 \sum_{i=1}^{\frac{n-1}{2}} B_j\right) = S_1^n + S_2^n = 1 \quad (13)$$

$$f = 3, \exp\left(A + 2 \sum_{\alpha=1}^{\frac{n-1}{3}} B_{3\alpha}\right) = S_1^3 + S_2^3 = \left[\exp\left(\sum_{\alpha=1}^{\frac{n-1}{3}} t_{3\alpha}\right)\right]^3 \quad (14)$$

If $S_1 = 1, S_2 = 0$ and $S_1 = 0, S_2 = 1$, then $A = B_j = 0$. Euler proved (13), therefore (11) has no rational solutions with $S_1 S_2 \neq 0$ (and so no integer solutions with $S_1 S_2 \neq 0$) for all odd exponents f . (11) and (13) can fit in the margin of Fermat book.

Let $n = 3p$ where p is an odd prime. From (3) and (7) we may derive Fermat equations

$$\exp\left(A + 2 \sum_{i=1}^{\frac{3p-1}{2}} B_j\right) = S_1^{3p} + S_2^{3p} = (S_1^p)^3 + (S_2^p)^3 = 1 \quad (15)$$

$$\exp\left(A + 2 \sum_{\alpha=1}^{p-1} B_p\right) = S_1^3 + S_2^3 = \left[\exp\left(\sum_{\alpha=1}^{p-1} t_{3\alpha}\right)\right]^3 \quad (16)$$

$$\exp\left(A + 2 \sum_{i=1}^{\frac{p-1}{2}} B_{3j}\right) = S_1^p + S_2^p = \left[\exp(t_p + t_{2p})\right]^p \quad (17)$$

Euler proved (15) and (16), therefore (17) have no rational solutions with $S_1 S_2 \neq 0$ (and so no integer solutions with $S_1 S_2 \neq 0$) for any odd prime $p > 3$. (15)-(17) can fit in the margin

Let $n = 5p$ where p is an odd prime. From (3) and (7) we may derive Fermat equations

$$\exp\left(A + 2 \sum_{j=1}^{\frac{5p-1}{2}} B_j\right) = S_1^{5p} + S_2^{5p} = 1 \quad (18)$$

$$\exp(A + 2B_p + 2B_{2p}) = S_1^5 + S_2^5 = [\exp \sum_{\alpha=1}^{p-1} t_{5\alpha}]^5 \quad (19)$$

$$\exp(A + 2 \sum_{j=1}^{\frac{p-1}{2}} B_{5j}) = S_1^p + S_2^p = [\exp(\sum_{\alpha=1}^4 t_{p\alpha})]^p \quad (20)$$

(18)-(20) can fit in the margin.

Let $n = 7p$ where p is an odd prime. From (3) and (7) we may derive Fermat equations

$$\exp(A + 2 \sum_{i=1}^{\frac{7p-1}{2}} B_j) = S_1^{7p} + S_2^{7p} = 1 \quad (21)$$

$$\exp(A + 2B_p + 2B_{2p} + 2B_{3p}) = S_1^7 + S_2^7 = [\exp \sum_{\alpha=1}^{p-1} t_{7\alpha}]^7 \quad (22)$$

$$\exp(A + 2 \sum_{i=1}^{\frac{p-1}{2}} B_{7j}) = S_1^p + S_2^p = [\exp \sum_{\alpha=1}^6 t_{p\alpha}]^p \quad (23)$$

(21)-(23) can also fit in the margin.

Using this method we proved FLT in 1991 [2-5].

Let $n = p$ where p is an odd prime. From (3) and (7) we have

$$\exp(A + 2 \sum_{i=1}^{\frac{p-1}{2}} B_j) = S_1^p + S_2^p = 1, e^{2B_1} = S_1^2 + S_2^2 - 2S_1S_2 \cos \frac{\pi}{p} \quad (24)$$

Let $a = S_1 e^{-B_1}$ and $b = S_2 e^{-B_1}$ From (24) we have

$$a^p + b^p = (e^{-B_1})^p \quad (25)$$

$$a^2 + b^2 - 2ab \cos \frac{\pi}{p} = 1 \quad (26)$$

The proof of (25) is transformed into studying (26). (26) has no rational solutions with $ab \neq 0$,

because $\cos \frac{\pi}{p}$ is an irrational number for $p > 3$. Therefore (25) has no rational solutions for any odd prime $p > 3$. (25) and (26) can also fit in the margin.

Remark. If $S_i \neq 0$, where $i = 1, 2, 3, \dots, n$, then (11)-(23) have infinitely many rational solutions [1].

Preprint (January 1994).

After Wiles was about to announce his proof of FLT to the world on June 23, 1993. Jiang wrote this paper.

Tepper Gill, Kexi Liu, and Eric Trel, Editors

Fundamental Open Problems in Science at the End of the Millennium

Proceedings of the Beijing Workshop, August 1997

Hadronic Press, Palm Harbor, FL 34682-1577, U. S. A

ISBN 1-57485-029-6, pp. 555-558.

References

1. Jiang, Chun-xuan. Hypercomplex variable theory, Preprints, 1989.
2. Jiang, Chun-xuan. Fermat last theorem has been

proved (Chinese, English summary) Qian Kexue, 2(1992)17-20. Preprints (English), December,

1991. (It is sufficient to prove $S_1^3 + S_2^3 = 1$ for

- FLT of odd exponents).
3. Jiang, Chun-xuan. More than 300 years ago Fermat last theorem was proved (Chinese, English summary). Qian Kexue, 6(1992) 18-20. (It is sufficient to prove $S_1^4 - S_2^4 = 1$ for FLT.)
 4. Jiang, Chun-xuan. Fermat proof for FLT. Preprints (English), March, 1992.
 5. Jiang, Chun-xuan. Factorization theorem for Fermat equation. Preprints (English), May, 1992.

Note.

Let one knew the important results, we gave out about 600 preprints in 1991-1992. There were my preprints in Princeton, Harvard, Berkeley, MIT, Uchicago, Columbia, Maryland, Ohio, Wisconsin, Yale,, England, Canada, Japan, Poland,

Germany, France, Finland,, Ann. Math., Mathematika, J. Number Theory, Glasgow Math. J., London Math. Soc., In. J. Math. Math. Sci., Acta Arith., Can. Math. Bull. (They refused the publicaitons of my papers). Both papers were published in Chinese. FLT is as simple as Pythagorean theorem. This proof can fit in the margin of Fermat book. We think the game is up. We sent dept of math (Princeton University) a preprint on Jan. 15, 1992. Wiles claims the second proof of FLT in England (not in U. S. A.) after two years. We wish Wiles and his supporters disprove my proof, otherwise Wiles work is only the second and complex proof of FLT. We believe that the Princeton is the fairest University and history will pass the fairest judgment on proofs of FLT and other problems. We are waiting for word from the experts who are studying this paper.

12/23/2015