

Automorphic Functions And Fermat's Last Theorem (1)

Jiang, Chun-Xuan (蒋春暄)

Institute for Basic Research, Palm Harbor, FL34682-1577, USA

And: P. O. Box 3924, Beijing 100854, China (蒋春暄, 北京 3924 信箱, 100854)

jiangchunxuan@sohu.com, cxjiang@mail.bcf.net.cn, jcxuan@sina.com, Jiangchunxuan@vip.sohu.com,
jcxxxx@163.com

Abstract: In 1637 Fermat wrote: "It is impossible to separate a cube into two cubes, or a biquadrate into two biquadrates, or in general any power higher than the second into powers of like degree: I have discovered a truly marvelous proof, which this margin is too small to contain." This means: $x^n + y^n = z^n$ ($n > 2$) has no integer solutions, all different from 0 (i.e., it has only the trivial solution, where one of the integers is equal to 0). It has been called Fermat's last theorem (FLT). It suffices to prove FLT for exponent 4. and every prime exponent P . Fermat proved FLT for exponent 4. Euler proved FLT for exponent 3. In this paper using automorphic functions we prove FLT for exponents $3P$ and P , where P is an odd prime. The proof of FLT must be direct. But indirect proof of FLT is disbelieving.

[Jiang, Chun-Xuan (蒋春暄). **Automorphic Functions And Fermat's Last Theorem (1)**. *Academ Arena* 2016;8(5):78-84]. ISSN 1553-992X (print); ISSN 2158-771X (online). <http://www.sciencepub.net/academia>. 8. doi:[10.7537/marsaaj08051608](https://doi.org/10.7537/marsaaj08051608).

Keywords: Automorphic Function; Fermat's Last Theorem; Jiang Chunxuan; number

In 1974 Jiang found out Euler formula of the cyclotomic real numbers in the cyclotomic fields

$$\exp\left(\sum_{i=1}^{n-1} t_i J^i\right) = \sum_{i=1}^n S_i J^{i-1} \quad (1)$$

where J denotes a n th root of unity, $J^n = 1$, n is an odd number, t_i are the real numbers.

S_i is called the automorphic functions (complex hyperbolic functions) of order n with $n-1$ variables [1-7].

$$S_i = \frac{1}{n} \left[e^A + 2 \sum_{j=1}^{\frac{n-1}{2}} (-1)^{(i-1)j} e^{B_j} \cos\left(\theta_j + (-1)^j \frac{(i-1)j\pi}{n}\right) \right] \quad (2)$$

where $i=1, 2, \dots, n$;

$$A = \sum_{\alpha=1}^{n-1} t_\alpha, \quad B_j = \sum_{\alpha=1}^{n-1} t_\alpha (-1)^{\alpha j} \cos \frac{\alpha j \pi}{n},$$

$$\theta_j = (-1)^{j+1} \sum_{\alpha=1}^{n-1} t_\alpha (-1)^{\alpha j} \sin \frac{\alpha j \pi}{n}, \quad A + 2 \sum_{j=1}^{\frac{n-1}{2}} B_j = 0 \quad (3)$$

(2) may be written in the matrix form

$$\begin{bmatrix} S_1 \\ S_2 \\ S_3 \\ \dots \\ S_n \end{bmatrix} = \frac{1}{n} \begin{bmatrix} 1 & 1 & 0 & \dots & 0 \\ 1 & -\cos \frac{\pi}{n} & -\sin \frac{\pi}{n} & \dots & -\sin \frac{(n-1)\pi}{2n} \\ 1 & \cos \frac{2\pi}{n} & \sin \frac{2\pi}{n} & \dots & -\sin \frac{(n-1)\pi}{n} \\ \dots & \dots & \dots & \dots & \dots \\ 1 & \cos \frac{(n-1)\pi}{n} & \sin \frac{(n-1)\pi}{n} & \dots & -\sin \frac{(n-1)^2\pi}{2n} \end{bmatrix} \begin{bmatrix} e^A \\ 2e^{B_1} \cos \theta_1 \\ 2e^{B_1} \sin \theta_1 \\ \dots \\ 2 \exp B_{\frac{n-1}{2}} \sin \theta_{\frac{n-1}{2}} \end{bmatrix} \quad (4)$$

where $(n-1)/2$ is an even number.

From (4) we have its inverse transformation

$$\begin{bmatrix} e^A \\ e^{B_1} \cos \theta_1 \\ e^{B_1} \sin \theta_1 \\ \dots \\ \exp(B_{\frac{n-1}{2}}) \sin(\theta_{\frac{n-1}{2}}) \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 & \dots & 1 \\ 1 & -\cos \frac{\pi}{n} & \cos \frac{2\pi}{n} & \dots & \cos \frac{(n-1)\pi}{n} \\ 0 & -\sin \frac{\pi}{n} & \sin \frac{2\pi}{n} & \dots & \sin \frac{(n-1)\pi}{n} \\ \dots & \dots & \dots & \dots & \dots \\ 0 & -\sin \frac{(n-1)\pi}{2n} & -\sin \frac{(n-1)\pi}{n} & \dots & -\sin \frac{(n-1)^2\pi}{2n} \end{bmatrix} \begin{bmatrix} S_1 \\ S_2 \\ S_3 \\ \dots \\ S_n \end{bmatrix} \quad (5)$$

From (5) we have

$$e^A = \sum_{i=1}^n S_i, \quad e^{B_j} \cos \theta_j = S_1 + \sum_{i=1}^{n-1} S_{1+i} (-1)^{ij} \cos \frac{ij\pi}{n}$$

$$e^{B_j} \sin \theta_j = (-1)^{j+1} \sum_{i=1}^{n-1} S_{1+i} (-1)^{ij} \sin \frac{ij\pi}{n}, \quad (6)$$

In (3) and (6) t_i and S_i have the same formulas. (4) and (5) are the most critical formulas of proofs for FLT. Using (4) and (5) in 1991 Jiang invented that every factor of exponent n has the Fermat equation and proved FLT [1-7] Substituting (4) into (5) we prove (5).

$$\begin{bmatrix} e^A \\ e^{B_1} \cos \theta_1 \\ e^{B_1} \sin \theta_1 \\ \dots \\ \exp(B_{\frac{n-1}{2}}) \sin(\theta_{\frac{n-1}{2}}) \end{bmatrix} = \frac{1}{n} \begin{bmatrix} 1 & 1 & 1 & \dots & 1 \\ 1 & -\cos \frac{\pi}{n} & \cos \frac{2\pi}{n} & \dots & \cos \frac{(n-1)\pi}{n} \\ 0 & -\sin \frac{\pi}{n} & \sin \frac{2\pi}{n} & \dots & \sin \frac{(n-1)\pi}{n} \\ \dots & \dots & \dots & \dots & \dots \\ 0 & -\sin \frac{(n-1)\pi}{2n} & -\sin \frac{(n-1)\pi}{n} & \dots & -\sin \frac{(n-1)^2\pi}{2n} \end{bmatrix} \times$$

$$\begin{aligned}
 & \begin{bmatrix} 1 & 1 & 0 & \dots & 0 \\ 1 & -\cos \frac{\pi}{n} & -\sin \frac{\pi}{n} & \dots & -\sin \frac{(n-1)\pi}{2n} \\ 1 & \cos \frac{2\pi}{n} & \sin \frac{2\pi}{n} & \dots & -\sin \frac{(n-1)\pi}{n} \\ \dots & \dots & \dots & \dots & \dots \\ 1 & \cos \frac{(n-1)\pi}{n} & \sin \frac{(n-1)\pi}{n} & \dots & -\sin \frac{(n-1)^2\pi}{2n} \end{bmatrix} \begin{bmatrix} e^A \\ 2e^{B_1} \cos \theta_1 \\ 2e^{B_1} \sin \theta_1 \\ \dots \\ 2 \exp(B_{\frac{n-1}{2}}) \sin(\theta_{\frac{n-1}{2}}) \end{bmatrix} \\
 &= \frac{1}{n} \begin{bmatrix} n & 0 & 0 & \dots & 0 \\ 0 & \frac{n}{2} & 0 & \dots & 0 \\ 0 & 0 & \frac{n}{2} & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & \frac{n}{2} \end{bmatrix} \begin{bmatrix} e^A \\ 2e^{B_1} \cos \theta_1 \\ 2e^{B_1} \sin \theta_1 \\ \dots \\ 2 \exp(B_{\frac{n-1}{2}}) \sin(\theta_{\frac{n-1}{2}}) \end{bmatrix} \\
 &= \begin{bmatrix} e^A \\ e^{B_1} \cos \theta_1 \\ e^{B_1} \sin \theta_1 \\ \dots \\ \exp(B_{\frac{n-1}{2}}) \sin(\theta_{\frac{n-1}{2}}) \end{bmatrix}, \tag{7}
 \end{aligned}$$

where $1 + \sum_{j=1}^{n-1} (\cos \frac{j\pi}{n})^2 = \frac{n}{2}$, $\sum_{j=1}^{n-1} (\sin \frac{j\pi}{n})^2 = \frac{n}{2}$.

From (3) we have

$$\exp(A + 2 \sum_{j=1}^{\frac{n-1}{2}} B_j) = 1 \tag{8}$$

From (6) we have

$$\exp(A + 2 \sum_{j=1}^{\frac{n-1}{2}} B_j) = \begin{vmatrix} S_1 & S_n & \dots & S_2 \\ S_2 & S_1 & \dots & S_3 \\ \dots & \dots & \dots & \dots \\ S_n & S_{n-1} & \dots & S_1 \end{vmatrix} = \begin{vmatrix} S_1 & (S_1)_1 & \dots & (S_1)_{n-1} \\ S_2 & (S_2)_1 & \dots & (S_2)_{n-1} \\ \dots & \dots & \dots & \dots \\ S_n & (S_n)_1 & \dots & (S_n)_{n-1} \end{vmatrix}, \tag{9}$$

$$(S_i)_j = \frac{\partial S_i}{\partial t_j} \tag{7}$$

where
 From (8) and (9) we have the circulant determinant

$$\exp\left(A + 2 \sum_{j=1}^{\frac{n-1}{2}} B_j\right) = \begin{vmatrix} S_1 & S_n & \cdots & S_2 \\ S_2 & S_1 & \cdots & S_3 \\ \cdots & \cdots & \cdots & \vdots \\ S_n & S_{n-1} & \cdots & S_1 \end{vmatrix} = 1 \quad (10)$$

If $S_i \neq 0$, where $i = 1, 2, \dots, n$, then (10) has infinitely many rational solutions.

Assume $S_1 \neq 0$, $S_2 \neq 0$, $S_i = 0$ where $i = 3, 4, \dots, n$. $S_i = 0$ are $n-2$ indeterminate equations with $n-1$ variables. From (6) we have

$$e^A = S_1 + S_2, \quad e^{2B_j} = S_1^2 + S_2^2 + 2S_1S_2(-1)^j \cos \frac{j\pi}{n} \quad (11)$$

From (10) and (11) we have the Fermat equation

$$\exp\left(A + 2 \sum_{j=1}^{\frac{n-1}{2}} B_j\right) = (S_1 + S_2) \prod_{j=1}^{\frac{n-1}{2}} (S_1^2 + S_2^2 + 2S_1S_2(-1)^j \cos \frac{j\pi}{n}) = S_1^n + S_2^n = 1 \quad (12)$$

Example[1]. Let $n = 15$. From (3) we have

$$A = (t_1 + t_{14}) + (t_2 + t_{13}) + (t_3 + t_{12}) + (t_4 + t_{11}) + (t_5 + t_{10}) + (t_6 + t_9) + (t_7 + t_8)$$

$$B_1 = -(t_1 + t_{14}) \cos \frac{\pi}{15} + (t_2 + t_{13}) \cos \frac{2\pi}{15} - (t_3 + t_{12}) \cos \frac{3\pi}{15} + (t_4 + t_{11}) \cos \frac{4\pi}{15} \\ - (t_5 + t_{10}) \cos \frac{5\pi}{15} + (t_6 + t_9) \cos \frac{6\pi}{15} - (t_7 + t_8) \cos \frac{7\pi}{15},$$

$$B_2 = (t_1 + t_{14}) \cos \frac{2\pi}{15} + (t_2 + t_{13}) \cos \frac{4\pi}{15} + (t_3 + t_{12}) \cos \frac{6\pi}{15} + (t_4 + t_{11}) \cos \frac{8\pi}{15} \\ + (t_5 + t_{10}) \cos \frac{10\pi}{15} + (t_6 + t_9) \cos \frac{12\pi}{15} + (t_7 + t_8) \cos \frac{14\pi}{15},$$

$$B_3 = -(t_1 + t_{14}) \cos \frac{3\pi}{15} + (t_2 + t_{13}) \cos \frac{6\pi}{15} - (t_3 + t_{12}) \cos \frac{9\pi}{15} + (t_4 + t_{11}) \cos \frac{12\pi}{15} \\ - (t_5 + t_{10}) \cos \frac{15\pi}{15} + (t_6 + t_9) \cos \frac{18\pi}{15} - (t_7 + t_8) \cos \frac{21\pi}{15},$$

$$B_4 = (t_1 + t_{14}) \cos \frac{4\pi}{15} + (t_2 + t_{13}) \cos \frac{8\pi}{15} + (t_3 + t_{12}) \cos \frac{12\pi}{15} + (t_4 + t_{11}) \cos \frac{16\pi}{15} \\ + (t_5 + t_{10}) \cos \frac{20\pi}{15} + (t_6 + t_9) \cos \frac{24\pi}{15} + (t_7 + t_8) \cos \frac{28\pi}{15},$$

$$B_5 = -(t_1 + t_{14}) \cos \frac{5\pi}{15} + (t_2 + t_{13}) \cos \frac{10\pi}{15} - (t_3 + t_{12}) \cos \frac{15\pi}{15} + (t_4 + t_{11}) \cos \frac{20\pi}{15} \\ - (t_5 + t_{10}) \cos \frac{25\pi}{15} + (t_6 + t_9) \cos \frac{30\pi}{15} - (t_7 + t_8) \cos \frac{35\pi}{15},$$

$$B_6 = (t_1 + t_{14}) \cos \frac{6\pi}{15} + (t_2 + t_{13}) \cos \frac{12\pi}{15} + (t_3 + t_{12}) \cos \frac{18\pi}{15} + (t_4 + t_{11}) \cos \frac{24\pi}{15}$$

$$\begin{aligned}
 &+(t_5 + t_{10}) \cos \frac{30\pi}{15} + (t_6 + t_9) \cos \frac{36\pi}{15} + (t_7 + t_8) \cos \frac{42\pi}{15} , \\
 B_7 = &-(t_1 + t_{14}) \cos \frac{7\pi}{15} + (t_2 + t_{13}) \cos \frac{14\pi}{15} - (t_3 + t_{12}) \cos \frac{21\pi}{15} + (t_4 + t_{11}) \cos \frac{28\pi}{15} \\
 &-(t_5 + t_{10}) \cos \frac{35\pi}{15} + (t_6 + t_9) \cos \frac{42\pi}{15} - (t_7 + t_8) \cos \frac{49\pi}{15} ,
 \end{aligned}$$

$$A + 2 \sum_{j=1}^7 B_j = 0, \quad A + 2B_3 + 2B_6 = 5(t_5 + t_{10}) \tag{13}$$

Form (12) we have the Fermat equation

$$\exp(A + 2 \sum_{j=1}^7 B_j) = S_1^{15} + S_2^{15} = (S_1^5)^3 + (S_2^5)^3 = 1 \tag{14}$$

From (13) we have

$$\exp(A + 2B_3 + 2B_6) = [\exp(t_5 + t_{10})]^5 \tag{15}$$

From (11) we have

$$\exp(A + 2B_3 + 2B_6) = S_1^5 + S_2^5 \tag{16}$$

From (15) and (16) we have the Fermat equation

$$\exp(A + 2B_3 + 2B_6) = S_1^5 + S_2^5 = [\exp(t_5 + t_{10})]^5 \tag{17}$$

Euler proved that (14) has no rational solutions for exponent 3[8]. Therefore we prove that (17) has no rational solutions for exponent 5[1].

Theorem 1. [1-7]. Let $n = 3P$, where $P > 3$ is odd prime. From (12) we have the Fermat's equation

$$\exp(A + 2 \sum_{j=1}^{\frac{3P-1}{2}} B_j) = S_1^{3P} + S_2^{3P} = (S_1^P)^3 + (S_2^P)^3 = 1 \tag{18}$$

From (3) we have

$$\exp(A + 2 \sum_{j=1}^{\frac{P-1}{2}} B_{3j}) = [\exp(t_P + t_{2P})]^P \tag{19}$$

From (11) we have

$$\exp(A + 2 \sum_{j=1}^{\frac{P-1}{2}} B_{3j}) = S_1^P + S_2^P \tag{20}$$

From (19) and (20) we have the Fermat equation

$$\exp(A + 2 \sum_{j=1}^{\frac{P-1}{2}} B_{3j}) = S_1^P + S_2^P = [\exp(t_P + t_{2P})]^P \tag{21}$$

Euler proved that (18) has no rational solutions for exponent 3[8]. Therefore we prove that (21) has no rational solutions for $P > 3$ [1, 3-7].

Theorem 2. In 1847 Kummer write the Fermat's equation

$$x^P + y^P = z^P \tag{22}$$

in the form

$$(x + y)(x + ry)(x + r^2y) \cdots (x + r^{P-1}y) = z^P \tag{23}$$

$$r = \cos \frac{2\pi}{P} + i \sin \frac{2\pi}{P}$$

where P is odd prime,

[8]. Kummer assume the divisor of each factor is a P th power. Kummer proved FLT for prime exponent $p < 100$

We consider the Fermat's equation

$$x^{3P} + y^{3P} = z^{3P} \quad (24)$$

we rewrite (24)

$$(x^P)^3 + (y^P)^3 = (z^P)^3 \quad (25)$$

From (24) we have

$$(x^P + y^P)(x^P + ry^P)(x^P + r^2y^P) = z^{3P} \quad (26)$$

$$r = \cos \frac{2\pi}{3} + i \sin \frac{2\pi}{3}$$

where

We assume the divisor of each factor is a P th power.

$$\text{Let } S_1 = \frac{x}{z}, S_2 = \frac{y}{z}. \text{ From (20) and (26) we have the Fermat's equation}$$

$$x^P + y^P = [z \times \exp(t_p + t_{2p})]^P \quad (27)$$

Euler proved that (25) has no integer solutions for exponent 3 [8]. Therefore we prove that (27) has no integer solutions for prime exponent P .

Fermat Theorem. It suffices to prove FLT for exponent 4. We rewrite (24)

$$(x^3)^P + (y^3)^P = (z^3)^P \quad (28)$$

Euler proved that (25) has no integer solutions for exponent 3 [8]. Therefore we prove that (28) has no integer solutions for all prime exponent P [1-7].

We consider Fermat equation

$$x^{4P} + y^{4P} = z^{4P} \quad (29)$$

We rewrite (29)

$$(x^P)^4 + ((y^P)^4 = (z^P)^4 \quad (30)$$

$$(x^4)^P + (y^4)^P = (z^4)^P \quad (31)$$

Fermat proved that (30) has no integer solutions for exponent 4 [8]. Therefore we prove that (31) has no integer solutions for all prime exponent P [2,5,7]. This is the proof that Fermat thought to have had.

Remark. It suffices to prove FLT for exponent 4. Let $n = 4P$, where P is an odd prime. We have the Fermat's equation for exponent $4P$ and the Fermat's equation for exponent P [2,5,7]. This is the proof that Fermat thought to have had. In complex hyperbolic functions let exponent n be $n = \Pi P$, $n = 2\Pi P$ and $n = 4\Pi P$. Every factor of exponent n has the Fermat's equation [1-7]. In complex trigonometric functions let exponent n be $n = \Pi P$, $n = 2\Pi P$

and $n = 4\Pi P$. Every factor of exponent n has Fermat's equation [1-7]. Using modular elliptic curves Wiles and Taylor prove FLT [9,10]. This is not the proof that Fermat thought to have had. The classical theory of automorphic functions, created by Klein and Poincare, was concerned with the study of analytic functions in the unit circle that are invariant under a discrete group of transformations. Automorphic functions are generalization of the trigonometric, hyperbolic, elliptic, and certain other functions of elementary analysis. The complex trigonometric functions and complex hyperbolic functions have a wide application in mathematics and physics.

Acknowledgments:

We thank Chenny and Moshe Klein for their help and suggestion.

References

1. Jiang, C-X, Fermat last theorem had been proved, Potential Science (in Chinese), 2.17-20 (1992), Preprints (in English) December (1991). <http://www.wbabin.net/math/xuan47.pdf>.
2. Jiang, C-X, Fermat last theorem had been proved by Fermat more than 300 years ago, Potential Science (in Chinese), 6.18-20(1992).
3. Jiang, C-X, On the factorization theorem of circulant determinant, Algebras, Groups and Geometries, 11. 371-377(1994), MR. 96a: 11023, <http://www.wbabin.net/math/xuan45.pdf>.
4. Jiang, C-X, Fermat last theorem was proved in 1991, Preprints (1993). In: Fundamental open problems in science at the end of the millennium, T. Gill, K. Liu and E. Trelle (eds). Hadronic Press, 1999, P555-558.
5. Jiang, C-X, On the Fermat-Santilli theorem, Algebras, Groups and Geometries, 15. 319-349(1998). <http://www.wbabin.net/math/xuan46.pdf>.
6. Jiang, C-X, Complex hyperbolic functions and Fermat's last theorem, Hadronic Journal Supplement, 15. 341-348(2000).
7. Jiang, C-X, Foundations of Santilli Isonumber Theory with applications to new cryptograms, Fermat's theorem and Goldbach's Conjecture. Inter, Acad. Press. 2002. MR2004c:11001, <http://www.wbabin.net/math/xuan13.pdf>. <http://www.i-b-r.org/docs/jjiang.pdf>.
8. Ribenboim, P, Fermat last theorem for amateur, Springer-Verlag, (1999).
9. Wiles, A, Modular elliptic curves and Fermat last theorem, Ann. of Math.,(2) 141(1995),443-551.
10. Taylor, R. and Wiles, A., Ring-theoretic properties of certain Hecke algebras, Ann. of Math (2),141(1995),553-572.

5/25/2016