

Santilli-Jiang Isomathematics for Changing Modern Mathematics.

Jiang, Chun-Xuan (蒋春暄)

Institute for Basic Research, Palm Harbor, FL34682-1577, USA

And: P. O. Box 3924, Beijing 100854, China (蒋春暄, 北京 3924 信箱, 100854)

jiangchunxuan@sohu.com, cxjiang@mail.bcf.net.cn, jcxuan@sina.com, Jiangchunxuan@vip.sohu.com,
jcxxxx@163.com

Abstract: The establishment of isomathematics, as proposed by Santilli thirty years ago in the USA, and contributed to by Jiang Chun-Xuan in China during the past 12 years, is significant and has changed the modern mathematics. At present, primary school mathematics consists of simple addition, subtraction, multiplication and division; middle school mathematics takes these four operations to a higher level; while university mathematics extends them to an even higher level. These four arithmetic operations form the foundation of modern mathematics.

[Jiang, Chun-Xuan (蒋春暄). **Santilli-Jiang Isomathematics for Changing Modern Mathematics.** *Academ Arena* 2016;8(6):40-43]. ISSN 1553-992X (print); ISSN 2158-771X (online). <http://www.sciencepub.net/academia>. 8. doi:[10.7537/marsaaj080616.08](https://doi.org/10.7537/marsaaj080616.08).

Keywords: Santilli-Jiang Isomathematics; Modern Mathematics; Jiang Chun-Xuan

Isomathematics is a generalisation of these four fundamental operations and heralds a great revolution in mathematics. Here, the four arithmetic operations of isoaddition, isosubtraction, isomultiplication and isodivision are established but only at the primary level of isomathematics. The material introduced here should be readily understandable by middle school pupils and university students.

Santilli's isomathematics is based on a generalisation of modern mathematics. Isomultiplication is defined by

$a \hat{\times} b = ab\hat{I}$, isodivision by $a \hat{\div} b = \frac{a}{b}\hat{I}$, where $\hat{I} \neq 1$ is called an isounit; $\hat{T}\hat{I} = 1$, where \hat{T} is the inverse of the isounit. If addition and subtraction remain unchanged, $(+, -, \hat{\times}, \hat{\div})$ are the four arithmetic operations in Santilli's isomathematics[1-5]. Isoaddition $a \hat{+} b = a + b + \hat{O}$ and isosubtraction $a \hat{-} b = a - b - \hat{O}$, where $\hat{O} \neq 0$ is called the isozero, together with the operations of isomultiplication and isodivision introduced above, form the four arithmetic operations $(\hat{+}, \hat{-}, \hat{\times}, \hat{\div})$ in Santilli-Jiang isomathematics[6].

Santilli [1] suggests isomathematics based on a generalisation of multiplication \times , division \div , and the multiplicative unit 1 of modern mathematics. It is an epoch making suggestion. From modern mathematics, the foundations of Santilli's isomathematics will be established.

(1) Division and multiplication in modern mathematics.

Suppose that

$$a \div a = a^0 = 1 \tag{1}$$

where 1 is the multiplicative unit and 0 is exponent zero.

From (1), division \div and multiplication \times are defined by

$$a \div b = \frac{a}{b}, b \neq 0, a \times b = ab \tag{2}$$

$$a = a \times (a \div a) = a \times a^0 = a \tag{3}$$

Now consider the multiplicative unit 1,

$$a \times 1 = a, a \div 1 = a, 1 \div a = 1/a \tag{4}$$

$$(1)^n = 1, (1)^{a/b} = 1, (-1)^n = (-1)^n, (-1)^{a/b} = (-1)^{a/b} \tag{5}$$

Addition +, subtraction -, multiplication \times , and division \div are the four operations forming the foundation of modern mathematics. The modern mathematics is generalised to establish the foundations of Santilli-Jiang isomathematics.

(2) Isodivision and isomultiplication in Santilli's isomathematics.

Isodivision $\hat{\div}$ and isomultiplication $\hat{\times}$ [1 – 5], which are generalisations of the division \div and multiplication \times of modern mathematics, are now defined.

$$a \hat{\div} a = a^{\bar{0}} = \hat{I} \neq 1, \quad \bar{0} \neq 0 \quad (6)$$

where \hat{I} is called the isounit and is a generalisation of the multiplicative unit 1 and $\bar{0}$ is the isoexponent zero which is a generalisation of the exponent zero 0. Then,

$$a \hat{\div} b = \hat{I} \frac{a}{b}, \quad b \neq 0, \quad a \hat{\times} b = a \hat{T} b \quad (7)$$

It is seen that

$$a = a \hat{\times} (a \hat{\div} a) = a \hat{\times} a^{\bar{0}} = a \hat{T} \hat{I} = a \quad (8)$$

from which it follows that

$$\hat{T} \hat{I} = 1 \quad (9)$$

where \hat{T} is the inverse of the isounit \hat{I} .

The isounit \hat{I} has the following properties[5,p93-95,isoexponents]:

$$a \hat{\times} \hat{I} = a, \quad a \hat{\div} \hat{I} = a, \quad \hat{I} \hat{\div} a = a^{-\hat{I}} = \hat{I}^2 / a \quad (10)$$

$$(+\hat{I})^{\hat{n}} = \hat{I}, \quad (+\hat{I})^{\hat{a}} = \hat{I}, \quad (-\hat{I})^{\hat{n}} = (-1)^{\hat{n}} \hat{I}, \quad (-\hat{I})^{\hat{a}} = (-1)^{\hat{a}} \hat{I} \quad (11)$$

With addition and subtraction unchanged, $(+, -, \hat{\times}, \hat{\div})$ are the four arithmetic operations in Santilli's isomathematics and these form the foundations of Santilli isomathematics. **When $\hat{I} = 1$, the operations revert to being those of the modern mathematics.**

(3) Addition and subtraction in modern mathematics.

These are defined by

$$x = a + b \quad \text{and} \quad y = a - b \quad (12)$$

$$a + a - a = a \quad (13)$$

$$a - a = 0 \quad (14)$$

Isoaddition and isosubtraction may be established using these results.

(4) Isoaddition and isosubtraction in Santilli-Jiang isomathematics.

Isoaddition $\hat{+}$ and isosubtraction $\hat{-}$ are defined by

$$a \hat{+} b = a + b + c_1, \quad a \hat{-} b = a - b - c_2 \quad (15)$$

$$\therefore a = a \hat{+} a \hat{-} a = a + c_1 - c_2 = a \quad (16)$$

Then, from (16), it follows that

$$c_1 = c_2 \quad (17)$$

Suppose that $c_1 = c_2 = \hat{0}$, where $\hat{0}$ is called the isozero which is a generalisation of the zero 0 of addition and subtraction[6]. Hence,

$$a \hat{+} b = a + b + \hat{0}, \quad a \hat{-} b = a - b - \hat{0} \quad (18)$$

When $\hat{0} = 0$, these equations are the usual laws of addition and subtraction of modern mathematics.

From the above results, the foundations of Santilli-Jiang isomathematics are readily established:

$$\hat{\times} = \times \hat{T} \times, \hat{+} = + \hat{0} +; \hat{\div} = \div \hat{I} \div, \hat{-} = - \hat{0} -; a \hat{\times} b = ab \hat{T}, a \hat{+} b = a + b + \hat{0};$$

$$a \hat{\div} b = \frac{a}{b} \hat{I}, a \hat{-} b = a - b - \hat{0}; a = a \hat{\times} a \hat{\div} a = a, a = a \hat{+} a \hat{-} a = a;$$

$$a \hat{\times} a = a^2 \hat{T}, a \hat{+} a = 2a + \hat{0}; a \hat{\div} a = \hat{I} \neq 1, a \hat{-} a = -\hat{0} \neq 0; \hat{T} \hat{I} = 1. \quad (19)$$

Here $(\hat{+}, \hat{-}, \hat{\times}, \hat{\div})$ are the four arithmetic operations in Santilli-Jiang isomathematics.

Remark:

$$a \hat{\times} (b \hat{+} c) = a \hat{\times} (b + c + \hat{0})$$

From the left-hand side, it follows

$$a \hat{\times} (b \hat{+} c) = a \hat{\times} b + a \hat{\times} \hat{+} + a \hat{\times} c = a \hat{\times} (b + \hat{+} + c) = a \hat{\times} (b + \hat{0} + c),$$

where $\hat{+} = \hat{0}$ is a number also.

Again,

$$a \hat{\times} (b \hat{-} c) = a \hat{\times} (b - c - \hat{0})$$

From the left-hand side of this relation, it is seen that

$$a \hat{\times} (b \hat{-} c) = a \hat{\times} b - a \hat{\times} \hat{-} - a \hat{\times} c = a \hat{\times} (b - \hat{-} - c) = a \hat{\times} (b - \hat{0} - c),$$

where $\hat{-} = \hat{0}$ is a number also.

The distributive laws are satisfied and $\hat{+}, \hat{-}, \hat{\times}, \hat{\div}$ are numbers.

This Santilli-Jiang isomathematics therefore, provides a new mathematical tool for studying the mathematical problems of the 21st century and helping in the understanding the mysteries of our universe.

(5) An illustrative example.

Consider the algebraic equation

$$y = a_1 \times (b_1 + c_1) + a_2 \div (b_2 - c_2) \quad (20)$$

(20) may represent a mathematical system, physical system, biological system, cryptogram system, IT system, or some other system. It may be written as the isomathematical equation

$$\hat{y} = a_1 \hat{\times} (b_1 \hat{+} c_1) \hat{+} a_2 \hat{\div} (b_2 \hat{-} c_2) = a_1 \hat{T} (b_1 + c_1 + \hat{0}) + \hat{0} + a_2 / \hat{T} (b_2 - c_2 - \hat{0}) \quad (21)$$

If $\hat{T} = 1$ and $\hat{0} = 0$ then $y = \hat{y}$.

Let $\hat{T} = 2$ and $\hat{0} = 3$. From (21) we have the isomathematical subequation

$$\hat{y}_1 = 2a_1 (b_1 + c_1 + 3) + 3 + a_2 / 2 (b_2 - c_2 - 3) \quad (22)$$

Let $\hat{T} = 5$ and $\hat{0} = 6$. From (21) we have the isomathematical subequation

$$\hat{y}_2 = 5a_1 (b_1 + c_1 + 6) + 6 + a_2 / 5 (b_2 - c_2 - 6) \quad (23)$$

Let $\hat{T} = 8$ and $\hat{0} = 10$. From (21) we have the isomathematical subequation

$$\hat{y}_3 = 8a_1 (b_1 + c_1 + 10) + 10 + a_2 / 8 (b_2 - c_2 - 10) \quad (24)$$

Therefore, (21) has infinitely many isomathematical subequations. Using (21) – (24), \hat{T} and $\hat{0}$, the stability and optimum structure of the algebraic equation (20) may be studied.

References.

1. R. M. Santilli, (1993), *Isonumbers and genonumbers of dimension 1, 2, 4, 8, their isoduals and pseudoduals, and "hidden numbers" of dimension 3, 5, 6, 7*; *Algebras, Groups and Geometries* 10, 273-322.
2. Chun-Xuan Jiang, (1998), *Foundations of Santilli's isonumber theory, Part I: Isonumber theory of the first kind*; *Algebras, Groups and Geometries*, 15, 351-393.
3. Chun-Xuan Jiang, (1998), *Foundations of Santilli's isonumber theory, Part II: Isonumber theory of the second kind*; *Algebras Groups and Geometries*, 15, 509-544.
4. Chun-Xuan Jiang, (1999), *Foundations of Santilli's isonumber theory*, in: *Fundamental open problems in sciences at the end of the millennium*, T. Gill, K. Liu and E. Trelle (Eds), Hadronic Press, USA, 105-139.
5. Chun-Xuan Jiang, (2002), *Foundations of Santilli's isonumber theory, with applications to new cryptograms, Fermat's theorem and Goldbach's conjecture*, International Academic Press, America- Europe- Asia.MR2004c:11001. (also available in the pdf file <http://www.i-b-r.org/docs/jiang.pdf>) (<http://vixra.org/pdf/1303.0088v1.pdf>).
6. Ma H. The Nature of Time and Space. *Nature and science* 2003;1(1):1-11. *Nature and science* 2007;5(1):81-96.
7. Chun-Xuan Jiang, (2008), Santilli-Jiang isomathematics for changing modern mathematics, middle school mathematics (Chinese), Dec. 46-48.

6/22/2016