

Genetic Diversity of *Shistosoma mansoni* Isolates Genome and Protein Analysis

Mohammed Yahia Araishi

Collage of health Science, Jazan Univ., Saudi Arabia.

Abstract: Five of *Shistosoma mansoni* isolates which differ in disease potential show genetic variability. Esterase isozymes, protein pattern and DNA finger-print analysis successfully revealed genetic diversity. DISC PAGE esterase isozymes revealed 5 bands which differ in density and relative mobility among SM1, SM3, SM4, SM5, while 4 bands with SM2. SDS-PAGE of protein finger print varied among the five isolates in number of protein species 19.18, 17.16 and 16 bands of SM1, SM2, SM3, SM4 and SM5 isolates respectively as well as in intensity. Molecular weight and reproducibility polymorphism analysis presented 12 common fragments (monomorphic) with 54%, 8 specific bands (polymorphic) with 45% and 4 bands unique with 20%. RAPD-PCR for identification of the 5 *S. mansoni* isolates were conducted on the screening of primers and on the extraction of template DNA. Three out of 10 sequence 10 mer primers were successful in identification of the 5 *S. mansoni* isolates. PCR amplification yielded reproducible RAPD patterns which differentiated the 5 isolates examined. The difference also recognized the RAPD patterns specially of SM5. Esterase isozymes, protein and DNA fragments conformed the genetic diversity among 5 *S. mansoni* isolates. [Journal of American Science 2010;6(8):104-110]. (ISSN: 1545-1003).

Key words: *S. mansoni*, esterase isozymes DISC-PAGE, RAPD-PCR, protein and DNA fragments.

1. Introduction

Schistosoma mansoni is one of the most severe tropical diseases in the world. Egypt is one of the most highly endemic areas in some localities in the Nile Valley. *Biomphalaria alexandrina* snail as specific intermediate host of *S. mansoni* are prevalent in both upper and lower Egypt, but during the last decade, it became the most dominant species in the Nile Delta forming a main threat for Schistosomiasis transmission in the North of Egypt. This wide distribution of the disease makes the study of genomic variability extremely important. There are well known two distinct geographical strains of *S. mansoni*, the Puerto Rican and the Egyptian strains (Saoud, 1966). However a new geographical strain namely, Saudi Arabian strain has been reported based on differences with the Egyptian strain shown by scanning electron microscopy (Shalaby et al., 1993).

Several techniques of molecular biology for studies S1, S2, S3, S4, S5, of genetic variability in Schistosomiasis vector have been used (Da Silva et al., 2004). An introduction of the random amplified polymorphic DNA-polymerase chain reaction (RAPD-PCR) technique (Jamjoom, 2006) has amplified possibilities of polymorphism analysis and biochemical markers, as enzyme isozymes (Lockyer et al., 2000).

The present study emphasizes our interest in soil genetics to detect genetic variability among five *S. mansoni* using RAPD-PCR and isozymes for genetic analysis of DNA genome of *S. mansoni* snails and protein enzymes.

2. Material and Methods

Parasite isolates: Five adult worms isolates of repeated twice *S. mansoni* from (Southern region). (Delta region) and (east region) at Egypt. They were obtained from theodor Bilharz Research Institute and originally selected on the basis of healthy different size and pathogenicity according to Longand and Marand (1995).

SDS-PAGE electrophoresis: Protein content of adult worms was determined by Lowry's method (Lowry et al., 1951) using bovine serum albumin as a standard.

Protein pattern was determined by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) as described by Laemmli (1970) under reducing condition using 12% pore gel with 3.7% stacking gel in a discontinuous electrode buffer system. Protein samples and molecular weight markers (Sigma) were denatured and reduced completely before electrophoresis by mixing the samples with an equal volume of 2x sample buffer and heating the mixture at 95°C in a water bath for 5 min. Treated samples were centrifuged at 12000 rpm for 10 min and chilled on ice before use. Each lane was charged with 25 µg protein of supernatant. For subsequent fractions, the lanes were filled up to their full capacity (50 µl). Electrophoresis was carried out a room temperature at a constant current of 25 mA for 1 hr followed by 30 Am for 4 hr. At the end of the run, gel was stained with commassie blue and destained in the stain solvent and photographed.

Isozyme electrophoresis:

The adult worms were homogenized in Tris. Glycine buffer and centrifuged at 10000 rpm for 10

min. The supernatant was kept at -20°C until electrophoresis according to Jonathan and Wendell, 1990). Isozyme variability was assayed by nature polyacrylamide slab gel electrophoresis using 8% acrylamide. Each sample was mixed with bromophenol blue and applied to each well, the wells were stained with the solution contain Na-phosphate and α , β naphthyl acetate and fast blue CR salt. The gel was inoculated at 4°C in work and adding the appropriate substrate and staining solution (Jonathan and Wendell, 1990).

DNA isolation:

DNA was extracted from individual frozen adult worms and homogenized with a pestle on Eppendorf tube containing 100 μl of 10 mM Tris-HCl pH 8.0; 1 mM EDTA, 10 mM NaCl and 70 mM sucrose. Sodium dodecyl sulfate (10%) and (12 μl) proteinase 10 mg ml⁻¹ were added to Eppendorf tube. The tubes were incubated at 57°C for 2 h. Then extracted once with an equal volume of phenol and once with chloroform. The DNA was precipitated by 0.3 M sodium acetate; 2 volume of absolute ethanol at 20°C overnight. The DNA was pelleted by centrifugation (15000 g/20 min at 4°C) then rinsed with ethanol 10% dried and finally resuspended in 100 μl of TE buffer (10 mM Tris HCl pH 8.0 and 1 mM EDTA). The purified DNA concentration was determined spectrophotometrically at 260 nm and 280 nm and also by using 2% agarose gel electrophoresis.

RAPD-PCR analysis:

Three arbitrary primers, A7 (5'-gaaocaaatg-3') A8 (5'-gtgacgtagg-3') and A6 (5'-gtgcaagtct-3') were used as described by Simpson et al. (1993) with modifications: 2 ng of DNA genome were amplified PCR thermal cycler). Each reaction was carried out in a final vol. Of 10 μl containing 1 unit Taq DNA polymerase 1x PCR buffer, 0.2 mM each dNTP and 7 p mol of each random primer (Gibco). The amplification conditions were as follows: 1 cycle at 95°C for 5 min, 3 steps, at 95°C for 30 sec, 30°C for 2 min and 72°C for 1 min and 72°C for 1 min and 33 cycles during which the annealing temperature was changed to 40°C and the time of the extension step was increased to 5 min during the final cycle.

The PCR products were visualized on 1.5% agarose gels and stained with ethidium bromide to resolve amplified fragments and photographed.

3. Results

Somaclonal variations among 5 isolates of *S. mansoni* were detected by protein pattern, esterase activity, isozymes and DNA fingerprint.

Protein content was determined in adult worm of each isolate related to BSA (Table 1). It was revealed that the protein content was varied among 5 isolates with 480, 425, 392, 405 and 385 mg/g fresh weight of adult worms.

Esterase activity was assayed of each isolate. The specific activities of enzyme was 4.25; 4.75; 3.82; 5.12 and 3.25 μmg protein for SM1; SM2; SM3; SM4 and SM5 respectively (Table 1). Esterase isozymes are shown in Fig. (1.A) and calculated in Table (2). The total number of isozymes 8 bands differ in number and density among five isolate such as 4 band of SM1 isolate and 5 bands of each SM2, SM3, SM4 and SM5 out of 8 total isozymes bands. The esterase isozyme variability among isolates showed some isozymes band disappeared in SM1 (0.25, 0.35, 0.55, 0.75 Rf); SM3 (0.25 M, 0.35, 0.75 Rf); SM3 (0.25, 0.55, 0.75 SM4 (0.35, 0.55, 0.75 Rf) and SM5 (0.25, 0.35, 0.55 Rf). As well as one unique band (genetic marker) at Rf 0.25 (SM4), 0.35 (SM3); 0.55 (SM2) and 0.75 (SM5) isolates and monomorphic (common isozyme band) in five isolates at Rf 0.42, 0.45, 0.61 and 0.64.

SDS-PAGE protein fractions was illustrated in Fig. (1-B) and Table (3). The results showed the variation of *S. mansoni* isolates in protein fractions and density bands. The variability analysis among isolates appeared 24 protein fractions. Out of them 19, 18, 17 polypeptides appeared in SM1, SM2, SM3 isolates respectively and 16 polypeptides in both SM4 and SM5. Some polypeptides bands disappeared among isolates (Table 3). The molecular weight of polypeptides were determined related to protein marker (Table 3). The most prominent alteration (polymorphic bands) among 5 isolates 122, 108, 60, 20 KDa with percentage 16%. The prominent polypeptide bands in all isolates (monomorphic or common fragments) were 156, 110, 100, 97, 85, 70, 54, 44, 36, 33, 24, 18 and 10 KDa with percentage 54%. The unique fragment (genetic markers) with 73, 39 (SM1); 15 (SM2); 65 (SM3), 175 (SM4) and 28, 26 (SM5) with percentage 30%.

Table (1):Protein content and esterase activity of five *Schistosoma mansoni* isolates.

<i>S. mansoni</i> isolates	Esterase activity		
	Protein content (mg/g)	Total activity (μ)	Specific activity (μmg)
SM1	480	2040.00	4.25
SM2	425	2018.75	4.75
SM3	392	1497.44	3.82
SM4	405	2673.60	5.12
SM5	385	1251.25	3.25

Table (2): Esterases isozymes analysis of the *Schistosoma monsoni* five isolates.

Schistosoma monsoni isolates							
No.	Rf bands	SM1	SM2	SM3	SM4	SM5	Polymorphism
1	0.25	-	-	-	+	-	Unique
2	0.35	-	-	+	-	-	Unique
3	0.42	++	++	++	++	++	Monomorphic
4	0.45	+++	+++	+++	+++	+++	Monomorphic
5	0.55	-	++	-	-	-	Unique
6	0.61	++++	++++	++++	++++	++++	Monomorphic
7	0.64	+++	+++	+++	+++	+++	Monomorphic
8	0.75	-	-	-	-	+	Unique
Total		4	5	5	5	5	

Rf = Relative mobility Unique = Genetic marker Monomorphic or common isozyme band
 Band density ++++ very strong; +++ Strong, ++ Moderate + Weak and - Absent band




Fig. (1): Electrogram protein and isozymes profiles of five *S. mansoni* isolate.

1-A: SDS-PAGE (12%) of denatured protein extracted from 5 isolates.

1-B: DISC-PAGE (12%) of native protein esterase isozymes.

M: Markers protein (KDa), S1, S2, S3, S4 and S5: *S. mansoni* isolates.

Table (3): Protein fractions of *Schistosoma monsoni* five isolates by SDS-PAGE.

Schistosoma monsoni isolates						
Molecular weight (KDa)	S1	S2	S3	S4	S5	Polymorphism
175	-	-	-	+	-	Unique
156	+++	+++	++++	+++	++	Monomorphic
122	++	++	++	+	-	Polymorphic
110	++	++	++	++	+	Monomorphic
108	+	+	+	-	-	Polymorphic
100	++	++	++	++	+	Monomorphic
97	+++	+++	+++	++	++	Monomorphic
85	+++	++++	+++	+++	+++	Monomorphic

73	+	-	-	-	-	Unique
70	+++	+++	++	++	++	Monomorphic
65	-	-	+	-	-	Unique
60	+	+	+	+	-	Polymorphic
54	++	++	++	++	++	Monomorphic
44	+++	+++	+++	+++	++	Monomorphic
39	+	-	-	-	-	Unique
36	++	++	++	+	++	Monomorphic
33	+++	+++	+++	+++	+++	Monomorphic
28	-	-	-	-	+	Unique
26	-	-	-	-	+	Unique
24	++++	+++	+++	+++	++++	Monomorphic
20	+	+	-	-	++	Polymorphic
18	++	++	+	+	+++	Monomorphic
15	-	+	-	-	-	Unique
10	++++	+++	++	++	++++	Monomorphic
Number protein	19	18	17	16	16	

DNA fingerprint: Total DNA preparation as found crucial for RAPD-PCR. The DNA yield was determined spectrophotometrically as 8 µg/ 0.15 g tissues. The DNA purity as indicated by 260/280 was 1.8. The reproducibility of RAPD analysis is known to be singly influenced by experimental conditions. It is therefore essential to optimize the PCR conditions to obtain reproducible and interpretable results before going on routine analysis. It was found that DNA quality was a good template per PCR amplification. However, treatments of DNA with RNase gave sharp and clear amplification products compared with untreated DNA.

The PCR reaction conditions, polymorphism among the five *S. mansoni* isolates were detected using different random primers by RAPD-PCR gave the best results of amplification expressed on average number of bands per primer. Of the three primers (Operon random primer) were screened in RAPD analysis for their ability to produce sufficient amplification products. Three random primer namely

these primers are summarized in Table (4, 5) and Fig. (2). The RAPD analysis of DNA extracted from 5 *S. mansoni* isolates revealed 22 amplified bands, (8, 7 and 7 bands) with different molecular weight ranged from 1.450 to 205 bp of primers A8, A7 and A6 respectively. The DNA amplified fragments of 5 isolates were varied in number, density and molecular weight. The variability analysis among 5 isolates showed some DNA amplified fragments absent or/and in some isolates (Table 3). The polymorphism analysis among isolates, 11 monomorphic amplified bands (common in all isolates) with 50%; 4 polymorphic amplified bands (specific bands) with 18% and 7 unique bands (genetic markers) with 32%. The genetic markers were 575 bp (S1 isolate), 275 bp (S2 isolate), 1025 bp (S3 isolate) and 1450, 1025 and 855 bp (S5 isolate).




Fig. (2): Electrogram of agarose gel 15% showing DNA polymorphism based on RAPD analysis from the studied five *S. mansoni* against random primers (A8, A7 and A6)

M: DNA molecular, S1, S2, S3, S4 and S5 isolates, A8, A7 and A6 random primers.

Table (5) : Polymorphism and genetic markers among 5 isolates of *S. mansoni* using random primers by RAPD-PCR.

Primer	Polymorphism				Genetic markers of isolates					
	TAF	MAF	PAF	Unique	M.W (bp)	S1	S2	S3	S4	S5
A8	8	4	2	2	855 275		+		- -	++++
A7	7	4	-	3	1450 1025 855				- - -	+ + +
A6	7	3	2	2	1025 575	- +	- -	+ -	- -	- -
Total bands	22	11	4	7	7	1	1	1	-	4
Percentage	-	50	18	32	32	4.5	4.5	4.5	-	18

4. Discussion

Schistosoma are dioecious digenetic trematodes carrying a large (270 Mb) genome. Gaining Knowledge about the genome of these parasites is of importance for the understanding of their biology, mechanisms of drug resistance and antigenic variation that determine escape from the host's immune system (Franco et al., 2006). This aim of the present study the genetic diversity of five *S. mansoni* isolates selected on the basis of different in disease potential (Longand and Morand, 1998).

Several techniques of molecular biology for studies of genetic diversity in Schistosomiasis vectors have been used (Spada et al., 2002; Florence et al., 2004; Hertel et al., 2004; Abdel-Hamid et al., 2006; Jamjoom, 2006 and Haggag and El-Sherbiny, 2006).

Electrophoretic protein banding pattern of an organism can be used to elucidate reliable biochemical genetic markers of this organism. It can also provide information about structural genes and their regulatory systems which control the biosynthetic pathways of that protein banding pattern (Abdel Salam et al., 1992).

In the present study, variation in protein banding patterns between five isolates were revealed using SDS-PAGE electrophoresis. The obtained protein profiles were analyzed and scanned for gel quantitations using Gel Doc 2000 instrument and quantity one. Software package (Bio-Rad). For comparisons among the five isolates of *S. mansoni* were analyzed in the same gel. The same procedures of scanning and quantitations were done. The observed changes in protein banding patterns in the present study could be reasonably interpreted to be the result of gene mutation. This conclusion is in accordance with Brown and Langley (1979) and Abdel Salam et al. (1992). However, other

investigators El-Gamal El-Din et al. (1988) traced such changes back to the induction of chromosomal abnormalities such as bridges breakes, laggards and micronuclei which can lead to loss of some of the genetic material. Therefore, some electrophoretic bands disappeared due to the deletion of their corresponding bands.

Disappearance of some bands could also be explained on the basis of a mutational event at the regulatory genes which are suppressed at transcription level. Meanwhile, the appearance of new bands could be explained on the basis of a mutational event at the regulatory system of unexpressed gene(s) that activate them (Abd El-Salam et al., 1992).

Gel electrophoresis is a widely used tool in studies of genetic variability. The electrophoretic differences reflect the allelic variations of *S. mansoni* enzymes which might due to mutational events occurring in the schistosoma under stress and affecting the loci controlling the synthesis of isozymes.

Several factors may be considered as primary determinants of the number of bands observed on a gel, including. (a) The number of coding genes, (2) Their allelic states (homozygous or heterozygous) and (3) The quaternary of the protein products. The simplest case involves a single region of salinity with variant electromorphs (allozymes) observed in different individuals. Because allozymes are usually codominantly inherited, the presence and number of bands are depending on the number of polypeptide subunits contained in the active enzyme (Kahler and Allard, 1970).

An electrophoretic analysis was made on the homogenate of progeny of five *S. mansoni* for esterase isozymes in order to determine isozymes. Since *S. mansoni* were collected and from different locations, therefore separate electrophoretic runs

were made for some *S. mansoni* from each location. Esterase isozymes are enzymes that characterized by their common activity on many naphthyl ester substrates (Korochkin et al., 1973). The group of esterase isozymes is one of the largest and most complicated system that has been intensively studied in many organisms of animal kingdom in vertebrates, esterase isozymes are mostly monomeric. Consequently, each band of esterase activity reflects a structure of one polypeptide chain. Hence, each band on the gel represents the end product of one locus (allele). As revealed from Est zymograms, all bands were anndelly migrated and their distributions were varied, some bands were present in a certain *S. mansoni* and were lacking in another. A total number of 4 were present in all five *S. mansoni* (minomorphic) and could considered as common bans for all the five *S. mansoni* which seem to be necessary ones for the enzyme constitution. Bands No. 1, No. 2 and No. 3 observed only in *S. mansoni* isolatges S4, S3, S2 and S5 respectively which could be a result of gene expressed under stress. The result indicates that there were allelic variants in this locus in all *S. mansoni*. Suggesting a polymorphic type of inheritance for this enzyme.

The results obtained after RAPD-PCR are presented in Fig. (3) and molecular weights of each primer was summarized in Table (3-4). From the RAPD profiles generated by these primers, bands with MW 1450, 1025, 855, 700, 75 and 205 bp were present in some *S. mansoni* and absent in others (polymorphic), while the two bands MW 510 and 325 bp were present in all the five isolates (monomorphic). One unique band was detected in S1 isolate with MW of 575 bp while was not seen in the other isolates.

Fortunately, the averages of similarities among five *S. mansoni* were 52.5, 76.5, 70.5, 40.5 and 82.2% for S1, S, S3, S4 and S5 respectively. This we expect the similarity of genetic backgrounds of many *S. mansoni* isolates.

The introduction of RAPD-PCR technique has amplified the possibilities of polymorphisms analysis, as it allowed the use of small arbitrary nucleotide segments without the need of a previous knowledge of genes and/or genomic sequences (Welsh and Mc Clelland, 1990). Also, RAPD assay clearly has certain proactively advantages for detecting DNA variation. It is technically less demanding, cheaper and quicker than other molecular techniques (Stothard and Rolliknson, 1996).

Rollinson et al. (1998) reported that, snails resistant to infection occur naturally and there is a genetic basis for this resistance, in *B. glabrata* resistance to *S. mansoni* is known to be polygenic trait. So, we have initiated a preliminary search for

snail genomic regions linked to, or involved in resistance by using a RAPD-PCR based approach in conjunction with progeny individual methods.

The present study, demonstrated the utility of RAPD-PCR method for the differentiation of *S. mansoni* isolates. The isolates studied were selected on the basis of different location and pathogenic by well characterized resistance, susceptibility phenotypes upon exposure to *S. mansoni*. Reproducible and inheritable stable polymorphic markes for *S. mansoni* were identified with one out of the ten arbitrary primers tested. Several of the primers produced monomorphic bands among the stocks, or the polymorphism identified were not reproducible. In previous studies, genetic diversity among *S. mansoni* snails was evaluated either by allozyme or restriction fragment length.

5. References:

1. Abdel Salam, A.Z.E.; S.A. Mansor and M. Abdel-Mordy (1992). Isozyme polymorphism in *Drosophila*. VII. Sensitivity of three major chromosomes to the effect of aromatic organophosphorus compounds as revealed by the induced mutations at twelve enzyme loci in *D. melanogaster*, Egypt J. Genet. Cytol. 21: 105-13.
2. Abdel-Hamid, Z.A.; S.M. Rawi and Azza, F. Arafa (2006). Identification of a genetic marker associated with the resistance to *Schistosoma mansoni* infection using random amplified polymorphic DNA analysis. Mem. Inst. Oswaldo Gruz, Rioo de Janeiro Vol. 101 (8): 863-868.
3. Brown, A.J. and C.H. Langley (1979). Reevaluation of level of genic heterozygosity in natural population of *D. melanogaster* by two dimensional electrophoresis. Proc. Nail. Acad. Sci, USA. 76 (6): 238-2384.
4. Da ilva, D.; R.G. Spada; S.S. Sobral-Hamaguchi; Z. Abdel-Hamid; N.R. Zuim; E.M. Zanotti-Magalhaes; A.L. Magalhaes and J.T. Riberiro-Paes (2004). *Biamphalaria tenagophila* Genetic Variability Within Intermediate Snail Hosts Susceptible and Resistant to *Schistosoma nansoni* Infection. Parasite, 11: 43-19.
5. Florcnce, M.R.; L.C. Roberta; C. Omar do Santos; B.G. Ana Lucia and Paulo Z.C. Marcos (2004). Dominant charaxcter of the molecular marker of a *Biophalarin tenagophila* strain(Mollusca planorbidae) resistant to *Schistosoma mansoni*. Mem. Inst. Oswaldo Cruz. 99: 85-87.

6. France, G.R.; A.F. Valadao; V. Azevedo and E.M. Rabelo (2006). The *Schistosoma* gene discovery programs state of the art. *Int. J. Parasitol.*; 10: 53-63.
7. Gamal El-Din, A.Y.; E.H.A. Hussein and M.A. Eweda (1988). Variation in chromosome number and its bearing on electrophoretic protein banding pattern in vien. *Bull. Fac. Agric., Cairo Univ*, 39 (1): 143-153.
8. Haggag, S.H. and M. El-Sherbiny (2006). Molecular markers associated with resistanc e to *Schistosoma mansoni* infection in the *Biomphalaria glabrata* snails. *Biotechnology* 5 (4): 404-412.
9. Hertel, J.; K. Keddves; A.H.M. Hassan; B. Haberele an W. Has (2004). Detection of *Schistosoma mansoni* cercariae in plankton samples by PCR. *Acta Tropica* 91: 43-46.
10. Jonathan, F.W. and N.F. Wondell (1990). Visualization and interpretation of plant isozymes. In *isozymes in Plant Biology-* D.E Saltis and P.S. Soltis (eds). London Chapman and Hall, pp. 5-45.
11. Kahler, A.L. and R.W. Allard (1970). Genetics of isozyme variants in barley 1. Esterases. *Crop Science* 10: 444-448.
12. Korochkin, L.I.; N.M. Matveev and M.B. Evageniev (1973). Expression of esterase isozyme in hybrids of *Drosophila virils*. *Genetika* 9 (12): 55-61.
13. Laemmli, J.K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T₄. *Nature* 227: 680-685.
14. Langand, J. and S. Morand (1998). Heritable non susceptibility in an allopatric host-parasitic system: *Biomphalaria glabrata*(Mollusca). *Echinotoma caproni* (*Platyhelminth digenea*). *J. Parasitol*, 84: 739-742.
15. Lockyer, A.E.; C.S. Jones; L.R. Noble and D. Riollinson (2000). Use of differential display to detect changes in gene expression in the intermediate snail host *Piomphalaria glabrata* upon infection with *Schistosoma mansoni*. *Parasitology*, 120: 399-407.
16. Lowry, O.H.; N.J. Resebriugh; A.L. Farr; R.J. Ranall (1951). Protein measurement with the folin phenol reagent. *J. Biol. Chem.* 193: 265-275.
17. Manal, B. Jamjoom (2006). Molecular identification of some *Schistosoma mansoni* isolates in Saudi Arabia. *World Journal of Medical Sciences* 1 (2): 102-107.
18. Saoudl, M.F. (1966). The infectivity and pathogenicity of geographical strains of *Schistosomi mansoni*. *Trans. Royal Soc. Trop. Med. and Hyq.* 69: 583-600.
19. Shalaby, I.M.; A.A. Banaja and A.M. Ghandour (1993). Comparative scanning electron microscopy study of the regimetal surface of *Schistosoma mansoni* (Egyptian and other strains) in mice. *J. Egypt. Ger. Soc. Zool.*, 12: 37-41. Da Silva, D., R.G. Spada; S.S. Sobral-Hamag.
20. Simpson, A.J.G.; E.D. Neto; M. Steindel; O.L.S.D. Gaballero; L.K.J. Pyssos and S.D.J. Pena (1993). The use of RAPDs for the analysis of parasites DNA Fingerprinting State of the Sciences 331-337.
21. Spada, R.G.; Dida Silva; A.Z. Abdel-Hamid, S.S. Sobral-Hamaguctu. N.R. Zuim; E.M. Zanotti- Magalhaes, L.A. Magalhaes and J.T. Riberio-Paes (2002). Genetic markers between *Biomphalaria glabrata* snails susceptible and resistant to *Schistosoma mansoni* infection. *Mem. Irst, Oswalde. Cruz*, 97: 53-58.
22. Stothard, J.R. and D. Rolliknson (1996). An evaluation of random amplified polymorphic DNA (RAPD) for identification and phylogeny of freshwater snails of the genus *Bulinus* (Gastrophoda *Planorbide*). *J. Mol. Stud.* 62: 165-176.
23. Welsh, J. and M. Mc. Clelland (1990). Fingerprinting genome using PCR with arbitrary, primers. *Nucleic acids. Res.* 18: 7213-7218.

5/5/2010