

The impact of genetic variability and smoking habits on the prevalence of periodontitis among adults

Faten S.Bayoumi¹, Fatehya.M.Metwaly², Hind M.Rashd² and E.H.A. Abouel-Ezz.³

¹Professor in Environmental &Occupational Medicine Department, National Research Center

²Assistant Professor in Environmental &Occupational Medicine Department, National Research Center

³Professor in genetic orodental Dep. National Research Center

fatenbayoumi@yahoo.co.uk

Abstract: Aim : Elucidate the effect of genetic variance of inflammatory mediators expression ,the influence of microbial expression, and smoking as a risk factors for periodontitis

Material &Methods: Sample of this study composed of 50 smokers & 50 non smoker volunteers (unrelated and of the same ethnic population) with 40-60 years old .Their periodontal status was estimated through periodontal examination (full mouth clinical attachment loss measurement ,probing depths ,plaque index scores, and bleeding on probing). Isolation and detection of certain oral pathogens; A.actinomycetemcomitans , Porphyromonas gingivalis ,and Prevotella intermedia was performed . Genotype for bi-allelic IL-1A+4845, IL-1B+3954 gene polymorphisms using mouth wash was detected by PCR based methods.

Results: There were a significant difference only between the two groups (smokers &non-smokers) as regards to colonization of A.actinomycetemcomitans & not among Porphyromonas gingivalis & Prevotella spp. There were no significant difference between the overall frequencies of carrying allele 2 of IL-1 A, IL-1B among smoker and non-smokers. The percentage of non smokers having healthy periodontal status was much higher than smokers. On the other hand, smokers recorded much higher percentage for mild, moderate and severe periodontitis. The difference was statistically significant concerning the percentage of those with severe periodontitis.

Conclusion: Environmental factors play either a direct (i.e., causative factor) or indirect (modifying factor) role as a risk factor for periodontitis. The association between genetic polymorphism of allele 2 of IL-1 A, IL-1B expression & smoking habits caused a synergistic effect for progression of periodontitis. Smoking initiated A.actinomycetemcomitans growth

Keywords: genetic polymorphism, periodontitis, Interleukin -1, periodontal pathogens, smoking

1. Introduction

The oral cavity is vulnerable to external agents as cigarette smoking exposure which causes oral changes in both hard and soft tissues Susin,et al.,(2004).

Periodontitis is a chronic inflammatory disease initiated by specific bacteria that activate host mechanisms destroying bone and connective tissues which support the teeth. Substantial data supported the current concept that specific bacteria are essential for initiation and progression of chronic periodontitis (Page et al., 1997), but the rate of progression and disease severity are determined by host modifiers such as smoking (Bergstrom, 1989),diabetes, (Collin et al., 1998) and genetic influences.(Kornman,2006)

Smoking has major effects on the host response, but there are also a number of studies recorded some microbiological differences between smokers and non-smokers (Gomes ,et al., 2006).

The pro-inflammatory cytokine interleukin-1 (IL-1) is a key regulator of the host responses to microbial infection and a major modulator of

extracellular matrix catabolism and bone desorption. It has been reported that variations in the IL-1 gene cluster on chromosome 2 are associated with increased susceptibility to severe periodontitis (Mc Devitt, et al., 2002).Therefore, a genetic test was being marketed to predict risk for periodontal disease progression (Higashi ,2002).

2.Aim of the work:

Elucidate the effect of genetic variability including the variance of inflammatory mediators expression ,the influence of microbial expression, and smoking as effects influencing risk for periodontitis.

3.Subjects and methods

Subjects:

Sample of this study composed of 50 smokers & 50 non smokers volunteers (unrelated and of the same ethnic population) with 40-60 years old. Both groups were interviewed and filled a detailed questionnaires for family history, dental, medical as well as smoking habits,. Cigarette consumption

was calculated (i.e. mean numbers of packs/day × number of years smoked)

Methods:

1-Periodontal Examination:

Criteria for assessment of the severity of periodontitis:

- 1- Mild periodontitis: Mean CAL ≥ 0.6 mm to 1.5 mm, no. interproximal sites with CAL ≥ 3 mm. No more than 3 missing teeth with the exception of orthodontic purpose, teeth lost as a result of extra oral trauma or extensive decay, or teeth that were congenitally missing.
- 2- Moderate periodontitis: Mean CAL ≥ 1.6 mm to 2.4 mm and ≤ 8 mm, interproximal sites with CAL ≥ 3 mm distributed through at least 3 quadrants or at 6 teeth. No more than 5 missing teeth with the exception of third molars, teeth extracted for orthodontic purpose, teeth lost as a result of extra oral trauma or extensive decay, or teeth that were congenitally missing
- 3- Severe periodontitis: Mean CAL ≥ 2.5 mm and 1 or more sites in 3 out of 4 quadrants with interproximal sites with CAL ≥ 5 mm. No more than 14 missing teeth with the exception of third molars, teeth extracted for orthodontic purpose, teeth lost as a result of extra oral trauma or extensive decay, or teeth that were congenitally missing

2-Microbiological Examination

Sampling : Paper-point samples were taken from the 4 deepest sub-gingival sites in each quadrant of the dentition (Mombelli et al., 1991, 1994). Samples were then placed in sterile pre-reduced anaerobically transport fluid (RTF) and transferred to the laboratory within 10 minutes.

Culturing: The samples were dispersed for 6 sec. with a vortex mixer and 10 fold serially diluted in RTF. Aliquots of 0.1 ml of appropriate dilutions were placed in duplicate onto specific media for different microorganisms in concern.

Samples were grown anaerobically (80% N₂, 10% H₂, 10% CO₂) at 37°C on 5% horse blood agar plates (Oxoid no. 2, Basingstoke, England) enriched with hemin (5 mg/L) and menadione (1 mg/L) for detection of *Porphyromonas gingivalis* and on Trypticase soy serum bacitracin-vancomycin (TSBV) medium in air with 5% CO₂ at 37°C for the selective isolation of *A. actinomycetemcomitans*.

KVLB-2 (kanamycin 75 μ g/ml-Vancomycin 2 μ g/ml laked blood agar) for isolation of pigmented and non pigmented *Prevotella* spp.

Identification: *Porphyromonas gingivalis* was identified on the basis of Gram stain, anaerobic growth, and the inability to ferment glucose, the production of indole, and a positive hemagglutination test with 3% sheep erythrocytes.

A. actinomycetemcomitans was identified on TSBV plates, based on typical colony morphology and positive catalase reaction. The percentage of the microorganisms of total colony-forming units (CFU) was counted.

3- PCR based methods:

a-DNA Isolation from Mouthwash

DNA from all subjects was isolated according to the method of de Vries et al. (1996) as modified and validated for the study of cytokine gene polymorphisms (Laine et al., 2000). In short, each individual rinsed out his/her mouth with 10 mL of 0.9% saline for 60 sec. Buccal epithelial cells were centrifuged at 300 x g for 10 min. The pellet was washed twice in 0.9% saline, re-suspended in 100 μ L of 50 mM NaOH, and boiled for 10 min. Samples were neutralized with 14 μ L of 1 M Tris (pH 7.5) and centrifuged at 14,000 x g for 3 min. Supernatants were collected and stored at 4°C until analysis.

b-Analysis of Polymorphisms in Genes of the IL-1A & B

The bi-allelic polymorphisms at position -889 within the promoter region of the IL-1A gene (McDowell et al., 1995) and at position +3954 (Taq I RFLP) within exon 5 of the IL-1B gene (Bioque et al., 1995), were determined according to previously described methods.

4.Results

Table (1): Percentage of bacterial species in plaque samples of smoker and non smoker individuals examined

Bacterial species	% of Non Smoker individuals	% of smokers individuals	Z.score	P
Gram (-) facultative rods	6%	30%		
A. actinomycetemcomitans	(3/50)	(15/50)*	1.96*	0.02
Gram (-) anaerobes	12%	24%	1.3	0.09
Porphyromonas gingivalis	(6/50)	(12/50)		
Gram (-) anaerobes	8%	16%	0.92	0.17
Prevotella intermedia	(4/50)	(8/50)		

The distribution of isolated microorganisms among examined groups was illustrated in table (1). Porphyromonas gingivalis, Prevotella spp and A.actinomycetemcomitans colonized 24%, 16%, and 30% of the smokers and 12%, 8% and 6% of non-smokers respectively.

There were significant difference between the two groups (smokers & non-smokers) as regards to colonization of Gram (-ve) facultative rods (A. actinomycetemcomitans) & no significant difference between the other two organisms.

Table (2): Distribution of smokers and non smokers of composite IL-1 genotype of allele 2 carriage of IL-1A(+4845) & IL-1B (+3953) among sam

IL-1 genotype of allele 2 carriage of IL-1A(+4845) & IL-1B(+3953)	IL-1A	IL-1B	Total carriers of allele 2
	Non smokers	14/50(28%)	3/50(6%)
Smokers	15/50 (30)	5/50(10%)	20/50(40%)
Total studied	29/100 (29%)	8/100(8%)	37/100(37%)

P value ;0.7

The percentage of smokers & non smokers carried allele 2 of IL-1A was more than those carried IL-1B (30%,28%& 10%,6%)

respectively, table (2). The difference was statistically non-significant.

Table(3): Distribution of allele 2 frequency of IL-1A(+4845) & IL-1B (+3953) composite genotype among studied individuals in relation to severity of periodontitis

	Total carriers of IL-1 genotype of allele 2 (IL-1A(+4845) & IL-1B(+3953))		Z scc	P.val
	Non smokers (n=)	Smokers (N=20)		
Healthy	(8/17) 47%*	0/20(0%)	3.0	0.00
Mild	Periodontal status	15%(3/20)	0.2	0.42
Moderate	(2/17) 11.8%	20%(4/20)	0.2	0.4
Sever	(5/17) 29.4%	65%(13/20)*	1.8	0.03

Smokers and non-smoker individuals carried allele 2 of IL-1A and IL-1 B were further divided according to the severity of the periodontitis. The percentage of non smokers having healthy periodontal status (47%) was much higher compared to smokers and the difference was significant. On the other hand, smokers & non smokers recorded nearly a similar percentage among those complained from mild, moderate (15%, 20% & 11.8%, 11.8%) with no statistical significant difference. Whereas, statistically significant difference was noted concerning the percentage of individuals in both groups

5. Discussion

Periodontitis is a multifactorial chronic inflammatory disease. However, it is difficult to ascertain the role of the different factors involved in its pathogenesis. Cigarette smoking is associated with increased prevalence and severity of destructive periodontitis in terms of periodontal pocketing, periodontal bone loss, and tooth loss (Gomes, 2006).

In microbiological study we focused on a number of microbial species e.g., *Actinobacillus actinomycetemcomitans*, *Porphyromonas gingivalis*, and *Prevotella intermedia* which were proposed to be a useful tool for the identification of susceptible individuals (Slots and Listgarten, 1988, Maiden et al., 1990 and Haffajee et al., 1994). Substantial data supported the current concept that specific bacteria are essential for initiation and progression of chronic periodontitis (Page et al., 1997), but the rate of progression and disease severity are determined by host modifiers such as smoking, diabetes, and genetic influences.

In the present study approximately half of the smokers and one fourth of the non smokers harbored these three microorganisms in their oral cavity. There was significant difference between the two groups (smokers & non-smokers) as regards to colonization of Gram (-ve) facultative rods (*A. actinomycetemcomitans*) & no significant difference between the other two organisms.

Genes which encode inflammatory cytokines were subject to polymorphisms in their regulatory regions that might affect both the level and the ratio of cytokines produced in response to exogenous stimuli. These variant alleles were observed in a large percentage of the population and were often associated with increased or decreased susceptibility or severity (modifiers) to infectious, immune or inflammatory diseases (Yucesoy et al., 2003). Axelsson (2002) reported that two factors, smoking and IL-1 genotype, significantly increased the risk of progression of alveolar bone loss and tooth loss due to progressive periodontitis. Moreover, the effect was synergistic: 41% of the IL-1 genotype-positive smokers lost 2 teeth,

complained from severe periodontitis (65% & 29.4% respectively), table (3) agreed with Kornman et al., (1997) who reported the same correlation and explained this finding as genetic mechanism by which some individuals, if challenged by bacterial accumulations, may have more vigorous immune-inflammatory response leading to more severe periodontitis. Moreover, Kornman (2006) added that monocytes from individuals homozygous for the IL-1 B +3953 allele 2 produce four-fold more IL-1 β and heterozygous cells produce approximately two-folds more IL-1 β from individuals homozygous for allele 1, compared with roughly 11% of those who had only one of the risk factors.

Our results correlated the severity of periodontitis to presence of carriers of allele 2 genotype in the IL-1A and IL-1B genes. A data

Our results showed that the nature of the host response is determined primarily by genetic factors, environmental and acquired factors (smoking). The complex interactions that occur between host-response mechanisms and oral pathogens in periodontal disease have made elucidation of genetic factors in disease susceptibility more difficult (Hassell et al., 1995).

6. Conclusion

Environmental factors play either a direct (i.e., causative factor) or indirect (modifying factor) role as a risk factor for periodontitis. The association between genetic polymorphism of allele 2 of IL-1 A, IL-1B expression & smoking habits caused a synergistic effect for progression of periodontitis. Smoking initiated *A. actinomycetemcomitans* growth

Corresponding Author

Dr. Faten S. Bayoumi
Assistant professor of microbiology
Immunogenetics Department
National Research Center, Cairo Egypt
E-mail: fatenbayoumy@yahoo.co.uk

References

1. Pociot F., Molvig J., Wogensen L., Worsaae H., and Nerup J: European Journal of Clinical Investigation, 1992; 22, 396.
2. Axelsson P.: Diagnosis and risk prediction of periodontal diseases. Vol 3. Chicago, IL: Quintessence Publishing Co 2002: 146-63.
3. Hassell TM, Harris EL: Genetic influences in caries and periodontal diseases. Crit Rev Oral Biol Med, 1995; 6: 319-342.
4. Gomes SC, Piccinin FB, Oppermann RV, Susin C, Nonnenmacher CI, Mutters R, Marcantonio

- RA. Periodontal status in smokers and never-smokers: clinical findings and real-time polymerase chain reaction quantification of putative periodontal pathogens Sep 2006;77(9):1483-90
5. Slots J, Listgarten MA :Bacteroides gingivalis, Bacteroides intermedius and Actinobacillus actinomycetemcomitans in human periodontal diseases. /Clin Periodontol 1988;15:85-93.
 6. McDowell TL, Symons JA, Ploski R, F0rre O, Duff GW. A genetic association between juvenile rheumatoid arthritis and a novel interleukin-1 alpha polymorphism. Arthritis Rheum J. 1995;38:221-228
 7. Bioque G, Crusius J A, Koutroubakis I, Bouma G, Kostense PJ, Meuwissen SGM, et al. Allelic polymorphism in IL-13 and IL-1 receptor antagonist (IL-1RA) genes in inflammatory bowel disease. Clin Exp Immunol. 1995; 102:379-383.
 8. Maiden MFJ, Carman RJ, Curtis MA, Gillett IR, Griffiths GS, Sterne JAC, et al. . Detection of high-risk groups and individuals for periodontal diseases: laboratory markers based on the microbiological analysis of subgingival plaque. / Clin Periodontol, 1990: 17:1-13.
 9. Katz, R.S., Premenko-Lanier M., McChesney M.B., Rotaa P.A. and Bellini W.J.: Detection of measles virus RNA in whole blood stored on filter paper. J. Med. Virol. 2002;67:596-602
 10. Higashi K.M.: The Cost-Effectiveness of Interleukin-1 Genetic Testing for Periodontal Disease. Journal of periodontology. J Periodontol, 2002;73:1474-1484.
 11. McDevitt M .J., Wang H. , Knobelmann C., Michael G. Francesco N., di Giovine S. , Timms J. , Duff G.W. , Kornman K.S. (2002): Interleukin-1 Genetic Association With Periodontitis in Clinical Practice. J Periodontol 2000;71:156-163
 12. Susin C, Vecchia CFD, Oppermann RV, Haugejorden O, Albandar JM: Periodontal attachment loss in an urban population of Brazilian adults: effect of demographic, behavioral, and environmental risk indicators. J Periodontol, 2004: 75:1033-41
 13. Bergstrom J.: Cigarette smoking as risk factor in chronic periodontal disease. Community Dent Oral Epidemiol, 1989: 17:245-247
 14. Kornman S.: Interleukin 1 genetics, inflammatory mechanisms, and nutrigenetic opportunities to modulate diseases of aging Am J Clin Nutr;83(suppl),2006::475S– 83S
 15. Page RC, Offenbacher S, Schroeder HE, Seymour GJ, Kornman KS. 1997:
 16. Advances in the pathogenesis of periodontitis: summary of developments, clinical implications and future directions. Periodontol 2000 ;,14:216–48.
 17. Lange DE,: Frequency, seriousness and treatment needs in periodontal diseases. ZWR, 1986: 95(4):402-6.
 18. Mombelli A, McNabb H, Lang NP . : Black-pigmenting Gram-negative bacteria in periodontal disease. II. Screening strategies for detection of P. gingivalis. J Periodontal Res, 1991: 26:308-313.
 19. Mombelli A, Gmuir R, Gobbi C, Lang NP . : Actinobacillus actinomycetemcomitans in adult periodontitis. I. Topographic distribution before and after treatment. J Periodontol, 1994: 65:820-826.
 20. Collin HL, Uusitupa M, Niskanen L, Kontturi-Narhi V, Markkanen H, Koivisto AM, et al. . Periodontal findings in elderly patients with non-insulin dependent diabetes mellitus. J Periodontol, 1998: 69:962-966.
 21. de Vries HG, Collee JM, van Veldhuizen MH, Achterhof L, Smit Sibinga CT, Scheffer H, et al. Validation of the determination of deltaF508 mutations of the cystic fibrosis gene in over 11 000 mouthwashes. Hum Genet, 1996: 97:334-336.
 22. Laine ML, Farre MA, Crusius JBA, van Winkelhoff A-J, Pefia AS,: The mouthwash: a non-invasive sampling method to study cytokine gene polymorphisms. J Periodontol, 2000: 71:13 15-1318.
 23. Albandar JM, Streckfus CF, Adesanya MR, Winn DM: Cigar, pipe, and cigarette smoking as risk factors for periodontal disease and tooth loss. J. Periodontol. 2000 Dec;71(12):1874-81