

## Micropropagation of Cassava (*Manihot esculantum* Crantz) Using Different Concentrations of Benzyaminiopurine (BAP)

Onuoha C.I.<sup>1</sup>, Onwubiko N.I.C<sup>2</sup>

1. Department of Plant Science and Biotechnology, Imo State University P.M.B. 2000 Owerri, Nigeria
2. Department of Crop Science and Technology, Federal University of Technology P.M.B. 1526 Owerri, Nigeria

Email: [onwubikouche@yahoo.com](mailto:onwubikouche@yahoo.com)

**Abstract:** An investigation conducted on the effect of Benzylaminopurine (BAP) on cassava plantlets cultures. Murashige and skoog medium containing different concentrations of BAP was used to subculture two varieties of cassava plantlets; TMS 98/0379 and TMS 98/0581, which were thereafter incubated in a culture room at 28°C ± 2 and exposed to artificial illumination of 2000 -2500 lux for sixteen hours daily. Results showed that BAP application had some inhibitory effects on the two cassava varieties as only the control treatment (no BAP application) recorded the best growth in most of the growth parameters observed (height, leaves, nodes), and thus differing significantly (P<0.05) from other BAP levels. Between the experimental units, TMS 98/0379 appeared to grow better in terms of height, leaf and number of nodes while TMS 98/0581 recorded the greatest fresh weight.[ Nature and Science. 2007;5(2):37-40] (ISSN:1545-0740).

### Introduction

Cassava improvement for increased production necessitates addressing various factors that beset its production, which include pests and diseases, requirement of large quantities of planting material for its propagation and low multiplication ratio (Dahniya and kellon, 1983; Okigbo, 1986). Although plant breeding and genetics has contributed immensely in the improvement of crop plants, it has been observed more recently that micropropagation has become an irreplaceable tool in the improvement and genetic manipulation of plants especially vegetatively propagated crops (Onwubiko and Mbanaso, 2006). And fortunately cassava is ranked fourth in order of importance after rice, wheat and maize in crops for micropropagation in Africa (Johan *et al*, 1998).

The establishment of culture media and adjustment of their concentrations are the keys to success in micropropagation. Generally, cassava researchers have used specific treatment to obtain good development (Guohua, 1998; Joseph *et al*, 1999; Peng *et al*; 2001; Matand *et al*, 2004). Also studies on other crops have reported controlling some Morphogenic developmental stages using a single treatment (Triagiano and Gray, 1996; Matand *et al*, 2004). This work therefore is an investigation on the effect of various levels of Benzylaminopurine (BAP) composition of culture media interacted with two cassava varieties for regeneration of cassava plants from stem explants.

### Materials and Method

This study was carried out in the tissue culture laboratory of National Root Crop Research Institute (NRCRI) Umudike, Abia State Nigeria. Umudike is located at latitude 5° 25<sup>1</sup>N, longitude 7° 35<sup>1</sup> and at 122m above sea level. The two cassava plantlets (TMS 98/0379 and TMS 98/0581) used for this study were collected from the tissue room in the tissue culture laboratory of NRCRI Umudike and the nutrient media (BAP) contained inorganic and organic constituents. Six different levels of BAP concentrations (0, 0.25, 0.5, 0.75, 1.0 and 1.25 Mgl<sup>-1</sup>) were used for the study.

The two cassava varieties used were five months old plantlets cultured on a basal medium with height above 5cm. The Plantlets were brought out on Petri dishes using sterile forceps and the tissues subdivided into one-node length sterile scalpel. The node cuttings were transferred into fresh media of different BAP concentrations. The necks of the culture vessels were flamed before replacing the cover. All these operations were carried out under strict aseptic conditions in the laminar airflow cabinet. The culture vessels were carefully labeled and incubated in the culture room at 28°C ±2 where they received artificial illumination of 2000-2500 Lux for 16 hours daily.

The experimental design used for this study was a 2x6 factorial in a completely randomized design (CRD) with 5 replications.

Data collection on plant height, number of leaves, number of developed nodes were collected on 2 weeks interval for 6 weeks after subculture (WAS). With electronic weighing balance fresh weight of the

plantlets was determined after removal of plantlets from culture vessels on the 6<sup>th</sup> week. The statistical analysis used in analyzing the result collected was genstat.

### Results

At 2,4 and 6 WAS, BAP application did not influence plant height in the two cassava varieties, rather in the control there was significant difference ( $p < 0.05$ ) in plant height as the greatest mean (3.88) was recorded (Table 1) Again at BAP level  $1.25 \text{ mg l}^{-1}$  TMS 98/0581 recorded a high percentage of callus formation showing *in vitro* recalcitrance.

The result on number of cassava leaves at 2,4 and 6 WAS was almost similar to that of plant height. Increase or decrease in BAP concentration was not consistent with leaf formation in the two cassava varieties. However at  $0.75 \text{ mg l}^{-1}$  BAP, TMS 98/0581 recorded the highest number of leaves at 6 WAS (Table 2) although this was not significantly ( $P < 0.05$ ) greater than the leaf formation in the control.

A significant difference in the number of nodes was however observed in 2,4 and 6 WAS between the two cassava varieties (Table 3) Apparently, TMS 98/0379 had the highest number of nodes. Generally, the highest number of nodes was observed at 4 and 6 WAS at  $0.00 \text{ Mg l}^{-1}$  BAP concentration, although this was not significantly different from other hormonal levels.

Similar to the result observed in the number of cassava leaf, the flesh weight of the two cassava varieties of constants increase or decrease in BAP concentration (Table 4). The highest fresh weight was recorded for the two cassava varieties at  $1.0 \text{ Mg l}^{-1}$  BAP concentration although it was not significantly different ( $P > 0.05$ ) with the result from the control and other levels of BAP. Comparatively TMS 98/0581 recorded a greater fresh weight.

Table 1. Effect of different concentration of BAP on plant height at 2, 4 and 6 WAS

Treatments	Weeks after Subculture		
	2	4	6
Cassava varieties			
TMS 98/0379(c <sub>1</sub> )	1.30	1.72	2.12
TMS 98/03819 (c <sub>2</sub> )	1.04	1.72	1.12
LSD (0.05)	0.28	1.19	1.47
LSD (0.05)			
BAP $\text{Mgl}^{-1}$			
0.00 (B <sub>0</sub> )	1.78	2.58	3.88
0.25 (B <sub>1</sub> )	1.26	1.47	1.53
0.50 (B <sub>2</sub> )	1.24	1.46	1.72
0.75 (B <sub>3</sub> )	1.10	1.49	1.67
1.00 (B <sub>4</sub> )	1.79	0.80	0.99
LSD 0.05	0.48	0.57	0.64

Table 2. Effect of different concentration of BAP on number of leaves at 2,4and 6 WAS

Treatments	Weeks after Subculture		
	2	4	6
Cassava varieties			
TMS 98/0379(c <sub>1</sub> )	1.65	2.84	2.87
TMS 98/03819 (c <sub>2</sub> )	0.95	2.26	2.78
LSD (0.05)	0.46	0.80	0.91
LSD (0.05)			
BAP $\text{Mgl}^{-1}$			
0.00 (B <sub>0</sub> )	2.20	3.70	4.40
0.25 (B <sub>1</sub> )	1.32	2.47	2.14
0.50 (B <sub>2</sub> )	1.02	2.95	3.43
0.75 (B <sub>3</sub> )	1.18	2.82	3.43
1.00 (B <sub>4</sub> )	1.00	2.00	2.00
LSD 0.05	1.07	1.35	1.54
LSD (0.05)	0.80	1.39	1.58

Table 3. Effect of different concentration of BAP on number of nodes at 2,4and 6 WAS

Treatments	Weeks after Subculture		
	2	4	6
Cassava varieties			
TMS 98/0379(c <sub>1</sub> )	3.04	3.75	4.74
TMS 98/03819 (c <sub>2</sub> )	1.60	2.64	3.75
LSD (0.05)	0.47	0.73	1.00
LSD (0.05)			
BAP Mgl <sup>-1</sup>	2.50	4.50	6.10
0.00 (B <sub>0</sub> )	2.88	3.36	4.22
0.25 (B <sub>1</sub> )	2.48	3.61	4.71
0.50 (B <sub>2</sub> )	2.34	3.46	4.33
0.75 (B <sub>3</sub> )	2.10	2.30	3.30
1.00 (B <sub>4</sub> )	1.63	1.94	2.82
LSD 0.05	0.81	1.27	1.74

Table 4. Effect of different concentration of BAP on number of nodes at 2, 4 and 6 WAS

Treatments	6 Weeks after Subculture
Cassava varieties	
TMS 98/0379(c <sub>1</sub> )	0.18
TMS 98/03819 (c <sub>2</sub> )	1.23
LSD (0.05)	2.06
LSD (0.05)	
BAP Mgl <sup>-1</sup>	0.25
0.00 (B <sub>0</sub> )	0.14
0.25 (B <sub>1</sub> )	0.19
0.50 (B <sub>2</sub> )	0.11
0.75 (B <sub>3</sub> )	0.42
1.00 (B <sub>4</sub> )	0.13
LSD 0.05	3.56

### Discussion

The *in vitro* application of different concentration of synthetic cytokinins (BAP) considerably inhibited the performance of the two cassava varieties used in the study. The fact that all the parameters evaluated were better in the control treatment (no BAP application) shows that the endogenous levels of BAP in these cassava varieties were adequate. Hence, exogenous application of BAP led to supra – optimal amounts which induced some inhibitory effects. This observation is consistent with the physiological behaviour of hormones which have two concentration maxima for promotive and inhibitory effects. Berrie (1984) reported that synthetic cytokinins are inhibitory at high concentration. Again, the effect of any particular exogenously applied growth hormone is influenced by a variety of other factors in the internal environment of the plant, especially other hormones in the plant (Curtis *et al*, 1985; Preece, 1987).

Comparatively, TMS 98/0379 appeared to have performed better in terms of height, leaves, and nodes. This is in agreement with the findings of Curtis *et al*. (1985), who reported that the response to a particular hormonal message does not only depend on its content but also upon how it is “read” by its recipient. The *in vitro* recalcitrance observed in TMS 98/0581 at BAP concentration 1.25 Mgl<sup>-1</sup> is also in agreement with the findings on *in vitro* propagation of cassava; that above 1.0 µm BAP, shoot lengths were decreased which made subculture of nodes more difficult (Smith *et al*, 1986). Also, Anura (2006) observed that higher levels of kinetin (Cytokinin) induced meristem cultures to form callus.

### Conclusion and Recommendation

Apparently, from the forgoing, the use of plant growth regulators like BAP has great potential and can offer meaningful and useful results. However, cassava cultures using basal media (no hormone) appears to be the most practical and effective method for maximum *in vitro* cassava performance.

**Received:** 6/22/2007

## References

1. Anura, H. (2006). Tissue culture and meristem culture in sweet Potato M. Sc. Thesis Unpublished. pp. 1-10.
2. Berrie, A.M. (1984) Germination and growth regulators. In: Advance Plant Physiology. Malcom. W.B. ed. English Language Book Society/Longman England. 987 pp.
3. Cutis, H. and Barnes, N. (1985) Invitation to Biology, Worth Publishers. New York 4<sup>th</sup> edition. pp 373.
4. Dahniya, N.T. and Kallon, S.N. (1983) Rapid Multiplication of Cassava by direct Planting. In: Tropical Root Crops Production and Uses in Africa. Terry, E.R., Doku, E.V., Arene, O.B. and Mahnungu, N.M. (eds) Proceedings of the second Triennial symposium of the International Society of Tropical Root Crops – Africa Branch held in Douala Cameroun, 14-19 August, 1983.
5. Guohua, M. (1998). Effects of Cytokinins and auxins on cassava shoot organogenesis and somatic embryogenesis from somatic embryo explants.I. Plant Cell Tissue and organo cultures 54 (1): 1-7.
6. Johan, A.B., Barbara, R.W. and Edgar J.D.(1998). Plant Biotechnology: a tool for development in Africa. Electronic Journal of Biotechnology Vol.1.N0 3.
7. Joseph, T., Yeoh, H.H. and Loh, C.S. (1999) cynogenesis in somatic embryos and plantlets of cassava (*Minohot esculentum* Crantz) J. sci food Agric 79:1071-1074.
8. Matand, K., Porobo, A.D., Prakash, C.S (1994) thidiazuron promotes high frequency regeneration system for peanut (*Arachis hypogaea* L.) plant *in vitro* plant cell reports 14:1-5
9. Onwubiko, O. and Mbanaso E.N.A (2006). Micropropagation for the enhancement of food production. In Repositioning Agriculture for Sustainable Millennium Development Goal. Asumugha, G.N., Olojede, A.O., Ikeorgu, J.G., Ano, A.O and Herbert U. (eds) proceedings of the 40<sup>th</sup> Annual conference of Agriculture Society of Nigeria held in Umudike Abia State, 16<sup>th</sup> -20<sup>th</sup>, October 2006.
10. Okigbo, B.N. (1986) Root and tuber in the Africa food crisis. In: Tropical Root crops. Terry E.R., Akoroda, M.O. and Arene, O.B.(eds). Proceedings of the third triennial symposium of Intentional Society for Tropical Root Crops- Africa Branch held in Owerri. Nigeria, 17-23 August, 1986
11. Peng, Z., Salak.P., Johanna, P.K. (2001). Improvement of cassava shoot organogenesis by the use of silver nitrate *in vitro* plant cell, Tissue and organ culture 67(1) 47-54.
12. Preece, J.E.(1987) treatment of the stock plant with plant growth regulators to improve propagation success. Hort science, 22(5) 54-759.
13. Trigiano, N.R. and Grey, J.D. (1996) plant tissue culture concepts and laboratory exercise. CRC press. N.Y 374 PP.