

Anthelmintic Effect of *Solanum lycocarpum* in Mice Infected with *Aspicularis tetraptera*

¹A.P.S. Cruz, ¹D.P.C. Costa, ³G.S.C. Valente, ³D.M.M. Mattos,
³D.J.A. Alexandre, ^{1,2,3}G.F. Diré and ¹H.R. Borba

¹Laboratório de Atividade Anti-helmíntica de Plantas, Departamento de Biologia Animal, Universidade Federal Rural do Rio de Janeiro, Instituto de Biologia, Br 465; Km 7-Seropédica, Rio de Janeiro, RJ 23890.000, Brazil

²Universidade Federal Rural do Rio de Janeiro, Instituto de Ciências Exatas, Setor de Bioquímica, Br 465, Km 7-Seropédica, Rio de Janeiro, RJ 23890.000, Brazil

³Campus R9/Taquara, Universidade Estácio de Sá, Centro de Ciências da Saúde, Rio de Janeiro, RJ 22710560, Brazil

[Nature and Science. 2008;6(2):20-23]. ISSN: 1545-0740.

Abstract: This approach intends to add new data on the helminthes parasites of laboratory mice. It has been investigated the anthelmintic activity of *Solanum lycocarpum* (*Solanaceae*) extracts against *Aspicularis tetraptera* in mice naturally infected. The extracts were applied for oral saw (intra-gastric), into the volume of 0.04 mL g⁻¹, with the employing of a dead and bend probe during three consecutive days. The fecal material, collected 24 h after each application, performing a total of four fecal collection, have been softened previously, transferred about to sieve of network of 125 µm and tested under microscope stereoscope, with the objective of behave the identification and counting from the worms eliminated of the second to the fifth day of the experimental. Tukey-Kramer Multiple Comparisons Test was applied to compare the results. According to the analysis of the results it was observed and an extremely significant difference between TM and C (from 5.64±3.16 to 1.56±3.16). It was published that medicinal plants which were reported as useful in the treatment of diabetes the *S. lycocarpum* was the sixth most frequently mentioned. According to the results obtained in the present study, we can speculate that the anthelmintic effect of *Solanum lycocarpum* was noticed due to the concentration of steroidal alkaloid oligoglycosides and short-chain fatty acids.

Key words: *Solanum lycocarpum*, *Aspicularis tetraptera*, anthelmintic, mice, extracts, medicinal plants

INTRODUCTION

Solanum lycocarpum was collected in the City of Três Marias, State of Minas Gerais and in the City of Seropédica, State of Rio de Janeiro. The botanical identification was carried through in the Department of Botany of the Rural Federal University of Rio de Janeiro, having been the exsiccates deposited under numbers RBR 28010 and RBR 14072.

S. lycocarpum is a plant which is shrubs ranging in height from 1.2 to 3 m. The fruit is yellow in color and resembles a medium sized tomato. Parts of the plant are poisonous if it gets in your system. When it is in bloom, it is medium blue. It blooms in the late winter, early spring, late fall, early winter and mid winter. It is velvety or fuzzy. It needs water regularly. It is found in the Brazilian savannah but has been said to grow in San Antonio, Texas. *S. lycocarpum* is commonly used in Brazilian folk

medicine. The Brazilian flora is one the world richest sources of bioactive material due to its biodiversity. Several plants are currently used in Brazilian traditional medicine to treat diabetes. *Solanum lycocarpum* St. Hill., *Solanaceae* has been widely used and commercialized as a hypoglycemic agent in Brazil. It was described by Vieira *et al.* (2003) the anti-inflammatory effects of the crude ethanol extract and its alkaloid fraction from *S. lycocarpum* fruits. Recently, it was carried out a chemical analysis of the starch and tried to correlate its supposed hypoglycemic activity with the polysaccharide content. However, these investigators did not conduct any experimental test to directly demonstrate the hypoglycemic effect attributed to the starch. *Solanaceae* or Lobeira is a plant used as a hypoglycemic agent. A study reported that the extract reduces glycoemia in alloxan induced diabetic rats. It was reported that the potential of *S. lycocarpum* as antioxidant was capable reduce in 27%

Corresponding Author: H.R. Borba, Laboratório de Atividade Anti-helmíntica de Plantas, Departamento de Biologia Animal, Universidade Federal Rural do Rio de Janeiro, Instituto de Biologia, Br 465; Km 7-Seropédica, Rio de Janeiro, RJ 23890.000, Brazil Fax: +552126821763/+552126821763

nitrate generation in diabetic animals. In literature has been demonstrated that *S. lycocarpum* is not ulcerogenic and restored haemoglobin and haematocrit to normal values in diabetic animals (Perez *et al.*, 2006).

It this plant contains steroidal glycoalkaloids that can be transformed into an intermediate for steroidal drug production. In this way, it is very possible that these glycoalkaloids and its aglycone, once in the body by ingestion of *S. lycocarpum* fruits, may act by disrupting the endocrine system. Because its fruits may be consumed by pregnant animals in the fields, various studies determined the possible toxic effects of exposure to *S. lycocarpum* fruit from gestation. The unripe fruits contained 0.6% of solamargine and 0.9% of solasonine. It was related that *S. lycocarpum*, during gestation and the beginning of lactation reduces intrauterine growth. It is known that during adulthood, female offspring showed impaired sexual behavior and male offspring showed prominent degeneration of testis germinative cells, characterized by a reduced number of germ cells and vacuolation. It has been documented that the exposed offspring showed reduced hypothalamic norepinephrine (NOR), vanillylmandelic acid (VMA), 3-methoxy-4-hydroxyphenylglycol (MHPG) and homovanillic acid (HVA) levels and reduced striatum NOR, HVA, VMA, MHPG, dopamine (DA), dihydroxyphenylacetic acid (DOPAC) and 5-hydroxyindolacetic acid (5-HIAA) levels. It is suggest that the fruit may act as an estrogen, with a long-term effect, impairing the receptive lordosis behavior of female offspring and promoting testis abnormalities in male offspring at adulthood. It appears to disrupt brain organization since important central monoamine level alterations were also related (Schwarz *et al.*, 2005a).

Rodents, as mice and rats are the most common laboratory animals used in research and testing. They are seldom investigated for autochthonous ecto and endoparasites prior their utilization in the experiments. Pinworms commonly infecting laboratory rodents include mainly the mice pinworms *Aspiculuris tetraptera* (Perec-Matysiak *et al.*, 2006; Gilioli *et al.*, 2000).

Some plant extract may act differently due to its action against the parasite. In a study the anthelmintic activity of the extracts obtained from *Luxemburgia octandra* was evaluated naturally infected mice with *Aspiculuris tetraptera* and *Vampirolepis nana*. The leaves extracts were given to the animals during three days. The ethanolic and ethyl acetate extracts presented significant increase of the *V. nana* elimination, but did not present the nematicide effect against *A. tetraptera* (Silva *et al.*, 2005).

In the present study we evaluated the anthelmintic activity of *Solanum lycocarpum* extracts in a concentration of 5% against *Aspiculuris tetraptera* in mice in naturally infected.

MATERIALS AND METHODS

Vegetal extracts: Dried leaves of units of had been used in the anthelmintic tests *Solanum lycocarpum* had been gotten by infusion (tea), submitted to the filtration in nylon and the express concentrations in g/100 mL (p/v).

Animals and anthelmintic tests: For anthelmintic test have been used lots of albinos mice, male and females weighted in media of 25 g and naturally infecting for *Aspiculuris tetraptera*, originated from Oswaldo Cruz Foundation-FIOCRUZ and held into the Institute of Biology from Rural Federal University from Rio de Janeiro. The animals have been held into bird cages individual of polypropylene (30×20×13 cm), it has at the bottom road of screen stark and stiff (network of 7×7 mm) upon a sheet of absorbent paper with the aim to facilitate the collection diary of excrement (Amorim *et al.*, 1987; Amorim and Borba, 1990). The referred study was conclude in the Laboratory of Activity Anthelmintic of Plants from Rural Federal University from Rio de Janeiro. The extracts were applied for oral saw (intra-gastric), into the volume of 0.04 mL g⁻¹, with the employing of a dead and bend probe during three consecutive days. The excrement, collected 24 h after each application, performing a total of four fecal collection, have been softened previously, transferred about to tames of network of 125 µm and evaluated under microscope stereoscope, with the objective of behave the identification from the worm eliminated of the second to the fifth day of the experimental. Into the fifth and last days from the tests, the mice have been sacrificing for inhalation of vapors of ether ethyl, examining humid weight of the contents of the small intestine, in order to access the number of the proglotes collected of *A. tetraptera* remnants (Amorim *et al.*, 1999). On the tests have been used the extracts of *S. lycocarpum* (leaves dried from Três Marias in the concentration of 10%) and (leaves dried from UFRRJ in the concentration of 10%). Additional lots of mice have been used with standard, they receiving doses of 20 mg kg⁻¹ day⁻¹ of mebendazol and 100 mg kg⁻¹ day⁻¹ of nitroscanato and they were submitted to the identical assessment anthelmintic description about to the animals treated with the plant extracts. A batch control, without a treatment served about to appraise the elimination spontaneous from the

helminthes studied. The outcome antinematode also was denominated in terms percentile average of roundworm eliminated, considering the number of roundworm eliminated in the excrement in relation to the total number. Statistical analysis were performed and Tukey-Kramer Multiple Comparisons Test was applied to compare the results.

RESULTS AND DISCUSSION

According to the analysis of the results it was observed that there were no differences ($p > 0.05$) in the % of elimination between TM and UR (from 5.64 ± 3.33 to 3.15 ± 3.16), UR and C (from 3.15 ± 3.16 to 1.56 ± 3.16) and an extremely significant difference between TM and C (from 5.64 ± 3.16 to 1.56 ± 3.16) (Table 1).

The extracts were applied for oral saw (intra-gastric), into the volume of 0.04 mL g^{-1} , with the employing of a dead and bend probe during three consecutive days. The excrements, collected 24 h after each application, performing a total of four fecal collection, have been softened previously, transferred about to tames of network of $125 \mu\text{m}$ and evaluated under microscope stereoscope, with the objective of behave the identification of the worm eliminated of the second to the fifth day of the experimental. Tukey-Kramer Multiple Comparisons Test was applied to compare the results.

Animal models have been exhaustively investigated regarding aspects related to their suitability for the development of experimental protocols under laboratory conditions. Nevertheless, in most of the adopted procedures, the prior detection of their ecto and endo parasites are generally overlooked related to the really effects of natural extracts in their biological cycle.

In the Brazilian cerate, a preparation obtained from the fruits of *Solanum lycocarpum* St.-Hill. (Solanaceae), popularly known as fruta-de-lobo (wolf-fruit), have been widely employed for diabetes management, obesity and to decrease cholesterol levels. The medicinal preparation consists of the green fruits which are ground in aqueous solution and filtered. The white gum deposited is decanted and slowly dried providing a powder which is

commercialized in capsules with the name of polvilho-de-lobeira. Through phytochemical analysis of this phytomedicine and the fruit of *S. lycocarpum* were found polysaccharides as the main component. Some polysaccharides slow gastric emptying and act on the endocrinous system affecting the liberation of gastrointestinal hormones, lowering blood glucose levels. According to Schwarz *et al.* (2005b) it is well known that this plant contain steroidal glycoalkaloids that can be transformed into an intermediate for steroidal drugs production, like oral contraceptives. In this way, it is very possible that these glycoalkaloids and its aglycone, once in the body by ingestion of *S. lycocarpum*, may act disrupting to the endocrine system as well as it may probably affect the reproductive system of helminthes. The hypocholesterolemic activity could be due to the increased fecal bile acid excretion as well as to the action of the short-chain fatty acids, coming from fermentation, on the synthesis of delta-aminolevulinic acid and by the increase of the cholesterol 7-alpha-hydroxylase and 3-hydroxy-3-methylglutaryl CoA reductase synthesis (Dall'Agnol and Von Poser, 2000).

Due to the effect related it may be possible that these fatty acids could act as an anthelmintic, although in the present study there was not observed differences between TM and UR extracts related to % of elimination in comparison one to another, although in comparison to the control group was evident a significative difference due to the TM group. Related to the obtained results due to the action of the TM extract it may be explained by their concentration as well as originated region which may explain the effect due to the biochemistry compounds in the equivalents proportions in spite of different conditions as soil composition, light and water availability.

The effect of TM extract may be support by possible modifications in ribosomal DNA spacer region suggesting that it could result in genetic and geographical variability as well as different bioactivity which may not be effective depend on the concentration of the extract (Arruda *et al.*, 2003).

Table 1: Anthelmintic activity of the extracts obtained of *Solanum lycocarpum* in the elimination of *Aspiculuris tetraptera* in mice naturally infected

Used parts	Administration form (%)	No. of animals	No. of Helminthes		
			Fecal exam	Necropsy	Elimination (%)
Leaves dried from três marias (TM)	10	10	61	1082	5.64 ± 3.33
Leaves died from UFRRJ (UR)	10	12	54	1717	3.15 ± 3.16
Nitroscanato (NIT)		12	499	282	64.00 ± 0.00
Mebendazol (MEB)		10	324	0.0	100.00 ± 0.00
Control (C)		10	45	2836	1.56 ± 3.16

We can speculate that the other effect would be related to the low concentration of steroidal alkaloid oligoglycosides which in a optimal concentration may suppress the transfer of sucrose from the stomach to the small intestine which could diminish the support of glucose to helminthes together with its antioxidant effect which is capable of reducing the nitrate generation which can be used in the protein synthesis.

CONCLUSION

Based on the results we can suggested that the anthelmintic effect of *Solanum lycocarpum*, TM extract, was observed related to the possible concentration of steroidal alkaloid oligoglycosides as well as the short-chain fatty acids presents in the extract. The similar action of the extracts may be explained by adaptation mechanisms related to the genetic and geographical variability.

ACKNOWLEDGMENT

We thanks to Thiago de Azevedo Amorim, technician of herbarium of the Botanical Department of Rural Federal University from Rio de Janeiro, by the identification of the studied species.

REFERENCES

- Amorim, A. and H.R. Borba, 1990. Action Anthelmintic III. Effect of the aqueous extracts of *Punica granatum* L. (romã) in the elimination of the *Vampirolepis nana* and de oxiurydeous in mice. Rev. Bras. Farmácia, 71: 85-87.
- Amorim, A., H.R. Borba, J.P.P. Carauta, D. Lopes and M.A.C. Kaplan, 1999. Anthelmintic activity of latex of *Ficus* species. J. Ethnopharmacol., 64: 255-258.
- Amorim, A., H.R. Borba and W.J. Silva, 1987. Action Anthelmintic of plants. Rev. Bras. de Farmácia, 68: 64-70.
- Arruda, M.C., M.A. Ferreira, R.N. Miller, M.L. Resende and M.S. Felipe, 2003. Nuclear and mitochondrial rDNA variability in *Crinipellis pernicioso* from different geographic origins and hosts. Mycol. Res., 107: 25-37.
- Dall'Agnol, R. and G.L. Von Poser, 2000. The use of complex polysaccharides in the management of metabolic diseases: the case of *Solanum lycocarpum* fruits. J. Ethnopharmacol., 71: 337-341.
- Gilioli, R., L.A.G. Andrade, L.A.C. Passos, F.A. Silva, D.M. Rodrigues and A.M.A. Guaraldo, 2000. Parasite survey in mouse and rat colonies of Brazilian laboratory animal houses kept under different sanitary barrier conditions. Arq. Bras. Med. Vet. Zootec., 52: 33-37.
- Perez, A.C., V. Franca, V.M. Jr. Daldegan and I.D. Duarte 2006. Effect of *Solanum lycocarpum* St. Hill on various haematological parameters in diabetic rats. J. Ethnopharmacol., 19: 442-444.
- Perec-Matysiak, A., A. Okulewicz, J. Hildebrand and G. Zalesny, 2006. Helminth parasites of laboratory mice and rats. Wiad Parazytol., 52: 99-102.
- Schwarz, A., E.C. Felipe, M.M. Bernardi and H.S. Spinosa, 2005a. Impaired female sexual behavior of rat offspring exposed to *Solanum lycocarpum* unripe fruits during gestation and lactation: Lack of hormonal and fertility alterations. Pharmacol. Biochem. Behav., 81: 928-934.
- Schwarz, A., M.R. Soares, J.C. Florio, M.M. Bernardi and H.S. Spinosa, 2005b. Rats exposed to *Solanum lycocarpum* fruit in utero and during lactation: Neurochemical, behavioral and histopathological effects. Neurotoxicol. Teratol., 27: 861-870.
- Silva, S.L.C., C.C. Alves, H.R. Borba, M.G. Carvalho and T.C. Bomfim, 2005. Evaluation of the anthelmintic activity of extracts from uxiemburgia octandra St. Hill. in mice naturally infected with *Aspiculuris tetraptera* and *Vampirolepis nana*. Rev. Bras. Parasitol. Vet., 14: 106-108.
- Vieira, G. Jr., P.M. Ferreira, L.G. Matos, E.C. Ferreira, W. Rodovalho, P.H. Ferri, H.D. Ferreira and E.A. Costa, 2003. Anti-inflammatory effect of *Solanum lycocarpum* fruits. Phytother. Res., 17: 892-896.