

Floristic structure and phytodiversity along an elevational gradient in Peepalkoti-Joshimath area of Garhwal Himalaya, India

Vishwapati Bhatt¹, Vijay Kant Purohit²

¹Department of Botany, Govt P. G. College, Gopeshwar, Chamoli-246 401, Uttarakhand, India

²High Altitude Plant Physiology Research Centre, HNB Garhwal University (A Central University), Srinagar (Garhwal), 246 1 74 Uttarakhand, India

vishwapati_bhatt@rediffmail.com, vijaykantpurohit@rediffmail.com.

Abstract: The present study was conducted in temperate Himalayan forests of Joshimath area in Chamoli district of Uttarakhand to understand the effect of altitudinal variation on structure and composition of the vegetation and to record the floristic diversity and economic utilities of the plants in the study area. Three altitudinal zones viz., upper zone (U) = 2000-2200m asl, middle zone (M) = 1800-2000m asl and lower zone (L) = 1600-1800m asl were selected for the study. In the present floristic survey the total of 74 families (72 Angiospermous and 2 Gymnospermous), 149 Genera (145 Angiospermous and 4 Gymnospermous) and 177 species (173 Angiospermous and 4 Gymnospermous) were recorded in the study area. Out of these 177 species identified in the study area 100, 47, 20 and 10 were herbs, shrubs, trees and climbers respectively. Rosaceae was the dominant family recorded with 16 species in the study area followed by the Asteraceae (15), Lamiaceae (11), Fabaceae (11) and Caryophyllaceae (5). In Ethnobotanical survey very useful information was recorded about the economic utility of the plants species present in the study area. Uses recorded were medicinal, fuel, fodder, edible and timber. Tree Species richness (SR) decreased from lower altitude to higher altitude. Species diversity (richness) and dominance (Simpson index) were found to be inversely related to each other. Tree density decreased from lower altitude to upper altitude, whereas TBC showed reverse trend. [Nature and Science. 2009;7(9):63-74]. (ISSN: 1545-0740)

Keywords: Phytosociology, Floristic composition, Diversity indices, Economic utility, Altitudes.

1. Introduction

The Indian Himalayan region occupies a special place in the mountain ecosystems of the world. These geodynamically young mountains are not only important from the stand point of climate and as a provider of life, giving water to a large part of the Indian subcontinent, but they also harbor a rich variety of flora, fauna, human communities and cultural diversity (Singh, 2006). The biodiversity which few years ago was considered unimportant by ecosystem ecologists, has now been shown to be significantly important for many aspects of ecosystem functioning. Diversity at all organizational levels, ranging from genetic diversity within populations to the diversity of ecosystems in landscapes, contributes to global biodiversity. The biodiversity has long been a source of amazement and scientific curiosity and increasingly a source of concern. Understanding of forest structure is a pre-requisite to describe various ecological processes and also to model the functioning and dynamics of forests (Elourard et al. 1997).

Species diversity has functional consequences, because the number and kinds of species present in any area determine the organismal traits, which influence ecosystem processes. The components of species diversity that determine the expression of traits include the number of species present (species richness), their relative abundance (species evenness), presence of the particular species (species composition), the

interactions among species (non-additive effects), and the temporal and spatial variation in these properties. In addition to its effects on current functioning of ecosystems, species diversity influences the resilience and resistance of ecosystems to environmental changes (Chapin et al. 2000).

The altitude and aspect play a key role in determining the temperature regime and atmospheric pressure of any site. Within one altitude the cofactors like topography, aspect, inclination of slope and soil type affect the forest composition (Shank and Noorie, 1950). The micro-environment of different aspects of hill slopes is influenced by the intensity and duration of available sunlight (Yadav and Gupta, 2006). This type of ecological knowledge is fundamental for conservation and sustainable utilization, and may provide important information for the policy makers for drafting management plans of fragile mountain ecosystems. Under the backdrop of the aforesaid facts, the present study was undertaken in temperate Himalayan forests of Peepalkoti-Joshimath area of Garhwal Himalaya, 1) to record plant species present in the study area along with their economic uses and 2) to understand the effect of altitude on the structure and composition of the vegetation of natural forests.

2. Material and Methods

The present study was conducted in temperate Himalayan forests of Joshimath area in Chamoli

district of Uttarakhand in year 2008. After the reconnaissance survey three altitudinal zones viz., upper zone (U) = 2200-2000m asl, middle zone (M) = 2000-1800m asl and lower zone (L) = 1800-1600m asl were identified to study the effect of altitudinal variation on structure and composition of the vegetation. The climate of the study area is typical temperate type. The year is represented by three main seasons; the cool and relatively dry winter (December to March); the warm and dry summer (mid-April to June); and a warm and wet period (July to mid-September) called as the monsoon or rainy season. The rainy season accounts for about three-quarters of the annual rainfall. Apart from these main seasons, the transitional periods interconnecting rainy and winter, and winter and summer are referred to as autumn (October to November) and spring (February to March). The mean annual rainfall was recorded as 1500mm and mean annual temperature between 5°C to 28°C.

The composition of the forest along the altitudinal gradient was analysed by using nested quadrat method or centre point quadrat method for trees, shrubs and herbs species as per Kent and Coker (1992). Three vegetation layers, (i.e., trees, shrubs and herbs) were analyzed for species richness, density and diversity. A total of 60 plots (twenty plots in each forest type) measuring 10m X 10m each were sampled. Trees (≥ 10 cm dbh) were analyzed by 10m x 10m sized quadrats, whereas shrubs by 5m x 5m sized quadrats. Further, quadrats of 1x1m size were randomly laid out with in each 10x10m sized quadrat at each site, to study plants in the herb layer. Circumference at breast height (cbh= 1.37m) was taken for the determination of tree basal area and was calculated as πr^2 , where r is the radius. Total basal area/cover is the sum of basal area/cover of all species present in the forest. The data were quantitatively analyzed for density, frequency and abundance following Curtis and McIntosh (1950). Species Richness was simply taken as a count of number of species present in that forest type. Basal area (m^2/ha) was used to determine the relative dominance of a tree species. Importance Value Index (IVI) was the sum of relative frequency, relative density and relative dominance (Phillips, 1959). The diversity (H') was determined by using Shannon-Wiener information index (Shannon and Weaver, 1963) as: $H' = - \sum n_i / n \log_2 n_i / n$; where, n_i was the IVI value of a species and n was the sum of total IVI values of all species in that forest type. The Simpson's concentration of dominance (Simpson, 1949) was measured as: $Cd = \sum P_i^2$, where, $\sum P_i = \sum n_i / n$, where, n_i and n are same as in Shannon-Wiener diversity

index. Simpson's diversity index (Simpson, 1949) was calculated as: $D = 1/Cd$, where, D = Simpson's diversity and Cd = Simpson's concentration of dominance. Species heterogeneity was calculated as under root of concentration of dominance (Cd).

To study the phytodiversity in the study area, regular field trips were undertaken in different seasons i.e., rainy, winter and summer, to collect the specimens of higher plants (Gymnosperms and Angiosperms). Identification of the specimens was done with the help of the existing Herbariums of Botany Department HNB Garhwal University (GUH), Forest Research Institute (DD) and Botanical Survey of India, Northern Circle (BSD). After identification, the enumeration of plants was done according to Bentham and Hooker's system of classification (1862-1883). The plants were divided into categories of common and uncommon according to their occurrence in the study area. An Ethnobotanical survey was also conducted in the villages nearby the study area to know the economic utility of various plant species encountered.

3. Results

Forest community structure and composition: Results of forest community structure and composition are given in tables 1 to 3.

Trees: At upper altitude *Cedrus deodara* was the dominant tree species with highest density (170 Ind/ha), TBC (98.82 m^2/ha) and IVI (155.96). At middle altitude *Pinus wallichiana* was the dominant tree species with highest density (180 Ind/ha), TBC (84.41 m^2/ha) and IVI (120.59). At lower altitude *Alnus nepalensis* was the dominant tree species with highest density (340 Ind/ha) and IVI (85.90), whereas highest TBC (3.78 m^2/ha) at this altitude was recorded for *Quercus semecarpifolia*. Tree Species richness (SR) decreased from lower altitude to higher altitude with highest SR at lower (19) altitude followed by middle (8) and upper (3) altitude. Highest (800 Ind/ha) tree density was recorded at lower zone followed by middle (600 Ind/ha) and lower (330 Ind/ha) altitudinal zone, where as highest (181.5 m^2/ha) TBC was recorded at upper altitude followed by middle (143.05 m^2/ha) and lower (9.63 m^2/ha) altitudes. Tree density decreased from lower altitude to upper altitude, whereas TBC showed reverse trend. Cd was found to be highest (0.4328) on upper altitude followed by middle (0.2561) and lower (0.1958) altitude whereas Simpson's diversity index showed reverse trend with highest (6.80) value at lower altitude followed by middle (6.74) and upper (2.57) altitude. Value H' was found to be highest (0.67) at upper altitude followed by middle (0.28) and lower (0.15) altitude.

Shrubs: At upper altitude *Rabdosia rugosa* was the dominant shrub species with highest density (520 Ind/ha) and TBC (0.3600 m²/ha), whereas highest IVI (82.38 m²/ha) at this altitude was recorded for *Corairia nepalensis*. At middle altitude *Rabdosia rugosa* was the dominant shrub species with highest density (680 Ind/ha), TBC (0.4310 m²/ha) and IVI (89.98). At lower altitude *Desmodium elegans* was the dominant shrub species with highest density (440 Ind/ha), TBC (0.1300 m²/ha) and IVI (70.27). Shrub Species richness (SR) decreased from lower altitude to higher altitude with highest SR at lower (22) altitude followed by middle (10) and upper (7) altitude. Highest (2420 Ind/ha) density was recorded at middle altitude followed by lower (2020 Ind/ha) and upper (1620 Ind/ha) altitudinal zone, where as highest TBC (1.21 m²/ha) was recorded at middle altitude followed by upper (0.75 m²/ha) and lower (0.39 m²/ha) altitudes. Cd was found to be highest (0.1996) on middle altitude followed by upper (0.1896) and lower (0.1138) altitude, whereas H' was found to be highest (0.17) at middle altitude followed by upper (0.14) and lower (0.06) altitude. Simpson's diversity index varied between 15.89 (lower altitude) to 7.81 (upper altitude).

Herbs: At upper altitude *Galium* sp. was the dominant herb species with highest density (15000 Ind/ha), TBC (0.0183 m²/ha) and IVI (54.36). At middle altitude *Geranium* sp. was the dominant herb species with highest density (28750 Ind/ha), TBC (0.0760 m²/ha) and IVI (70.52). At lower altitude *Pilea umbrosa* was the dominant herb species with highest density (16250 Ind/ha), TBC (0.0191 m²/ha) and IVI (44.31). Herb Species richness (SR) decreased from lower altitude to higher altitude with highest SR at lower (19) altitude followed by middle (16) and upper (7) altitude. Highest (174375 Ind/ha) density was recorded at middle altitude followed by lower (136250 Ind/ha) and upper (112500 Ind/ha) altitudinal zone, where as highest TBC (0.17 m²/ha) was recorded at middle altitude followed by lower (0.08 m²/ha) and upper (0.06 m²/ha) altitudes. Cd was found to be highest (0.0961) on middle altitude followed by upper (0.0777) and lower (0.0711) altitude, whereas H' was found to be highest (0.05) at middle altitude followed by upper

(0.03) and lower (0.02) altitude. Simpson's diversity index varied between 21.90 (middle altitude) to 18.93 (lower altitude).

Phytodiversity: In the present floristic survey the total of 74 families (72 Angiospermous and 2 Gymnospermous), 149 Genera (145 Angiospermous and 4 Gymnospermous) and 177 species (173 Angiospermous and 4 Gymnospermous) were recorded in the study area (table 4). Out of these 177 species identified in the study area 100, 47, 20 and 10 were herbs, shrubs, trees and climbers respectively. Rosaceae was the dominant family recorded with 16 species in the study area followed by the Asteraceae (15), Lamiaceae (11), Fabaceae (11) and Caryophyllaceae (5). Families with only one species were Agavaceae, Anacardiaceae, Aquifoliaceae, Araliaceae, Asclepidaceae, Berberidaceae, Betulaceae, Buxaceae, Cannabinaceae, Chenopodiaceae, Coriariaceae, Crassulaceae, Cucurbitaceae, Cuperasaceae, Cuscutaceae, Dioscoreaceae, Dipsacaceae, Elaeagnaceae, Ericaceae, Euphorbiaceae, Fabaceae, Gentianaceae, Geraniaceae, Hippocastanaceae, Hydrangeaceae, Juglandaceae, Lythraceae, Malvaceae, Meliaceae, Mimosaceae, Nictaginaceae, Orchidaceae, Oxalidaceae, Philadelphaceae, Phytolaccaceae, Plantaginaceae, Polygalaceae, Primulaceae, Rhamnaceae, Rutaceae, Saxifragaceae, Smilacaceae and Vitaceae. Families with two species were Boraginaceae, Brassicaceae, Campanulaceae, Caprifoliaceae, Onagraceae, Salicaceae, Thymelaeaceae, Ulmaceae, Urticaceae and Violaceae. Families with three species were Acanthaceae, Amaranthaceae, Araceae, Balsaminaceae, Cyperaceae, Moraceae, Oleaceae, Pinaceae, Rubiaceae, Scrophulariaceae and Solanaceae. Families with four species were Apiaceae, Hypericaceae, Poaceae, Polygonaceae and Ranunculaceae. In Ethnobotanical survey of the plant species present in the study area, very useful information was recorded about the economic utility of the plants. Uses recorded were medicinal, fuel, fodder, edible and timber and results are shown in the Table 4.

Table 1: Analytical characters for different forest types.

Trees	Density (Ind/ha)			TBC (m ² /ha)			IVI		
	U	M	L	U	M	L	U	M	L
<i>Aesculus indica</i>	-	40	-	-	3.41	-	-	19.58	-
<i>Alnus nepalensis</i>	30	120	340	0.28	3.76	0.96	25.91	33.15	85.80
<i>Cedrus deodara</i>	170	140	-	98.82	49.32	-	155.96	78.86	-
<i>Celtis australis</i>	-	-	30	-	-	0.35	-	-	15.72
<i>Lyonia ovalifolia</i>	-	40	50	-	1.20	1.26	-	18.03	27.67

<i>Pinus wallichiana</i>	130	180	-	82.40	84.41	-	118.13	120.59	-
<i>Populus ciliata</i>	-	-	40	-	-	0.84	-	-	22.06
<i>Pyrus pashia</i>	-	30	80	-	0.25	1.49	-	10.44	37.97
<i>Quercus semecarpifolia</i>	-	-	190	-	-	3.78	-	-	79.67
<i>Salix alba</i>	-	50	70	-	0.70	0.95	-	19.35	31.12
	330	600	800	181.50	143.05	9.63	300.00	300.00	300.00
Shrubs	U	M	L	U	M	L	U	M	L
<i>Berberis sp</i>	-	120	60	-	0.0040	0.0040	-	10.55	8.26
<i>Buddleja paniculata</i>	-	-	80	-	-	0.0100	-	-	10.80
<i>Corairia nepalensis</i>	340	440	-	0.3300	0.3100	-	82.38	59.61	-
<i>Cotoneaster baccularis</i>	-	60	-	-	0.0020	-	-	5.28	-
<i>Cotoneaster microphyllus</i>	40	-	60	0.0004	-	0.0100	8.40	-	9.81
<i>Daphne retusa</i>	-	-	80	-	-	0.0050	-	-	11.64
<i>Dapnae sp.</i>	100	140	-	0.0020	0.0050	-	35.85	14.09	-
<i>Desmodium elegans</i>	360	520	440	0.3600	0.4300	0.1300	75.82	72.84	70.27
<i>Deutzia compacta</i>	-	-	160	-	-	0.0200	-	-	23.73
<i>Elaeagnus conferta</i>	-	-	140	-	-	0.0600	-	-	28.82
<i>Lonicera quinquelocularis</i>	-	-	40	-	-	0.0030	-	-	4.88
<i>Princepia utilis</i>	-	60	100	-	0.0020	0.0200	-	7.91	16.50
<i>Rabdosia rugosa</i>	520	680	320	0.0500	0.4310	0.0500	47.55	89.98	39.40
<i>Rhamnus persica</i>	-	-	60	-	-	0.0040	-	-	8.26
<i>Rhamnus sp.</i>	40	60	-	0.0010	0.0040	-	20.25	8.07	-
<i>Rhamnus virgatus</i>	-	-	40	-	-	0.0020	-	-	4.62
<i>Rubus foliolosus</i>	-	-	40	-	-	0.0030	-	-	4.88
<i>Rubus niveus</i>	-	-	80	-	-	0.0050	-	-	11.64
<i>Sorbaria tomentosa</i>	120	200	240	0.0100	0.0200	0.0600	20.50	20.45	35.90
<i>Wikstroemia canescens</i>	100	140	80	0.0010	0.0020	0.0010	9.25	11.21	10.60
	1620	2420	2020	0.7544	1.2090	0.3870	300.00	300.00	300.00
Herbs	U	M	L	U	M	L	U	M	L
<i>Ajuga paviflora</i>	-	-	5000	-	-	0.0008	-	-	9.25
<i>Arisaema sp.</i>	2500	1875	-	0.0004	0.0002	-	5.49	3.44	-
<i>Artemisia capillaris</i>	-	-	2500	-	-	0.0004	-	-	5.40
<i>Bidens pilosa</i>	-	-	5000	-	-	0.0018	-	-	8.93
<i>Chenopodium album</i>	3750	-	-	0.0006	-	-	9.33	-	-
<i>Chenopodium sp.</i>	-	5000	2500	-	0.0016	0.0004	-	9.19	5.40
<i>Circium sp.</i>	-	3750	-	-	0.0011	-	-	6.09	-
<i>Circium verutum</i>	2500	-	-	0.0004	-	-	5.49	-	-
<i>Clinopodium sp.</i>	6250	5000	4375	0.0030	0.0016	0.0008	15.63	8.15	7.26
<i>Conyza japonica</i>	-	-	4375	-	-	0.0011	-	-	7.62
<i>Cynoglossum glochidium</i>	3750	7500	8750	0.0006	0.0046	0.0062	8.14	12.41	20.09
<i>Elsholtzia sp.</i>	5000	8750	-	0.0018	0.0062	-	12.48	12.01	-
<i>Eriophorum comosum</i>	-	-	6250	-	-	0.0029	-	-	11.18
<i>Fragarea sp.</i>	5000	-	-	0.0016	-	-	12.14	-	-
<i>Fragaria nubicola</i>	-	5625	-	-	0.0011	-	-	8.21	-
<i>Galium sp.</i>	15000	17500	10000	0.0183	0.0183	0.0046	54.36	27.54	20.61
<i>Geranium sp.</i>	11250	28750	-	0.0058	0.0760	-	27.42	70.52	-
<i>Hypericum elodeoides</i>	-	3750	-	-	0.0008	-	-	4.88	-
<i>Impatiens sp.</i>	6250	8750	7500	0.0029	0.0062	0.0050	16.65	14.08	17.72

<i>Lactuca sp.</i>	2500	2500	-	0.0004	0.0004	-	5.49	3.92	-
<i>Malva verticillata</i>	-	-	5000	-	-	0.0018	-	-	8.93
<i>Micromeria biflora</i>	-	-	6250	-	-	0.0023	-	-	11.99
<i>Origanum vulgare</i>	8750	11250	9375	0.0050	0.0103	0.0072	23.84	19.18	21.76
<i>Oxalis acetocella</i>	7500	23125	10000	0.0030	0.0220	0.0062	16.74	31.76	22.55
<i>Phytolacca acinosa</i>	5000	5000	-	0.0050	0.0026	-	17.91	9.77	-
<i>Pilea umbrosa</i>	-	-	16250	-	-	0.0191	-	-	44.31
<i>Pimpinella sp.</i>	3750	11250	8750	0.0006	0.0080	0.0046	9.33	17.80	18.15
<i>Plantago sp.</i>	-	2500	-	-	0.0002	-	-	3.80	-
<i>Polygonum sp.</i>	-	-	11250	-	-	0.0109	-	-	29.16
<i>Prunella vulgare</i>	7500	5000	5625	0.0050	0.0018	0.0018	21.33	8.27	12.46
<i>Salvia mocoocroftiana</i>	-	-	7500	-	-	0.0046	-	-	17.23
<i>Salvia sp.</i>	3750	3750	-	0.0006	0.0006	-	9.33	6.83	-
<i>Stellarea sp.</i>	5000	7500	-	0.0016	0.0029	-	13.34	11.40	-
<i>Thalictrum sp.</i>	-	2500	-	-	0.0002	-	-	3.80	-
<i>Viola sp.</i>	7500	3750	-	0.0023	0.0008	-	15.55	6.95	-
	112500	174375	136250	0.0589	0.1675	0.0825	300.00	300.00	300.00

Abbreviations: U= Upper altitude; M= Middle altitude; L= Lower altitude; TBC= Total Basal Cover; IVI= Importance Value Index.

Table 2: Diversity Indices of different forest types.

Trees	Cd			SDI			H'			Heterogeneity		
	U	M	L	U	M	L	U	M	L	U	M	L
<i>Aesculus indica</i>	-	0.0043	-	-	0.9957	-	-	0.00	-	-	0.07	-
<i>Alnus nepalensis</i>	0.0075	0.0122	0.0818	0.9925	0.9878	0.9182	0.0021	0.00	0.08	0.09	0.11	0.29
<i>Cedrus deodara</i>	0.2703	0.0691	-	0.7297	0.9309	-	0.4666	0.06	-	0.52	0.26	-
<i>Celtis australis</i>	-	-	0.0027	-	-	0.9973	-	-	0.00	-	-	0.05
<i>Lyonia ovalifolia</i>	-	0.0036	0.0085	-	0.9964	0.9915	-	0.00	0.00	-	0.06	0.09
<i>Pinus wallichiana</i>	0.1550	0.1616	-	0.8450	0.8384	-	0.2027	0.22	-	0.39	0.40	-
<i>Populus ciliata</i>	-	-	0.0054	-	-	0.9946	-	-	0.00	-	-	0.07
<i>Pyrus pashia</i>	-	0.0012	0.0160	-	0.9988	0.9840	-	0.00	0.01	-	0.03	0.13
<i>Quercus semecarpifolia</i>	-	-	0.0705	-	-	0.9295	-	-	0.06	-	-	0.27
<i>Salix alba</i>	-	0.0042	0.0108	-	0.9958	0.9892	-	0.00	0.00	-	0.06	0.10
	0.4328	0.2561	0.1958	2.5672	6.7439	6.8042	0.6715	0.28	0.15	1.00	1.00	1.00
Shrubs	U	M	L	U	M	L	U	M	L	U	M	L
<i>Berberis sp</i>	-	0.0012	0.0008	-	0.9988	0.9992	-	0.00	0.00	-	0.04	0.03
<i>Buddleja paniculata</i>	-	-	0.0013	-	-	0.9987	-	-	0.00	-	-	0.04
<i>Corairia nepalensis</i>	0.0754	0.0395	-	0.9246	0.9605	-	0.0688	0.03	-	0.27	0.20	-
<i>Cotoneaster baccularis</i>	-	0.0003	-	-	0.9997	-	-	0.00	-	-	0.02	-
<i>Cotoneaster microphyllus</i>	0.0008	-	0.0011	0.9992	-	0.9989	0.0001	-	0.00	0.03	-	0.03
<i>Daphne retusa</i>	-	-	0.0015	-	-	0.9985	-	-	0.00	-	-	0.04
<i>Dapnae sp.</i>	0.0143	0.0022	-	0.9857	0.9978	-	0.0057	0.00	-	0.12	0.05	-
<i>Desmodium elegans</i>	0.0639	0.0590	0.0549	0.9361	0.9410	0.9451	0.0536	0.05	0.04	0.25	0.24	0.23
<i>Deutzia compacta</i>	-	-	0.0063	-	-	0.9937	-	-	0.00	-	-	0.08
<i>Elaeagnus conferta</i>	-	-	0.0092	-	-	0.9908	-	-	0.00	-	-	0.10
<i>Lonicera quinquelocularis</i>	-	-	0.0003	-	-	0.9997	-	-	0.00	-	-	0.02
<i>Princepia utilis</i>	-	0.0007	0.0030	-	0.9993	0.9970	-	0.00	0.00	-	0.03	0.06
<i>Rabdosia rugosa</i>	0.0251	0.0900	0.0172	0.9749	0.9100	0.9828	0.0132	0.09	0.01	0.16	0.30	0.13
<i>Rhamnus persica</i>	-	-	0.0008	-	-	0.9992	-	-	0.00	-	-	0.03

<i>Rhamnus sp.</i>	0.0046	0.0007	-	0.9954	0.9993	-	0.0010	0.00	-	0.07	0.03	-
<i>Rhamnus virgatus</i>	-	-	0.0002	-	-	0.9998	-	-	0.00	-	-	0.02
<i>Rubus foliolosus</i>	-	-	0.0003	-	-	0.9997	-	-	0.00	-	-	0.02
<i>Rubus niveus</i>	-	-	0.0015	-	-	0.9985	-	-	0.00	-	-	0.04
<i>Sorbaria tomentosa</i>	0.0047	0.0046	0.0143	0.9953	0.9954	0.9857	0.0011	0.00	0.01	0.07	0.07	0.12
<i>Wikstroemia canescens</i>	0.0009	0.0014	0.0012	0.9991	0.9986	0.9988	0.0001	0.00	0.00	0.03	0.04	0.04
	0.1896	0.1996	0.1138	7.8104	9.8004	15.8862	0.1435	0.17	0.06	1.00	1.00	1.00
Herbs	U	M	L	U	M	L	U	M	L	U	M	L
<i>Ajuga paviflora</i>	-	-	0.0010	-	-	0.9990	-	-	0.00	-	-	0.03
<i>Arisaema sp.</i>	0.0003	0.0001	-	0.9997	0.9999	-	0.0000	0.00	-	0.02	0.01	-
<i>Artemisia capillaris</i>	-	-	0.0003	-	-	0.9997	-	-	0.00	-	-	0.02
<i>Bidens pilosa</i>	-	-	0.0009	-	-	0.9991	-	-	0.00	-	-	0.03
<i>Chenopodium album</i>	0.0010	-	-	0.9990	-	-	0.0001	-	-	0.03	-	-
<i>Chenopodium sp.</i>	-	0.0009	0.0003	-	0.9991	0.9997	-	0.00	0.00	-	0.03	0.02
<i>Cirsium sp.</i>	-	0.0004	-	-	0.9996	-	-	0.00	-	-	0.02	-
<i>Cirsium verutum</i>	0.0003	-	-	0.9997	-	-	0.0000	-	-	0.02	-	-
<i>Clinopodium sp.</i>	0.0027	0.0007	0.0006	0.9973	0.9993	0.9994	0.0005	0.00	0.00	0.05	0.03	0.02
<i>Conyza japonica</i>	-	-	0.0006	-	-	0.9994	-	-	0.00	-	-	0.03
<i>Cynoglossum glochidium</i>	0.0007	0.0017	0.0045	0.9993	0.9983	0.9955	0.0001	0.00	0.00	0.03	0.04	0.07
<i>Elsholtzia sp.</i>	0.0017	0.0016	-	0.9983	0.9984	-	0.0002	0.00	-	0.04	0.04	-
<i>Eriophorum comosum</i>	-	-	0.0014	-	-	0.9986	-	-	0.00	-	-	0.04
<i>Fragaria sp.</i>	0.0016	-	-	0.9984	-	-	0.0002	-	-	0.04	-	-
<i>Fragaria nubicola</i>	-	0.0007	-	-	0.9993	-	-	0.00	-	-	0.03	-
<i>Galium sp.</i>	0.0328	0.0084	0.0047	0.9672	0.9916	0.9953	0.0198	0.00	0.00	0.18	0.09	0.07
<i>Geranium sp.</i>	0.0084	0.0553	-	0.9916	0.9447	-	0.0025	0.04	-	0.09	0.24	-
<i>Hypericum elodeoides</i>	-	0.0003	-	-	0.9997	-	-	0.00	-	-	0.02	-
<i>Impatiens sp.</i>	0.0031	0.0022	0.0035	0.9969	0.9978	0.9965	0.0006	0.00	0.00	0.06	0.05	0.06
<i>Lactuca sp.</i>	0.0003	0.0002	-	0.9997	0.9998	-	0.0000	0.00	-	0.02	0.01	-
<i>Malva verticillata</i>	-	-	0.0009	-	-	0.9991	-	-	0.00	-	-	0.03
<i>Micromeria biflora</i>	-	-	0.0016	-	-	0.9984	-	-	0.00	-	-	0.04
<i>Origanum vulgare</i>	0.0063	0.0041	0.0053	0.9937	0.9959	0.9947	0.0017	0.00	0.00	0.08	0.06	0.07
<i>Oxalis acetocella</i>	0.0031	0.0112	0.0056	0.9969	0.9888	0.9944	0.0006	0.00	0.00	0.06	0.11	0.08
<i>Phytolacca acinosa</i>	0.0036	0.0011	-	0.9964	0.9989	-	0.0007	0.00	-	0.06	0.03	-
<i>Pilea umbrosa</i>	-	-	0.0218	-	-	0.9782	-	-	0.01	-	-	0.15
<i>Pimpinella sp.</i>	0.0010	0.0035	0.0037	0.9990	0.9965	0.9963	0.0001	0.00	0.00	0.03	0.06	0.06
<i>Plantago sp.</i>	-	0.0002	-	-	0.9998	-	-	0.00	-	-	0.01	-
<i>Polygonum sp.</i>	-	-	0.0094	-	-	0.9906	-	-	0.00	-	-	0.10
<i>Prunella vulgare</i>	0.0051	0.0008	0.0017	0.9949	0.9992	0.9983	0.0012	0.00	0.00	0.07	0.03	0.04
<i>Salvia mocroftiana</i>	-	-	0.0033	-	-	0.9967	-	-	0.00	-	-	0.06
<i>Salvia sp.</i>	0.0010	0.0005	-	0.9990	0.9995	-	0.0001	0.00	-	0.03	0.02	-
<i>Stellarea sp.</i>	0.0020	0.0014	-	0.9980	0.9986	-	0.0003	0.00	-	0.04	0.04	-
<i>Thalictrum sp.</i>	-	0.0002	-	-	0.9998	-	-	0.00	-	-	0.01	-
<i>Viola sp.</i>	0.0027	0.0005	-	0.9973	0.9995	-	0.0005	0.00	-	0.05	0.02	-
	0.0777	0.0961	0.0711	18.9223	21.9039	18.9289	0.0291	0.05	0.02	1.00	0.80	0.00

Abbreviations: U= Upper altitude; M= Middle altitude; L= Lower altitude; Cd= Simpson's Concentration of Dominance; SDI= Simpson's Diversity Index; H'= Shannon-Wiener Diversity Index.

Table 3: Total Diversity Indices of different forest types.

		Density (Ind/ha)	TBC (m ² /ha)	Cd	SWDI	H'	SR
Trees	Upper	330	181.50	0.4328	2.57	0.67	3
	Middle	600	143.05	0.2561	6.74	0.28	8
	Lower	800	9.63	0.1958	6.80	0.15	19
Shrubs	Upper	1620	0.75	0.1896	7.81	0.14	7
	Middle	2420	1.21	0.1996	9.80	0.17	10
	Lower	2020	0.39	0.1138	15.89	0.06	22
Herbs	Upper	112500	0.06	0.0777	18.92	0.03	7
	Middle	174375	0.17	0.0961	21.90	0.05	16
	Lower	136250	0.08	0.0711	18.93	0.02	19

Table 4: Details and uses of the plant species recorded in the Chamoli-Joshimath study area

Botanical Name	Family	Occurrence	Economic Utility	LF
<i>Abelia triflora</i>	Caprifoliaceae	UC	Fu	S
<i>Abies pindrow</i>	Pinaceae	C	Tm, Me	T
<i>Achyranthes bidentata</i>	Amaranthaceae	UC	-	H
<i>Adenocaulon himalaicum</i>	Asteraceae	C	Me	H
<i>Aesculus indica</i>	Hippocastanaceae	UC	Fo, Me	T
<i>Agave americana</i>	Agavaceae	C	Me	S
<i>Agrimonia pilosa</i>	Rosaceae	C	Me	H
<i>Albizia julibrissin</i>	Mimosaceae	UC	Me	T
<i>Alnus nitida</i>	Betulaceae	C	Fu	T
<i>Alpuda mutica</i>	Poaceae	C	Me	H
<i>Amaranthes viridis</i>	Amaranthaceae	C	Me	H
<i>Ammi majus</i>	Asteraceae	UC	-	H
<i>Anaphalis contrata</i>	Asteraceae	C	Me	H
<i>A. triplinervis</i>	Asteraceae	C	-	H
<i>Andropogon controtus</i>	Poaceae	C	-	H
<i>Androsace</i> sp.	Primulaceae	UC	-	H
<i>Anemone</i> sp.	Rosaceae	UC	Me	H
<i>A. vitifolia</i>	Rosaceae	UC	Me	H
<i>Aquilegia pubiflora</i>	Aquifoliaceae	UC	Me	H
<i>Arabis</i> sp.	Brassicaceae	C	Me	H
<i>Arctium lappa</i>	Asteraceae	C	Me	H
<i>Arenaria</i> sp.	Caryophyllaceae	C	-	H
<i>Arisaema</i> sp.	Araceae	C	-	H
<i>A. concinnum</i>	Araceae	UC	Me	H
<i>Artemisia capillaris</i>	Asteraceae	C	-	H

<i>A. roxburghiana</i>	Asteraceae	UC	Me	S
<i>Astragalus chlorostachys</i>	Fabaceae	C	Me	S
<i>Barleria cristata</i>	Acanthaceae	UC	Me	S
<i>Berberis angulosa</i>	Berberidaceae	C	-	S
<i>Bergenia ciliata</i>	Saxifragaceae	C	Me	H
<i>Boerhavia diffusa</i>	Nictaginaceae	C	Me	H
<i>Buddleja paniculata</i>	Scrophulariaceae	C	Fu	S
<i>Bupleurum falcatum</i>	Apiaceae	UC	Me	H
<i>Campanula</i> sp.	Campanulaceae	UC	-	H
<i>C. pallida</i>	Campanulaceae	UC	-	H
<i>Canabis sativa</i>	Cannabinaceae	C	Me	S
<i>Carex</i> sp.	Cyperaceae	C	-	H
<i>Cedrus deodara</i>	Pinaceae	UC	Tm, Me	T
<i>Celtis australis</i>	Ulmaceae	C	Ed, Fu	T
<i>Cerastrium</i> sp.	Caryophyllaceae	C	Fo	H
<i>Chenopodium album</i>	Chenopodiaceae	C	Ed	H
<i>Cichorium intybus</i>	Asteraceae	C	-	H
<i>Circium verutum</i>	Asteraceae	UC	Me	H
<i>Clematis connata</i>	Ranunculaceae	C	Me	C
<i>Clinopodium</i> sp.	Lamiaceae	C	-	H
<i>Corairia nepalensis</i>	Coriariaceae	C	Fu	S
<i>Cotoneaster baccilaris</i>	Rosaceae	C	Fu	S
<i>C. microphyllus</i>	Rosaceae	C	-	S
<i>Cupressus torulosa</i>	Cuperasaceae	UC	Tm, Me	T
<i>Cuscuta reflexa</i>	Cuscutaceae	C	Me	C
<i>Cyathula tomentosa</i>	Amaranthaceae	C	Me	S
<i>Cynodon dactylon</i>	Poaceae	C	Me	H
<i>Cynoglossum glochidiatum</i>	Boraginaceae	C	Me	H
<i>C. lanceolatum</i>	Boraginaceae	UC	Me	H
<i>Daphne retusa</i>	Thymelaeaceae	C	-	S
<i>Datura</i> sp.	Solanaceae	UC	Me	S
<i>Delphinium danudatum</i>	Ranunculaceae	UC	Me	H
<i>Desmodium elegans</i>	Fabaceae	C	Me, Fu	S
<i>D. multiflorum</i>	Fabaceae	C	Fu	S
<i>Deutzia compacta</i>	Hydrangeaceae	C	Fu	S
<i>Dioscorea deltoidea</i>	Dioscoreaceae	UC	Me	C
<i>Dipsacus mitis</i>	Dipsacaceae	UC	-	H
<i>Elaeagnus conferta</i>	Elaeagnaceae	C	Ed	S
<i>Elsholtzia</i> sp.	Lamiaceae	C	-	H

<i>E. fruticosa</i>	Lamiaceae	C	Fu	S
<i>E. flava</i>	Lamiaceae	C	Me	S
<i>Epilobium</i> sp.	Onagraceae	UC	-	H
<i>Erigeron</i> sp.	Asteraceae	UC	-	H
<i>Eriophorum comosum</i>	Cyperaceae	C	Fo	H
<i>Erysimum hieraciifolium</i>	Brassicaceae	UC	-	H
<i>Euphorbia</i> sp.	Euphorbiaceae	C	-	H
<i>Euphrasia himaliana</i>	Scrophulariaceae	C	-	H
<i>Fagopyrum dibotryis</i>	Polygonaceae	C	Ed	H
<i>Fallopia pterocarpa</i>	Polygonaceae	UC	-	H
<i>Ficus</i> sp.	Moraceae	UC	Ed, Fu	T
<i>F. hederacea</i>	Moraceae	C	Fo	C
<i>Fragaria nubicola</i>	Rosaceae	UC	Ed	H
<i>F. vestita</i>	Rosaceae	UC	Ed	H
<i>Fraxinus micrantha</i>	Oleaceae	C	Me	T
<i>Galium</i> sp.	Rubiaceae	C	-	H
<i>Geranium</i> sp.	Geraniaceae	C	Me	H
<i>Girardinia diversifolia</i>	Urticaceae	C	Me	S
<i>Hedera nepalensis</i>	Araliaceae	C	Fo	C
<i>Heracleum canascens</i>	Apiaceae	UC	Me	H
<i>Hypericum</i> sp.	Hypericaceae	UC	-	H
<i>H. perforatum</i>	Hypericaceae	UC	-	S
<i>H. elodeoides</i>	Hypericaceae	UC	-	H
<i>H. uralum</i>	Hypericaceae	UC	-	S
<i>Impatiens</i> sp.	Balsaminaceae	C	-	H
<i>I. falconerii</i>	Balsaminaceae	C	-	H
<i>I. sulcata</i>	Balsaminaceae	UC	Me	H
<i>Indigofera heterantha</i>	Fabaceae	C	Fu	S
<i>Inula cuspidata</i>	Asteraceae	UC	Me	H
<i>Jasminum</i> sp.	Oleaceae	UC	-	S
<i>J. humile</i>	Oleaceae	UC	Me	S
<i>Juglans regia</i>	Juglandaceae	UC	Ed, Me	T
<i>Kylinga</i> sp.	Cyperaceae	C	-	H
<i>Leptodermis lanceolata</i>	Rubiaceae	UC	Me	S
<i>Lespedeza gerardiana</i>	Fabaceae	C	Me	S
<i>L. juncea</i>	Fabaceae	C	Me	H
<i>Lonicera quinquelocularis</i>	Caprifoliaceae	UC	Fu	S
<i>Lotus corniculatus</i>	Fabaceae	UC	-	H
<i>Lyonia ovalifolia</i>	Ericaceae	UC	Me	T

<i>Malva verticilata</i>	Malvaceae	UC	Me	H
<i>Mentha longifolia</i>	Lamiaceae	UC	Me	H
<i>Micromeria biflora</i>	Lamiaceae	C	Me	H
<i>Morus serrata</i>	Moraceae	C	Me	T
<i>Nepeta</i> sp.	Lamiaceae	UC	-	H
<i>N. laevigata</i>	Lamiaceae	UC	-	H
<i>Oenothera rosea</i>	Onagraceae	UC	-	H
<i>Origanum vulgare</i>	Lamiaceae	C	Me	H
<i>Oxalis acetosella</i>	Oxalidaceae	C	Ed	H
<i>Paspalum paspalodes</i>	Poaceae	C	-	H
<i>Peristrophe paniculata</i>	Acanthaceae	UC	-	H
<i>Philadelphus tomentosus</i>	Philadelphaceae	UC	-	S
<i>Phytolacca acinosa</i>	Phytolaccaceae	C	Ed, Me	H
<i>Pimpinella</i> sp.	Apiaceae	UC	-	H
<i>Pinus wallichiana</i>	Pinaceae	UC	Tm, Me	T
<i>Plantago himalaica</i>	Plantaginaceae	C	Me	H
<i>Polygala</i> sp.	Polygalaceae	UC	-	H
<i>Populus ciliata</i>	Salicaceae	UC	Fo, Me	T
<i>Potentilla</i> sp.	Rosaceae	UC	Me	H
<i>Princepia utilis</i>	Rosaceae	C	Me, Fu	S
<i>Pteracanthus alatus</i>	Acanthaceae	C	-	S
<i>Pyrus pashia</i>	Rosaceae	C	Ed, Fu	T
<i>Quercus semecarpifolia</i>	Fagaceae	C	Fo, Me	T
<i>Rabdosia rugosa</i>	Lamiaceae	C	Fu	S
<i>Ranunculus</i> sp.	Ranunculaceae	UC	Me	H
<i>Rhamnus persica</i>	Rhamnaceae	UC	-	S
<i>R. virgatus</i>	Rhamnaceae	UC	Fu	S
<i>Rhus javanica</i>	Anacardiaceae	UC	Fo, Me	T
<i>Rosa brunonii</i>	Rosaceae	C	-	S
<i>Rosularia</i> sp.	Crassulaceae	UC	-	H
<i>Rubia cordifolia</i>	Rubiaceae	UC	Me	C
<i>Rubus ellipticus</i>	Rosaceae	C	Ed	S
<i>R. foliolosus</i>	Rosaceae	C	Ed	S
<i>R. prostrata</i>	Rosaceae	C	-	S
<i>Rumex hastatus</i>	Polygonaceae	C	Ed, Me	H
<i>R. nepalensis</i>	Polygonaceae	C	Me	H
<i>Salix alba</i>	Salicaceae	C	Fo	T
<i>Salvia</i> sp.	Lamiaceae	UC	-	H
<i>S. mcroftianna</i>	Lamiaceae	C	-	H

<i>Sarcococca saligna</i>	Buxaceae	C	-	S
<i>Saussurea albscens</i>	Asteraceae	C	-	S
<i>Sedum multicaule</i>	Crassulaceae	UC	-	H
<i>Selinum vaginatum</i>	Apiaceae	UC	Me	H
<i>Senecio chrysanthamoides</i>	Asteraceae	UC	-	H
<i>Silene</i> sp.	Caryophyllaceae	C	Fo	H
<i>S. edgeworthii</i>	Caryophyllaceae	C	-	H
<i>Smilax aspra</i>	Smilacaceae	UC	Me	C
<i>Solanum</i> sp.	Solanaceae	C	-	H
<i>S. nigrum</i>	Solanaceae	UC	Me	H
<i>Solena heterophylla</i>	Cucurbitaceae	UC	Ed, Fo	C
<i>Sorbaria tomentosa</i>	Rosaceae	C	Fu	S
<i>Spiraea canascens</i>	Rosaceae	C	Fu	S
<i>Spiranthes sinensis</i>	Orchidaceae	UC	Me	H
<i>Stellaria media</i>	Caryophyllaceae	C	-	H
<i>Swertia angustifolia</i>	Gentianaceae	UC	Me	H
<i>Tagetes minuta</i>	Asteraceae	C	-	H
<i>Thalictrum</i> sp.	Ranunculaceae	UC	Me	H
<i>Thymus linearis</i>	Lamiaceae	UC	Me	H
<i>Toona serrata</i>	Meliaceae	UC	Me	T
<i>Trifolium repens</i>	Fabaceae	C	-	H
<i>Trigonella corniculata</i>	Fabaceae	UC	Me, Ed	H
<i>Typhonium diversifolium</i>	Araceae	UC	-	H
<i>Ulmus villosa</i>	Ulmaceae	UC	Tm, Me	T
<i>Urtica dioica</i>	Urticaceae	C	Me	S
<i>Verbascum thapsus</i>	Scrophulariaceae	UC	Me	H
<i>Vigna</i> sp.	Fabaceae	UC	Fo	C
<i>Vincetoxicum hirundinaria</i>	Asclepidaceae	UC	Me	H
<i>Viola betonicifolia</i>	Violaceae	C	Me	H
<i>V. pilosa</i>	Violaceae	UC	Me	H
<i>Vitis</i> sp.	Vitaceae	C	-	C
<i>Wikstroemia canescens</i>	Thymelaeaceae	UC	-	S
<i>Woodfordia fruticosa</i>	Lythraceae	UC	Me	S
<i>Youngia</i> sp.	Asteraceae	UC	-	H
<i>Zanthoxylum armatum</i>	Rutaceae	UC	Me	S

Abbreviations: C= Climber; C= Common; Ed= Edible; Fo= Fodder; Fu= Fuel; H= Herb; LF= Life Form; Me= Medicinal; S= Shrub; T= Tree; Tm= Timber; UC= Uncommon.

4. Discussion

The diversity of trees is fundamental to total forest biodiversity, because trees provide resources and

habitat for almost all other forest species (Huang et al. 2003). At large scales, species diversity generally was found related to climate and productivity (Rahbek,

2005). Franklin *et al.* (1989) proposed that long-term productivity of natural forest ecosystems with high tree species diversity may be greater than that of forests with low diversity as a result of increased ecosystem resilience to disturbance. Slobodkin and Sanders (1969) opined that species richness of any community is a function of severity, variability and predictability of the environment in which it develops. Therefore, diversity tends to increase as the environment becomes more favourable and more predictable (Putman, 1994). Tree species diversity varied greatly from place to place mainly due to variation in biogeography, habitat and disturbance (Sagar *et al.* 2003), which have also been considered as the important factors for structuring the forest communities (Burslem and Whitmore, 1999). Srivastava *et al.* (2005) reported that the community characters differ among aspect, slope and altitude even in the same vegetation type. In our study we found that tree diversity decreased from lower altitude to higher altitude which means in our study area the environment at lower altitude was favourable for increasing tree diversity as compared to higher altitude.

In many other studies, the mean H' values for the other forests of temperate Himalaya varied from 0.4 to 2.8 (Singh *et al.* 1994), 0.08 to 1.29 (Shivnath *et al.* 1993) and 1.55 to 1.97 (Mishra *et al.* 2000), whereas in our study it varied between 0.67 to 0.15. Whittaker (1965) and Risser and Rice (1971) have reported the range of values of Cd for certain temperate vegetation from 0.19 to 0.99. The values of concentration of dominance (Cd) of the present study were more or less similar to the earlier reported values for temperate forests. Mean Cd values of 0.31 to 0.42 (Mishra *et al.* 2000) and 0.07 to 0.25 (Shivnath *et al.* 1993) were reported earlier from other parts of Indian Himalaya. The higher value of Cd in the forest growing on upper altitude was due to lower species richness. According to Baduni and Sharma (1997) the Cd or Simpson's index was strongly affected by the IVI of the first three relatively important species in a community. Species diversity (richness) and dominance (Simpson index) are inversely related to each other (Zobel *et al.* 1976).

The Himalayan region is bestowed with a variety of natural resources which have been exploited by mankind since time immemorial. The link between forest management and the well-being of communities in forested areas has traditionally been defined by forest sector employment opportunities (Sharma and Gairola, 2007). Ethnobotanical studies typically focus on recording the knowledge of traditional societies in remote places (Hodges and Bennett, 2006). Indigenous people have a vast knowledge of, and capacity for, developing innovative practices and products from

their environment. Indigenous knowledge grows from close interdependence between knowledge, land, environment and other aspects of culture in indigenous societies, and the oral transmission of knowledge in accordance with well understood cultural principles and rules regarding secrecy and sacredness that govern the management of knowledge (Tripathi *et al.* 2000). In the present study the traditional uses of various plant species by indigenous people have been recorded, which can be utilized in the future for technological advancement, economic prosperity and providing employment opportunity to the local people.

Acknowledgements:

Authors gratefully acknowledged the Head, Department of Botany, HNB Garhwal University, Srinagar, Directors, G. B. Pant Institute of Himalayan Environment and Development and High Altitude Plant Physiology Research Centre, Principal Govt. P.G. College Gopeshwar, and Prof. D. P. Vasistha, Department of Botany, HNB Garhwal University, Srinagar for encouragement and Support throughout the study periods.

*Correspondence to:

V. P. Bhatt, Ph.D., Assistant Professor
Department of Botany, Govt P. G. College, Gopeshwar
246401, Chamoli, Uttarakhand, India
vishwapati_bhatt@rediffmail.com,
bhattvp3@yahoo.com, Phone: 9412364460

References

- [1] Baduni NP and Sharma CM. Flexibility fitness compromise in some moist temperate forests of Garhwal Himalaya. *Annals of Forestry*, 1997;5:126-135..
- [2] Bentham G and Hooker JD. *Genera Plantarum*. London, 1862-1883; 3 Vols.
- [3] Burslem DF and Whitmore TC. Species diversity susceptibility to disturbance and tree population dynamics in tropical rain forest. *Journal of Vegetation Science*, 1999; 10: 767-776.
- [4] Chapin III, Erika FS, Zavaleta S, Eviner VT, Naylor RL, Vitousek PM, Reynolds HL, Hooper DU, Lavorel S, Sala OE, Hobbie SE, Mack MC and Diaz S. Consequences of changing biodiversity. *Nature*, 2000; 405: 234-242.
- [5] Curtis JT and McIntosh RP. Interrelations of certain analytical and synthetic phytosociological characters. *Ecology*, 1950; 31: 434-455.
- [6] Elourard C, Pascal JP, Pelissier R., Ramesh BR, Houllier F, Durand M, Aravajy S, Moravie MA and Gimaret-Carpentier C. Monitoring the structure and

dynamics of a dense moist evergreen forest in the Western Ghats (Kodagu District, Karnataka, India). *Tropical Ecology*, 1997; 38: 193-214.

[7] Franklin JF, Perry DA, Schowaltr ME, Harmon ME, Mckee A, Spies TA. Importance of ecological diversity in maintaining long-term site productivity. 1989; In: Maintaining the long-term productivity of Pacific Northwest forest ecosystems, Perry DA, Meurisse R, Thomas B, Miller R, Boyle J, Means J, Perry CR, Powers RF. (Eds.). Timber Press, Portland OR, pp. 82-97.

[8] Hodges S and Bennett BC. The ethnobotany of *Pluchea carolinensis* (Jacq.) G. Don (Asteraceae) in the Botanicas of Miami, Florida. *Economic Botany*, 2006; 60(1): 75-84.

[9] Huang W, Pohjonen V, Johansson S, Nashanda M, Katigula and Luvkkanen O. Forest structure, Species composition and diversity of Tanzanian rain forest. *Forest Ecology and Management*, 2003; 173: 11-24.

[10] Kent M and Coker P. Vegetation description and Analysis, 1992; Belhaven Press, London.

[11] Mishra A, Sharma CM, Sharma SD and Baduni NP. Effect of aspect on the structure of vegetation community of moist bhabar and tarai *Shorea robusta* forest in Central Himalaya. *Indian Forester*, 2000; 126(6): 634-642.

[12] Phillips EA. Methods of vegetation study. 1959; Henry Holt and Co. Inc., New York.

[13] Putman RJ. Community Ecology. 1994; Chapman & Hall, London.

[14] Rahbek C. The role of spatial scale and the perception of large-scale species-richness patterns. *Ecology Letters*, 2005; 8: 224-239.

[15] Risser PG and Rice EL. Diversity in tree species in Oklahoma upland forest. *Ecology*, 1971; 52: 876-880.

[16] Sagar R., Ragubanshi AS and Singh JS. Tree species composition, dispersion and diversity along a disturbance gradient in a dry tropical forest region of India. *Forest Ecology and Management*, 2003; 186: 61-71.

[17] Shank R.E and Noorie EN. Microclimate vegetation in a small valley in eastern Tennessee. *Ecology*, 1950; 11: 531-539.

[18] Shannon CE and Weaver W. The Mathematical Theory of Communication. University of Illinois Press, Urbana, Illinois, USA. 1963; p. 117.

[19] Sharma CM and Gairola Sumeet. Prospects of Carbon Management in Uttarakhand: An Overview. *Samaj Vigyan Shodh Patrika, Special Issue (Uttarakhand-1)*, 2007; 23-34.

[20] Shivnath S, Gupta K and Rajwar GS. Analysis of forest vegetation in a part of Garhwal Himalaya. *Recent Researches in Ecology, Environment and Pollution*, 1993; 6: 47-58.

[21] Simpson EH. Measurement of diversity. *Nature*, 1949; 163: 688.

[22] Singh JS. Sustainable development of the Indian Himalayan region: Linking ecological and economic concerns. *Current Science*, 2006; 90(6):784-788.

[23] Singh SP, Adhikari BS and Zobel DB. Biomass productivity, leaf longevity and forest structure in the central Himalaya. *Ecological Monograph*, 1994; 64: 401-421.

[24] Slobodkin LB and Sanders HL. On the contribution of environmental predictability to species diversity. *Brookhaven Symposia in Biology*, 1969; 22: 82695.

[25] Srivastava RK, Khanduri VP, Sharma CM and Pankaj Kumar. Structure, diversity and regeneration potential of Oak dominant conifer mixed forest along an altitudinal gradient of Garhwal Himalaya. *Indian Forester*, 2005; 131(12): 1537-1553.

[26] Tripathi S, Varma S and Goldey P. Using plants for health: indigenous knowledge in health care in a tribal region of Bihar, India. *International Journal of Sustainable Development and World Ecology*, 2000; 7:321-332.

[27] Whittaker RH. Dominance and diversity in land plant communities. *Science* 1965;147:250-60.

[28] Yadav AS and Gupta SK. Effect of micro-environment and human disturbance on the diversity of woody species in the Sariska Tiger Project in India. *Forest Ecology and Management*, 2006; 225: 178-189.

[29] Zobel DB, Mckee A, Hawk GM and Dyrness CT. Relationship of environment to the composition, structure and diversity of forest communities of the central western cascades of Oregon. *Ecological Monograph*, 1976; 46: 135-156.