

# Influence of Drying Methods, Extraction Time, and Organ Type on Essential Oil Content of Rosemary (*Rosmarinus officinalis* L.)

Khorshidi Jalal<sup>1,\*</sup>, Mohammadi Rahmat<sup>2</sup>, Fakhr Tabatabaei Mohammad<sup>1</sup>, Nourbakhsh Himan<sup>3</sup>

1. Department of Horticulture Science, University of Tehran, Karaj, Iran

2. Department of Biotechnology, University of Tehran, Karaj, Iran

3. Department of Food Science and Engineering, University of Tehran, Karaj, Iran

[jalal\\_khorshidi2006@yahoo.com](mailto:jalal_khorshidi2006@yahoo.com)

**Abstract:** Rosemary (*Rosmarinus officinalis* L.) is belonging to *Lamiaceae* family which grows wild in most Mediterranean countries. In order to investigate the influence of different methods of drying, extraction time, and type of organ on the essential oil percentage of this plant an experiment was carried out in college of agriculture of Karaj during the year 2009. Three drying methods investigated were oven drying (45°C), shade drying and sun drying. Four extraction times were: 1, 2, 3, 4 hours and three organ type were leaf, stem, mixed leaf and stem. Essential oil was obtained by water distillation method. Results showed that effect of drying methods, extraction time, and organ type on the essential oil percentage were significant. The maximum essential oil percentage (1.8%) was obtained to leaf sample, 3h of extraction, and shade drying. While the minimum essential oil percentage (0.12%) was obtained to stem sample, 1h of extraction, and oven drying. [Nature and Science. 2009;7(11):42-44]. (ISSN: 1545-0740).

**Key words:** *Rosmarinus officinalis* L.; Drying Methods; Extraction; Essential oil

## 1. Introduction

Rosemary (*Rosmarinus officinalis* L.) is a spice and medicinal herb widely used around the world. Of the natural antioxidants, rosemary has been widely accepted as one of the spices with the highest antioxidant activity (Wang et al, 2008., Peng et al, 2005., Oluwatuyi et al, 2004). Rosemary essential oil is also used as an antibacterial, antifungal and anticancer agent. Major constituents described for the oil are  $\alpha$ -pinene, 1, 8-cineole and camphor. The post-harvesting process of medicinal plants has great importance in the production chain, because of its direct influence on the quality and quantity of the active principles in the product sold. Drying has been one of the most important processes in pre-processing of agricultural products. Aromatic plants are often dried before extraction to reduce moisture content. The aim of drying is to reduce the moisture content of the product from actively growing in the field to a level that prevents deterioration of the product and allows storage in a stable condition. Proper drying of medicinal plants is fundamental to the achievement of a high quality product. A literature search was undertaken on effects of different methods of drying on essential oil content and chemical composition of the essential oil plants. The results showed that drying method had a significant effect on oil content and composition of aromatic plants (Okoh et al, 2008., Asekun et al, 2006., Ahmadi et al, 2008., Ronicely et al, 2008., Diana et al, 2008). Also duration of essential oil extraction affected on the quantity and quality of essential oil. Jamshidi et al. (2004) reported that essential oil percentage and essential oil component of fennel were affected by

duration of essential oil extraction. The objective of this study was to evaluate the influence of drying methods, extraction time, and type of organ on essential oil content of *Rosmarinus officinalis* L.

## 2. Materials and Methods

*Rosmarinus officinalis* L. were collected in college of agriculture of Karaj in during the year 2009 (longitude: 50° 59'E, Latitude: 35° 47'N, height of sea level (m): 1312/5, average annual rainfall (mm): 230, and mean annual temperature: 14.3 °C). The fresh plant materials were carefully separated into leaves, stems, and mixed of stems and leaves. Three drying methods investigated were oven drying, shade drying and sun drying. One of samples (100gr) was dried at room temperature and shade for 5 days. One of samples (100gr) was dried at sun for 3 days and other sample (100gr) was dried at oven (45°C) for 24 hours. This work performed for each sample with three replications. In order to extraction of essential oil, 100gr from each sample powdered and then essential oil isolated by water distillation for 1, 2, 3, 4 hour and to three replications. The essential oils were separated from the aqueous layer, dried over anhydrous sodium sulfate and calculated average of essential oil yield for three replications.

## 3. Results and Discussion

Results showed that drying methods, extraction time, and type of organ had a significant effect on the essential oil content of Rosemary. The maximum essential oil percentage (1/8%) was obtained in leaf,

shade drying, and 3h. The minimum essential oil percentage (0.12%) was obtained in stem, oven drying, and 1h. The samples that dried in shade had the higher essential oil as compared to samples that dried in sun and oven. Also leaf samples as compared to stem and mixed leaf and stem had the higher essential oil (Table 1). The oil content of shade-dried *Roman chamomil* flowers was found to be larger (1.9% w/w) than there of sun-dried (0.4%) and oven-dried at 40 °C (0.9%) (Omidbaigi et al, 2004). The drying method also had a significant effect on the proportion of the various components. It was also reported that the chemical composition, physical properties and antioxidant activities of yam flours were affected by drying methods to different extents. Yuan Zhang and Zhezhi Wang (2007) reported that in *Glechoma longituba* different drying methods caused some variation of the relative proportions of the components and the higher amount of germacrene D (19.0%) was obtained by shade-drying. Only sun drying brought about significant losses of the major compounds ( $\alpha$ -cadinol, germacrene B, germacrene D-4-ol, and  $\alpha$ -caryophyllene) in the essential oil when compared to the fresh plant material. Yuan Zhang and Zhezhi Wang showed that it could be concluded that drying of leaves of *G. longituba* under normal air and at room temperature conditions is most

suitable for a high-percentage of sesquiterpene, especially for germacrene D, but the oven-drying and silica gel-drying method are recommended for fast drying and similar components compared to the fresh plant material. Fatemeh Sefidkon et al. (2006) reported that the highest essential oil content of *Satureja hortensis* (1.06%) was obtained to oven-dried sample and the lowest essential oil content (0.87%) was obtained to sun-dried sample. In the *Mentha longifolia* L. only oven drying brought about significant losses of the major compounds (menthone, pulegone and 1,8-cineole) in the essential oil when compared to the fresh plant material (Asekun et al, 2007). The maximum essential oil percentage was obtained at 3h duration of extraction and with increasing duration of extraction not observed increasing in essential oil percentage. Barazandeh, M.M. (2005) reported that duration of essential oil extraction had significant effect on essential oil content in *Eucalyptus globules* and essential oil percentage was increased with the increasing of duration of extraction. Jamshidi et al. (2004) suggested that in *Foeniculum vulgare* Mill the highest essential oil percentage was obtained in 2.5hour.

Table 1. Effect of drying methods, extraction time, and type of organ on the essential oil content of *Rosmarinus officinalis* L.

Time (hour)	Shade drying			Sun drying			Owen drying		
	Leaf	Stem	Mixed and stem	Leaf	Stem	Mixed leaf and stem	Leaf	Stem	Mixed leaf and stem
1	0.9	0.21	0.7	0.8	0.19	0.5	0.7	0.12	0.4
2	1.3	0.3	0.9	1.2	0.26	0.7	1.1	0.22	0.7
3	1.8	0.45	1.3	1.62	0.35	1.1	1.5	0.3	0.9
4	1.8	0.46	1.3	1.63	0.36	1.1	1.5	0.3	0.9

#### 4. Conclusions

*Rosmarinus officinalis* L. is one of the most interesting research plants because it is between medicinal and aromatic plant. The post harvesting process (drying methods and extraction time) have a great importance in the production of essential oil and influenced on the quantity and quality of essential oil. Generally results showed that by shade drying, leaf samples and 3 hour a highest of essential oil percentage was obtained.

#### Acknowledgement

The authors are grateful to University of Tehran, Mr Kaveh Mollazade and Khaled Ahmad Aali for their valuable assistance and technical support.

#### Correspondence to:

Jalal Khorshidi

Department of Horticulture Science, University of Tehran, Iran

Cell phone: +98-914-816-5105

Email: [jalal\\_khorshidi2006@yahoo.com](mailto:jalal_khorshidi2006@yahoo.com)

#### References

- [1] Ahmadi K, Sefidkon F, Osareh MH. Effect of drying methods on quantity and quality of essential oil three genotype of *Rosa damascene* Mill. Iranian Journal of Medicinal and Aromatic Plants 2008; 24(2): 162-176.
- [2] Asekun OT, Grierson DS, Afolayan, AS. Influence of Drying Methods on the Chemical Composition and Yield of the Essential oil of *Leonotis Leonurus*. Journal Sci. Res 2006; 10: 61-64.
- [3] Asekun OT, Grierson DS, Afolayan AJ. Effects of drying methods on the quality and quantity of the essential oil of *Mentha longifolia* L. subsp. Capensis. Food chemistry 2007; 101: 995-998.
- [4] Barazandeh MM. Influence of extraction method and time of extraction on yield and chemical composition of *Eucalyptus globules*. Iranian Journal of Medicinal and Aromatic Plants 2005; 21: 75-93.
- [5] Diana L, Ronicely R, Evandro M, Evan V, Antonio P. Influence of drying air temperature on the essential oil content from *Melaleuca alternifolia* Cheel. International Conference of Agricultural Engineering 2008; Brazil, Aug 31 to Sep 4.
- [6] Jamshidi A, Ardakani M, Hajiakhundi A, Abdi Kh. Effect of extraction time on chemical composition of the essential oil of *Foeniculum vulgare*. Journal of Medicinal Plants 2004; 11: 68-73.
- [7] Okoh OO, Sadimenko AP, Asekun OT, Afolayan AJ. The effects of drying on the chemical components of essential oils of *Calendula officinalis* L. African Journal of Biotechnology 2008; 7(10): 1500-1502.
- [8] Oluwatuyi M, Kaatz, GW, Gibbons S. Antibacterial and resistance modifying activity of *Rosmarinus officinalis* L. Phytochemistry 2004; 65: 3249-3254.
- [9] Omidbaigi R, Sefidkon F, Kazemi F. Influence of drying methods on the essential oil content and composition of *Roman chamomile*. Flavour and Fragrance Journal 2004; 19: 196-198.
- [10] Peng Y, Yuan J, Liu F, Ye J. Determination of active components in rosemary by capillary electrophoresis with electrochemical detection. Journal of Pharmaceutical and Biomedical Analysis 2005; 39: 431-437.
- [11] Ronicely R, Evandro M, Lauril R, Luiz B, Jose M. Effect of drying air temperature upon the essential oil content from Guaco. International Conference of Agricultural Engineering 2008; Brazil, Aug 31 to Sep 4.
- [12] Sefidkon F, Abbasi Kh, Bakhshi Gh. Influence of drying and extraction methods on yield and chemical composition of the essential oil of *Satureja hortensis*. Food Chemistry 2006; 99: 19-23.
- [13] Wang W, Wu N, Zu Y, Fu Y. Antioxidative activity of *Rosmarinus officinalis* L. essential oil compared to its main components. Food Chemistry 2008; 108: 1019-1022.
- [14] Yuan Z, Zhezhi W. Influenced of drying methods on chemical composition of the essential oil of *Glechoma longitudo*. Chemistry of Natural Compounds 2007; 43 (5): 625-628.

8/10/2009