

Carotenoids Accumulation in the Green Alga *Scenedesmus* sp. Incubated with Industrial Citrate Waste and Different Inductions Stress

El-Sayed, A.B

Fertilization Technology Department, National Research Centre, Dokki, Cairo, Egypt
bokhair@msn.com

Abstract: A laboratory experiment was conducted in the algae-station of the Fertilization Technology Department, National Research Centre, Dokki, Cairo, Egypt with the green alga *Scenedesmus* sp. to study the effect of citrate waste in the nutrient solution on vegetative growth and pigments accumulation. The alga was incubated in 2000 ml culture with 10 ml citric acid waste characterized by high CNPK contents. After maximum growth rate was achieved induction was performed (by the day ten) and separate addition of NaCl, FeSO₄ and chelated FeSO₄ was performed and citrate volume was raised to be 50ml.l⁻¹. Growth measurements were dry weight, total chlorophyll and carotenoids. Results showed that maximum vegetative growth was reached after approximately 10 days of incubation. The maximum growth rate during this period was 0.29 and 0.26d⁻¹, while average growth rate was 0.14 and 0.16d⁻¹ for dry weight and total chlorophyll, respectively. Except of cultures received chelated FeSO₄, all other treated cultures (control, metal FeSO₄ and NaCl) resulted in complete degradation of chlorophyll and carotenoids massive accumulation. The cultures were completely greenish yellow and yellow colored. Maximum chlorophyll ratio was found in control cultures which received citrate only (50ml.l⁻¹), however maximum carotenoids ratio was detected in the algae supplied with metal FeSO₄, followed by 2% NaCl. [Nature and Science 2010;8(9):209-215]. (ISSN: 1545-0740).

Key words: Citrate wastes; Green algae; Vegetative growth; Caroteno

1. Introduction

Recycling of particulate organic wastes into marketable production matrix is highly recommended to meet the safe re-use and utilization of wastes daily increased due to the high industrial development (Stirling and Okumug, 1995). A promising technique in this field is the leaching with microorganisms. With this technique both energy requirements and environmental damage are kept low (Schinner and Burgstaller, 1989).

The use of autotrophic microorganisms is advantageous in this respect, because no organic carbon source is needed for their growth. On the other hand, heterotrophic microorganism can be used with higher pHs (alkaline, acid-consuming materials). Another advantage of leaching with heterotrophic microorganisms is the precipitations avoidance when high concentrations of metals and alkaline pHs are present because of the formation of complexes between metals and metabolites. At the same time, complication often reduces the toxicity of metal ions (Avakyan and Rabotnova 1971 & Babich and Stotzky 1980). Because of their capacity for oxidizing the substrate only partially and secreting it again in partly oxidized form, fungi are of particular interest for leaching. This "incomplete

oxidation" is strongly influenced by the composition of the medium and causes the accumulation of organic acids, which are able to extract metals from solid substrates. The most important primary metabolite for leaching is citric acid (Bosecker, 1987).

A pilot plant called 'Biocoil' has been set up in the UK and in Australia (Robinson, 1987). Their biocoil system is claimed to be used in conjunction with bacterial or algal growth to break down toxic wastes, to extract metals from liquids.

Different microorganisms, fungi, algae and higher plants are able to synthesize carotenoids. Production of secondary carotenoids from microalgae surpasses all other sources including yeast and fungi. Production of carotenoids from micro-algae for human consumption is of more interest during the last years,. However, more studies should be done to make it economically competitive (Bhosale, 2004, Pulz and Gross, 2004 and Spolaore ., *etal* 2006).

Factors affecting the bioaccumulation of such pigments were fully understood. Carotenoids accumulation by microalgae depend on both nutritional status (Cero'n., *et al* 2005 and Tittel,

2005) and environmental conditions, such as high light intensity (Bhosale, 2004), type of light (Janhke, 1999). In all cases, induced algae cells must be used by the log phase growth to avoid the dry weight failure. Also, shifting of photosynthetic metabolism to carotenoids accumulation by lipid biosynthesis should be considered (El-Shafey *et al.*, 1999).

In the present work, the green alga *Scenedesmus* sp. was used to study the effect of citrate wastes in the medium on both vegetative growth and carotenoids accumulation in the presence or absence of some induction factors.

2 - Materials and Methods

2.1. Alga and growth conditions

Polyethylene cylinders containing 2000 ml of nutrient growth media were used for incubating *Scenedesmus* sp. Growth media was composed of (g.l⁻¹) 0.56: 0.1:0.01 from urea, phosphoric acid and potassium sulphate; respectively plus 10 ml of citrate wastes. Illumination was provided by day light lamps (12x40w) reflexes from one side. Aeration was performed by compressed air from the upper hold throughout 3mm polyethylene tube. Chemical characteristics of the used waste are listed in Table 1.

Table 1. Chemical prosperities of citrate waste

pH	E.C dS.m ⁻¹	O.M	O.C	T.N	C:N ratio
4.87	4.95	0.80	0.47	0.65	0.73:1
P	K	Fe	Zn	Mn	Cu
%		ppm			
0.013	0.13	395	21	9	13

By the 10th day as cultures reached their maximum vegetative growth, induction growth was performed by discarding the upper clear layer of growth media and kept the sediment water algal bulk to the next induction growth. Induction growth media composed of tap enriched by 50 ml⁻¹ of citrate wastes with different induction factors including NaCl (2%) metal FeSO₄ and chelated FeSO₄ (Table 2). Sodium acetate was added in both growth periods at a final concentration of 45mM (per each) aiming at enhancing of chlorophyll formation during the vegetative growth period and decomposition as well as carotenoids bioaccumulation during induction.

Table 2. Induction factors used and their concentrations

Control	FeSO ₄	*Fe(chelate)	NaCl
50m.l ⁻¹ citrate wastes	50m.l ⁻¹ citrate wastes + 0.001 g.l ⁻¹ FeSO ₄	50m.l ⁻¹ citrate wastes + 0.001 g.l ⁻¹ chelated Fe	50m.l ⁻¹ citrate wastes + 2%NaCl

*Micronutrients suspension (3% Fe, Zn, Mn and 14%N)

2.2. Growth measurements

Daily measurement of dry weight was routinely carried out by filtering a defined volume of the algal slurry (5-10 ml) over pre-weighted dried membrane filter (0.45 μm). Filters were dried at 105°C for 30 minutes, kept over anhydrous calcium chloride till room temperature and then re-weighted. The difference between weights was recorded. .

Total chlorophyll was determined based on the methods described by Burnison 1980 on DMSO 95% extract. A modification was done by precipitating the algal biomass over membrane filter after dry weight determination. At 70°C, filters were soaked within 5-10ml of 95% DMSO, filtered and re-extracted if necessary. Chlorophyll absorbance was measured at 666nm and concentration was calculated according to Seely *et al.*, 1972. Carotenoids were determined based on the modification done by Boussiba *et al.*, 1992, Filters containing algal biomass were soaked into 5% (w/v) KOH in 30% (v/v) methanol, homogenized and heated in water bath to 70°C for 5 min. Supernatant was discarded and 5 drops of concentrated acetic acid were added to recover carotenoids and the residual was dissolved again by 5-10ml of 95% DMSO. Absorbance was measured at 468nm and concentration was calculated according to Davies 1976. Growth analysis was performed using the methods adopted by Pirt, 1973.

3. Result and Discussion

3.1. Vegetative growth

Citrate wastes were previously used to enhance vegetative growth of the green alga *Scenedesmus* sp. in NPK depleted media. Here, full media were used to obtain the biggest bulk which resist algal decay during induction period. Daily yield analysis showed that dry weight accumulation was obviously recorded their log phase by the day eleven (11). Chlorophyll content represented the same by the 10th day. Thus, the recommended retention time could be delivered by such time as a result of chlorophyll development curve. Otherwise, the relationship between chlorophyll and carotenoids defining the rate of nutrient consumption or biomass

accumulation, where low ratio explain the high growth rate which in turn monitored the high nutrient consumption rate with high dry weight accumulation.

As for growth analysis, specific growth rate (μ) showed the long exponential phase as dry weight accumulation, which in turn, explains the high suitability of such media to *Scenedesmus* sp due to its chemical composition or to the adaptation potential of this alga to the given nutritional conditions. Of this result, maximum growth rate by the third day was 0.29 d^{-1} and the calculated average was 0.13 d^{-1} . A slight decline in the growth rate was observed by the days 5 and 7, which explains the difference between both μ values (maximum and average). As for total chlorophyll, maximum accumulation rate was 0.27 d^{-1} with calculated average of 0.16 d^{-1} . The effect of citrate and other like-wastes on algal growth was considerably studied (Ruiz-Marin *et al.*, 2010). The main effect on growth enhancement could be attributed to the initial content from macro and micro-nutrients especially carbon and nitrogen.




Fig.1. a) Dry weight, b) total chlorophyll and c) growth rate of *Scenedesmus* sp. grown with 10 ml.l^{-1} of citrate wastes

3.2. Induction growth

Different hypothesis on carotenoids accumulation by green micro-algae were mentioned. Previous studies on carotenoids accumulation by *Scenedesmus*, *Chlorella* and *Haematococcus* showed that addition of at least 10% of nutrients mainly nitrogen are required to overcome the dry weight accumulation failure (El-Shafey 1999). Here, growth as dry weight was enhanced at first by the addition of citrate wastes even at the higher concentration. The main reason could be ascribed to the presence of organic carbon that allow fast carotenoids accumulation on the expense of carbohydrate metabolism or green pigments. Addition of extra citrate amount increased the salt potential of the growth medium, even though they are serve as utilizable nutrients as a source of different macronutrients including carbon, nitrogen, phosphorous, potassium and some micronutrients. Growth slightly delayed due to such high salt potential, but the growth; in general; might be goes back to the high initial biomass. Full BG-II medium which received the recommended NPK ratio when enriched by 10 ml.l^{-1} of citrate wastes represented the same afore-obtained result concerning dry weight and total chlorophyll. By the 10th day, all cultures were supplied by extra amount of citrate wastes (40 ml.l^{-1}). Different treatments including salinity (NaCl) and different iron forms represented wide results due to the kind of stress factor used.

When control cultures received only 50 ml^{-1} of citrate waste represented a slight dry weight decreases as compared with the time of citrate addition (following to the 10th day). The decrease on dry weight accumulation due to salting effect was accompanied by chlorophyll decomposition and carotenoids accumulation.

Dry weight: chlorophyll: carotenoids ratio was found as (1g: 19.37mg: 7.63mg) at the first cultivation time that explain the high initial biomass as dry weight or total chlorophyll. At the end of vegetative growth, such ratio was found as (1:17.06:13.31). Comparing both data showed the slight decrease on chlorophyll content against the increase on carotenoids content. The ratios here are not conditioned by the same rate of dry weight acceleration (Fig.1 and Tab. 3).

The ratio showed the drastic effect of NaCl on growth metabolites, where chlorophyll decreased and carotenoids increased. Under conditions of 2%NaCl, complete chlorophyll decomposition was observed by Boussiba *et al.*, 1992 with *Haematococcus pluvialis*. Here, the inhibitory effect on chlorophyll decomposition by NaCl could be ascribed to the presence of citrate wastes with its initial properties.

Cultures received 50 ml⁻¹ of citrate waste plus sodium chloride (2%) represented an extra inhibitory effect on dry weight as compared with citrate induced cultures. Chlorophyll decomposition was also engaged with extra carotenoids accumulation rate. The afore-mentioned ratio was found as (1:14.52:20.64) at the end of induction period. Sodium induced cultures which became nearly orange in color suggesting the extra salting-out potential that affect all the determined algal growth parameters including dry weight and pigments.

Table 3. Dry weight, total chlorophyll and carotenoids ratios of *Scenedesmus* sp grown under different carotenogenesis conditions with citrate wastes.

Treatments	Ratio	D.W	T.Ch	T.Car
Control	*V	1	19.37	7.63
	**C	1	17.06	13.31
NaCl (2%)	*V	1	19.37	7.63
	**C	1	14.52	20.64
FeSO ₄	*V	1	19.37	7.63
	**C	1	3.57	44.9
Fe-Chelate	*V	1	19.37	7.63
	**C	1	12.14	37.45

*V = ratio at the end of vegetative growth

**C= ratio at the end of induction growth

The effect of salinity against growth metabolism is fully established by many investigations even in halotolerant algal species such as *Dunaliella*. In *Haematococcus pluvialis*, carotenoids accumulation (astaxanthin) was triggered by the addition of 2% of NaCl to growth medium (Boussiba *et al.*, 1992). Furthermore, the biomass of *Botryococcus braunii* was increased with increasing concentration of sodium chloride and maximum biomass was achieved in 17mM and 34mM salinity, while phosphate decrease was observed due to its utilization by the alga (Ranga Rao *et al.*, 2007). During haloadaptation of such alga their hydrocarbon profile was highly increased while decreases in protein content and biomass yield. Carbohydrates and lipids remain unchanged may be due to the decrease in photosynthetic rate (Hart *et al.*, 1991) or probably due to less adaptability of the organism to high salinity (Vazquez-Duhalt and Arredondo-Vega 1991 and Ben-Amotz *et al.* (1985).

Ferrous sulfate induced cultures were drastically changed and the fast yellow color was observed by the fourth day of induction period. The fast appearing of visual carotenoids suggesting the rapid chlorophyll decomposition, however dry weight was

slightly inhibited. The main reason for such results could be ascribed to the effect of ferrous ion on active hydroxyl group (active oxygen species). The ferrous form of Fe in particular is known to give rise to free radical formation (especially hydroxyl radicals, HO^{*}) via Fenton chemistry. Radical initiation of cellular processes via signal cascade systems is thought to be an integral part of many biological systems and it was suggested that free radicals may play a role in the accumulation of β -carotene in *Dunaliella* (Ben-Amotz & Avron, 1983) and astaxanthin in the yeast *Paffia rhodozyma* (Schroeder & Johnson, 1995). Kobayashi *et al.* (1993) reported that increased astaxanthin formation caused by Fe⁺² was inhibited by potassium iodide which scavenges HO^{*}, suggesting the HO^{*} formed by an iron-catalyzed Fenton reaction is required for enhanced astaxanthin biosynthesis in *Haematococcus pluvialis*. As observed previously by Droop (1954) and Borowitzka *et al.* (1991), acetate- at small quantities- appeared to be an important carbon source, enhancing both growth and carotenogenesis. However, the effect of acetate was concentration dependent. Higher concentrations inhibiting growth, but markedly increasing astaxanthin content per cell. Acetate addition in excess may generate a relative shortage of nitrogen inducing cyst formation and astaxanthin accumulation triggered by a high carbon/nitrogen (C/N) ratio (Kakizono *et al.*, 1992). The algal cells seem to reduce their nitrogen uptake and begin to use the cellular nitrogen as in typical N-deficiency, although nitrogen exists in the culture medium. Citrate wastes at high concentration (50 ml.l⁻¹) support growth media by high organic carbon and other organic acids which stimulate both vegetative growth and carotenoids accumulation. As mentioned before, in the absence of essential nutrients including nitrogen and phosphorous, increasing of acetate concentrations also rise the salinity potential of the growth media.

Comparing the ratio of dry weight: total chlorophyll: carotenoids showed the highly drastic effect of metal Fe on cell metabolism, where it was found as (1: 3.57: 44.9). (Tab. 3 and Fig.2) These Results mean that complete chlorophyll decomposition and extra accumulation of carotenoids are possibility happened. As mentioned before, the stability of dry weight could be attributed to the effect of extra citrate addition.

Cultures received 50 ml⁻¹ citrate waste with chelated Fe represented a moderate response and the rate of dry weight accumulation surpass all other cultures except those of control cultures. Chlorophyll decomposition was slightly inhibited as well as inhibition of carotenoids accumulation was

occurred. The massive effect could be oriented as Fe, Na, Fe-chelate and control. It is well known that chelating prevent the free charge of ions and make it easy to penetrate the life cells and also used to remove toxicity of heavy metals even in human body.




Fig.2. a) Dry weight, b) total chlorophyll and c) growth rate (dry weight and total chlorophyll) of *Scenedesmus sp.* grown on 50ml.l⁻¹ of citrate wastes with different induction factors.

Thus, it may be concluded that chelating prevent the high solubility of ferrous in which became slow released ion and delay the free radical generation. In this context, when the chelated iron form was used in this action, no Fenton reaction could be ascribed as mentioned by Borowitzka *et al.*, 1991, where he stated that when EDTA-chelated FeCl₃.6H₂O did not produce any significant differences in astaxanthin

formation when added at three different concentrations. Our data could be supported by what obtained by El-Sayed 1999, where, FeCl₃ triggered the carotenoids accumulation with *Haematococcus pluvialis*, *Scenedesmus sp.* and *Chlorella sp.* It can be also mentioned that EDTA and other chelating agents neutralize the ionic charges of the used ions which in turn prevent the photo reaction produce free radicals. The afore-obtained ratio found as (1: 12.14:37.14) explains the inhibitory effect of EDTA against drastic chlorophyll decomposition. Concerning the chelated Fe form used, it is accompanied with the same content of Zn, which minimizes the toxicity of extra Fe addition and also contain 14% of N delivered from urea that enhance vegetative growth and support growth medium by alternative carbon source (El-Sayed *et al.*, 2001).

Nitrate concentration played a very important role in the cell division rate and in the accumulation of secondary carotenoids of *H. pluvialis* (Boussiba and Vonshak, 1991). This suggests that the synthesis of astaxanthin requires nitrogen, and most likely reflects the need for continuous synthesis of protein in order to support the massive accumulation of pigments. Nitrogen starvation is an effective way to enhance astaxanthin accumulation in *Haematococcus*. The optimum nitrate concentration to avoid this problem seems to be 0.15 g/l (Orosa *et al.*, 2005). The ratio chlorophyll a/total carotenoids could be a good indicator of the physiological status of the culture. Boussiba and Vonshak, 1991 added that in favorable growth conditions, this value was about 4; indicating nutrient replete conditions; however, a value below 3 implied limitation of growth by reduced nutrient supply or another kind of stress. Thus, it is possible to quickly check the N status of the cell as an indicator to astaxanthin accumulation.

4. Conclusion

As it supports growth media by organic carbon and several nutrients, utilizing of citrate wastes for enhancement of algae vegetative growth is very advisable to reduce production cost and enhancement of the induction growth. In addition, the induction effect suggested to mainly factoring kind dependent rather than the salt concentration.

Acknowledgment

This work was carried out as a part of the activities of the Egypto-German Project "Micronutrients and other Plant Nutrition Problems in Egypt" conducted by National Research Centre, Cairo (Coordinator: Prof. Dr. M.M. El-Fouly) and the Institute of Plant Nutrition, Technical University of Munich (Prof. Dr. A. Amberger).

Corresponding Author

El-Sayed, A.B

Fertilization Technology Department, National Research Centre, Dokki, Cairo, Egypt
bokhair@msn.com**5. References**

1. Aksu, Z. (1988). Biosorption of heavy metals by microalgae in batch and continuous systems. in: Wong, Y.S. and Tam, N.F.Y. ed(s),. *Algae for Waste Water Treatment*. Springer-Verlag and Landes Bioscience, p. 37–53.
2. Avakyan, Z. A., and Rabotnova, I. L. (1971). Comparative toxicity of free ions and complexes of copper with organic acids for *Candida utilis*. *Microbiology* 40:262-266.
3. Babich, H., and Stotzky, G. (1980). Environmental factors that influence the toxicity of heavy metal and gaseous pollutants to microorganisms. *Crit. Rev. Microbiol.* 8:99-145.
4. Ben-Amotz, A. & Avron, M. (1983). On the factors which determine massive β -carotene accumulation in the halotolerant alga *Dunaliella bardawil*. *Plant Physiol.*, 72,.
5. Ben-Amotz, A.; Tornabene, T.G. and Thomas, W.H. (1985). Chemical profile of selected species of microalgae with special emphasis on lipids. *J. Phycol.* 21, 72–81.
6. Bhosale, P. (2004). Environmental and cultural stimulants in the production of carotenoids from microorganisms. *Appl. Microbiol Biotechnol.*;63:351 – 61.
7. Borowitzka, M.A.; Huisman, J.M. and Osborn, A. (1991). Culture of the astaxanthin-producing green alga *Haematococcus pluvialis*. I. Effects of nutrients on growth and cell type. *J. Appl. Phycol.* 3, 295–304.
8. Bosecker, K. (1987). Laugung lateritischer Nickelerze mit heterotrophen Mikroorganismen. *Acta Biotechnol.* 7:389-399.
9. Boussiba, S. and Vonshak, A. (1991). Astaxanthin accumulation in the green alga *Haematococcus pluvialis*. *Plant Cell Physiol.*, 32, 1077-1082.
10. Boussiba, S.; Fan, L. and Vonshak, A. (1992). Enhancement and determination of astaxanthin accumulation in green alga *Haematococcus pluvialis*. *Methods in Enzymology*, 213, Carotenoids Part A, Lester Packer (ed.), Academic Press: 386-371
11. Burnison, B. K. (1980). Modified dimethyl sulfoxide (DMSO) extraction for chlorophyll analysis of phytoplankton. *Can. J.Fish. Aquat. Sci.*, 37:729-733.
12. Cero'n, M.C.; Sa'nchez, A.; Ferna'ndez, J.M.; Molina, E., and Garc'a, F. (2005). Mixotrophic growth of the microalga *Phaeodactylum tricorutum*. Influence of diferent nitrogen and organic carbon source on productivity and biomass composition. *Proc Biochem* 2005;40:297 – 305.
13. Crist, H.R.; Oberholser, K.; Shank, N. and Nguyen, M. (1981). Nature of bonding between metallic ions and algal cell walls. *Environ Sci. Technol.*;15:1212–7.
14. Davis, B. H. (1976). Carotenoids. In *Chemistry and Biochemistry of Plant Pigments*. 2nd Eddition, vol.2. ed. Goodwin. T. W. pp.38-165. Academic Press.
15. Droop, M.R. (1954). Conditions governing haematochrome formation in *Haematococcus pluvialis*. *Arch. Mikrobiol.* 20, 391.
16. Durrer, H., 1986. Coloration. In: Bereiter-Hahn, J., Matolsky, A.G., Richards, K.S. (Eds.), *Biology of the Integument*, vol. 2. Springer-Verlag, Berlin, Germany, pp. 239–247.
17. El-Sayed, A.B. (1999). Some physiological studies on green algae. Ph.D. Thesis, Faculty of Agric., Department of Agriculture Botany, Cairo University.
18. El-Sayed, A. B. (2004). Circulation of Quaron Lake wastes. II-Growth of *Scenedesmus* sp. under Mg residences. *Egypt. J. Biotechnol.* Vol.17, 477-485, June., 2004.
19. El-Sayed, A.B.; F.E. Abdalla and A.A. Abdel-Maguid (2001). Use of some commercial fertilizer compounds for *Scenedesmus* cultivation. *Egyptian J. Phycology*, Vol.2:9-16.
20. El-Shafey, Y.H.; El-Fouly, M.M.; Khalil, M.M.; Abdallah, F.E. and El-Sayed, A.B. (1999). Secondary carotenoids accumulation by some green algae species. *The First Congress on the Recent Technologies in Agriculture*, 27-29 Nov., 1999; Faculty of Agriculture , Cairo University, Egypt.
21. Garbayo, I.; Cuaresma, M.; lchez, C.V. and Vega, J.M.(2008). Effect of abiotic stress on the production of lutein and b-carotene by *Chlamydomonas acidophila*. *Process Biochemistry* 43,1158 – 1161
22. Gardea-Torresdey, J.L.; Becker-Hapak, M.K.; Hosea, J.M. and Darnall, D.W. (1990). Effect of chemical modification of algal carboxyl groups on metal ion binding. *Environ Sci Technol.*;24:1372–8.
23. Hart, B.T.; Bailey, P.; Edwards, R.; Hortlek, K.; James, K.; McMohan, A.; Meredith, C. and Swading, K. (1991). A review of the salt sensitivity of the Australian fresh water biota. *Hydrobiologia* 210, 105–144.
24. Hasimoto, S. and Furakawa, K. (1989). Nutrient removal from secondary effluent by filamentous

- algae. *J. Ferment. Bioeng.*, 67(1), 62-69.
25. Inborr, J. and Lignell, Å. (1997). Effect of feeding astaxanthin-rich algae meal (*Haematococcus pluvialis*) on performance and carotenoid concentration of different tissues of broiler chickens. In: Proceedings of the XIII WPSA Conference on Poultry Meat Quality in Poznan, Poland, September 22–25, Session M1, pp.39–43.
 26. Janhke, LS. (1999). Massive carotenoid accumulation in *Dunaliella bardawil* induced by ultraviolet-A radiation. *J Photochem Photobiol B* 1999;48:68 - 74.
 27. Kakizono, T.; Kobayashi, M. and Nagai, S. (1992). Effect of carbon/nitrogen ratio on encystment accompanied with astaxanthin formation in a green alga, *Haematococcus pluvialis*. *J. Ferment. Bioeng.* 74, 403 - 405.
 28. Kobayashi, M.; Kakizono, T. and Nagai, S. (1991). Astaxanthin production by a green alga, *Haematococcus pluvialis* accompanied with morphological changes in acetate media. *J. Ferment. Bioeng.* 71:335-339.
 29. Kobayashi, M.; Toshihide, K. and Nagai, S. (1993). Enhanced carotenoid biosynthesis by oxidative stress in acetate-induced cyst cells of a green unicellular alga *Haematococcus pluvialis*. *Appl. Environ. Microbiol.*, 59, 867-873.
 30. Martinez, M. E; Jimenez, J. M. and El-Yousfi, F. (1999). Influence of phosphorous concentration and temperature on growth and phosphorous uptake by the microalga *Scenedesmus obliquus*. *Bioresource Biotechnology*, vol. 67, pp233-240.
 31. Martinez, M. E; Sanchez, S. Jimenez, J. M., El-Yousfi, F. and Mounz, L.(2000). Nitrogen and phosphorous removal from urban wastewater by the microalga *Scenedesmus obliquus*. *Bioresource Biotechnology*, vol.73, pp263-272.
 32. Orosa, M.; Franqueira, D.A. and Abalde, C.J.(2005). Analysis and enhancement of astaxanthin accumulation in *Haematococcus pluvialis*.*Bioresource Technology* 96 (2005) 373 -378
 33. Pirt, S.J. (Ed.) (1973). Principle of Microbe and Cell Cultivation. Blackwell Scientific Publication, pp: 4-7.
 34. Pulz, O. and Gross, W. (2004). Valuable products from biotechnology of microalgae. *Appl. Microbiol Biotechnol.*; 65:635 - 48.
 35. Ranga Rao, A.; Dayananda, C.; Sarada, R.; Shamala, T. R. and Ravishankar, G.A. (2007). Effect of salinity on growth of green alga *Botryococcus braunii* and its constituents *Bioresource Technology* 98560–564
 36. Robinson, L.F. (1987). Improvements relating to biomass production. European Patent 0,239,272.
 37. Ruiz-Marin, A.; Mendoza-Espinosa, L.G. and Stephenson, T. (2010). Growth and nutrient removal in free and immobilized green algae in batch and semi-continuous cultures treating real wastewater. *Bioresource Technology* 101, 58–64.
 38. Schinner, F. and Burgstaller, W. (1989). Extraction of Zinc from Industrial Waste by a *Penicillium* sp. *Applied and Environmental Microbiology*, 55, 5, 1153-1156.
 39. Schroeder, W. A. and Johnson, E. A. (1995). Singlet oxygen and peroxy radicals regulate carotenoid biosynthesis in *Phaffia rhodozyma*. *J. Biol Chem.*, 270, 18374-18379.
 40. Seely, G. R.; Doncan, M. J. and Widaver, W. E. (1972). Preparative and analytical extraction of pigments from brown algae with dimethyl sulfoxide. *Mar. Biol.*, 12: 184188.
 41. Spolaore, P.; Joannis-Cassan, C.; Duran, E. and Isambert A.(2006). Commercial applications of microalgae. *J Biosci. Bioeng.*; 101:87 - 96.
 42. Stirling, H. P. and Okumug, I. (1995). Growth and production of mussels (*Mytilus edulis* L.) suspended at salmon cages and shellfish farms in two Scottish sea lochs. *Aquaculture* 134 (1995) 193-210
 43. Tittel, J.; Bissinger, V.; Gaedke, U. and Kamjunke, N. (2005). Inorganic carbon limitation and mixotrophic growth in *Chlamydomonas* from an acidic mining lake. *Protist.*;156:63 - 75.
 44. Vazquez-Duhalt, R. and Arredondo-Vega, B.O. (1991). Halo-adaptation of the green alga *Botryococcus braunii* ('A' Race). *Phytochemistry* 30 (9), 2919–2925.
 45. Wilde, E.W. and Benemann, J.R. (1993). Bioremoval of heavy metals by the use of microalgae. *Biotech. Adv.*;11:781–812.

7/21/2010