

Evaluation of pattern of shell utilization of the hermit crab *Clibanarius africanus* (Aurivillus, 1898) (Decapoda: Diogenidae) in the Lagos lagoon: an example of bioecological relationship in benthos.

Aderonke Lawal-Are, Roland Efe Uwadiae* and Olayemi Ruth Owolabi

Department of Marine sciences, University of Lagos, Akoka , Yaba, Lagos, Nigeria

*Corresponding author: eferoland@yahoo.com. Tel: +2348063145723

Abstract. Evaluation of pattern of shell utilization by *Clibanarius africanus* was investigated in the intertidal area of the Lagos lagoon, between March and August, 2008 at five study sites. A total of 663 specimens of hermit crab in gastropod shells were collected. Shell occupation of *C. africanus* was limited to gastropod shells belonging to *Pachymelania* and *Tympanotonus* spp. The gastropod shell most inhabited was those of *Pachymelania* spp, which accounted for 60.96% of inhabited shells, while about 39% of the shell inhabited belonged to *Tympanotonus* spp. Shell preferences were characterized by shell length, weight and aperture width. Positive and statistically significant correlations were obtained between morphometric characteristics of *C. africanus* and those of the shells inhabited, suggesting that fitness of shell to crab dimension constitutes mainly the determinant for *C. africanus* shell utilization. Spatiotemporal variations in the type of shell occupied were not significant in this study. Analysis of the abundance of *C. africanus* in the study area indicates that, a relatively higher abundance of *C. africanus* was observed in site 3, due probably to the favourable environmental conditions provided by large percentage of sand fractions in the sediment. From the data recorded in this study, it may be concluded that shell utilization by hermit crabs involves individual preferences related to the shell features that best provide protection, survival and opportunity for the enhancement of behavioural attributes that are necessary for the maximization of bio-ecological relationships.

[Aderonke Lawal-Are, Roland Efe Uwadiae and Olayemi Ruth Owolabi. Evaluation of pattern of shell utilization in the hermit crab *Clibanarius africanus* (Aurivillus, 1898) (Decapoda: Diogenidae) in the Lagos lagoon: aspects of behavioural and bio- ecology of benthos. Nature and Science 2011;9(3):7-15]. (ISSN: 1545-0740). <http://www.sciencepub.net>.

Key words: Bioecological relationships, shell utilization, hermit crabs, *Clibanarius africanus*

1. Introduction

Bioecological relationship is an important phenomenon in benthic assemblages and therefore plays significant roles in the pattern of structural parameters of benthic communities such as density, biomass, richness, species diversity and spatiotemporal distribution (Tait and Dipper, 1998). Benthic taxocoenosis of the Lagos lagoon system is controlled by the populations of two gastropod mollusc taxa; *Tympanotonus* and *Pachymelania* (Oyekan, 1975; Brown, 2000; Uwadiae, 2009), which provide important biotope for encrusting benthic species and cover for hermit crabs.

Hermit crabs (Anomura: Diogenidae) are crustaceans adapted to use empty gastropod shells as shelter, preventing mechanical damage to their abdomen (Barnes, 2003), and as protection against predation (Leonard *et al.*, 2001). They need increasingly larger shells during their life cycle, a fact that keeps them in constant activity searching for suitable shells (Bertness, 1981b). The life cycle of hermit crabs, therefore depends mostly on the availability of suitable gastropod shells (Hazlett, 1981). Shell utilization appears to be based on a complex and interactive factors, including shell

weight, architecture, volume, height, width, colour and aperture size (Garcia and Mantelatto, 2000, 2001).

Often, availability of empty gastropod shells is a limiting factor to populations of many species of hermit crabs (Scully, 1979) and the sizes of shells occupied by hermit crabs are usually well correlated with crab size owing to mechanisms such as mutual gain shell exchange (Hazlett, 1981, 1989). Hermit crabs seem to utilize among the available empty gastropod shells the most suitable one for their size and shape (Koutsoubas *et al.*, 1993). Shell selection and utilization are not by chance, but based on adequacy and availability of resources (Reese, 1969, Conover, 1978), and is affected by both shell size and species (Conover, 1978; Lively, 1988; Ohmori *et al.*, 1995; Rodrigues *et al.* 2000; Mantelatto *et al.*, 2007). Moreover, the fitness of a particular shell may differ between hermit crab species, reflecting several selection pressures associated with different habitats which act in different ways on each crab species (Bertness, 1981a) in different areas.

As hermit crabs grow they require larger shells. Since suitable intact gastropod shells are sometimes a limited resource, there is often vigorous competition among hermit crabs for shells. The availability of

empty shells at any given place depends on the relative abundance of gastropods and hermit crabs, matched for size. An equally important issue is the population of organisms that prey upon gastropods and leave the shells intact (Tricarico and Gherardi, 2006).

A hermit crab with a shell that is too small cannot grow as fast as those with well-fitted shells, and is more likely to be eaten if it cannot retract completely into the shell (Angel, 2000). Most species of hermit crabs have long, spirally curved abdomens, which are soft, unlike the hard, calcified abdomens seen in related crustaceans. The vulnerable abdomen is protected from predators by shell carried by the hermit crab, into which its whole body can retract (Ingle, 1997). The tip of the hermit crab's abdomen is adapted to clasp strongly onto the columella of the gastropod shell (Chapple, 2002).

This habit of living in a second hand shell gives rise to the popular name "hermit crab", by analogy to a hermit who lives alone. Several hermit crab species use "vacancy chains" to find new shells: when a new, bigger shell becomes available, hermit crabs gather around it and form a kind of queue from largest to smallest. When the largest crab moves into the new shell, the second biggest crab moves into the newly vacated shell, thereby making its previous shell available to the third crab, and so on (Randi *et al.*, 2010).

Although the utilization and selection process of gastropod shells by hermit crabs have been investigated by many authors (Bertness, 1980, 1981a, b, 1982; Blackstone, 1985, 1989; Gherardi and Vannini, 1989; Hazlett, 1989, 1990, 1992; Gherardi, 1991; Ohmori *et al.* 1995; Garcia and Mantelatto, 2001; Meireles and Mantelatto, 2005) around the world, there is scarcity of information on shell utilization in hermit crabs in Nigerian aquatic environment. *Clibanarius africanus* (Aurivillus, 1898) is a species of hermit crab in the family Diogenidae

common on the shores of the Lagos lagoon. Despite its abundance, wide distribution and easy accessibility to its habitat, ecological information on the organism is still scarce. In this study, shell utilization by *C. africanus* in a stretch of intertidal area in the western part of the Lagos lagoon is evaluated.

2. Description of study area

Lagos lagoon is located in the West African Coast of Nigeria. It lies between longitude 3° 54" and 4° 13"E and latitude 6° 25" and 6° 35" N. The Lagoon is more than 50 km long and between 3 to 13 m in width, it is separated from the Atlantic Ocean by a long sand spit of 2 to 5 km wide. The lagoon is characterised by fresh and brackish water conditions occasioned by the heavy input of rainfall run-offs and river discharges during the rainy season, and the influences of tidal incursion and increased surface water evaporation during the dry seasons. The Lagoon is fed by a number of rivers including Ogun and Majidun rivers and Agboyi creek. These water bodies altogether have a drainage area of 103, 637 km² (Oyewo, 1998).

The study sites (Table 1; Figure 1) used for this study were selected along the intertidal area of the Lagos lagoon close to the University of Lagos. This part of the lagoon has been under the stress of human activities because of its accessibility to man. Major stressors include garbage overload arising from indiscriminate dumping of litter during major events such as picnics and students' activities around the lagoon. Modification of the shoreline arising from the construction of brick wall to serve as shoreline protection is evident in the study area. Another prominent feature of the study area is the presence of burrows of crabs scattered all over the shoreline which are most visible at low tides. This area was chosen for this study because of the high biological activities that can be observed in this area. Table 1 presents some physical characteristics of the study sites.

Table 1: Some physical characteristics of the study sites.

Station	Longitude	Latitude	Water colour	Sediment colour
1	003°24'01E	06°31'42"	Brown	Brown
2	003°24'02E	06°31'13"	Brown	Brown
3	003°24'06E	06°31'07"	Brown	Brown
4	003°24'08E	06°31'04"	Brown	Brown
5	003°24'10E	06°31'00"	Gray	Black




Figure 1: Map showing sampling sites (1-5)

3. Materials and methods

3.1 Collection and analysis of samples

Specimens of *C. africanus* in gastropod shells were collected using a van Veen grab at five locations along the lagoon. Three grab hauls were taken from each site and the collected materials washed through a 0.5mm mesh sieve. The residue in the sieve for each station was preserved in 10% formalin solution and kept in labelled plastic containers for further laboratory analysis. Preserved benthic samples were washed with tap water to remove the preservative and any remaining sediment. Specimens of *C. africanus* were sorted out, counted and recorded for all the sampling months and study sites, gastropod species were identified based on the works of Olaniyan (1975), Edmund (1978) and Yoloye (1994).

Specimens of *C. africanus* in gastropod shell and after removal from the shell were weighed with an electronic scale of 0.001g sensitivity. Prior to weighing, the animals were drained on a fine sieve, air dried for 5 minutes on absorbent paper and exposed to air until liquid is no longer visible. The lengths of gastropod shell, width of aperture and length of *C. africanus* were measured in centimeter using a graduated meter rule. Collection of sediment samples and grain size analysis followed the methods described in Holme and McIntyre (1970).

3.2 Statistical analysis

To determine morphometric relationships and correlations between characteristics of hermit crabs and their preferred shells, regression analyses were computed. All statistical analyses were performed using SPSS 10 and Microsoft Excel 2003 for Windows.

4. Results

4.1 Shell type preference of *C. africanus*

The spatial and temporal variations in the frequency of shell type used by *C. africanus* are presented in figure 2. A total of 663 specimens of hermit crab in gastropod shells were collected. Shell occupation of *C. africanus* was limited to gastropod shells belonging to *Pachymelania* and *Tympanotonus* spp. The gastropod shell mostly inhabited were those of *Pachymelania* spp, which accounted for 60.96% of inhabited shells, while about 39% of the shell inhabited belonged to *Tympanotonus* spp. Occurrence of *C. africanus* in the shells of *Pachymelania* spp shell was highest (43) in the month of March while its lowest occurrence (30) in the same shell type was observed in the months of August and April. The number of shells of *Tympanotonus* spp inhabited by *C. africanus* was highest (34) in the month of April and least number (9) of shells occupied in June.

There were slight variations (41- 64) in the total monthly number of individuals of *C. africanus* recorded during the sampling months. The month of April recorded the highest number (64) of *C. africanus* in monthly samples, while the least number of individuals of *C. africanus* was observed in June.

Sixty - one and forty - eight individuals were recorded in the months of July and August respectively, while 59 specimens of *C. africanus* were observed for the months of March and May respectively.




Fig. 2. Spatial and temporal variations in the frequency of shell type selected by *C. africanus*.

4.2 Shell size preference

Specimens of *C. africanus* collected during this study varied between 0.5 and 4.3 cm in length while the length of shell occupied ranged from 1.8 to 5.5 cm. These figures show an overlap indicating that the lengths of *C. africanus* observed are closely related to the length of gastropod shell inhabited. This relationship is clearly depicted in figure 3a where a positive correlation between lengths of shell and that of *C. africanus* is shown in a regression model. Statistically, the relationship between the length of *C. africanus* and the length of shell it occupied was found to be significant ($F = 145.58$, $df = 1$, $P < 0.001$). The length of *C. africanus* was also found to be related to the width of the aperture of the mollusc shell it occupied. Regression analysis indicates that length of *C. africanus* correlated positively (Figure 3b) with width of shell aperture, and this relationship was observed to be statistically significant ($F = 38.58$, $df = 1$, $P < 0.05$).

This study also revealed that another factor of importance in the utilization of shell by *C. africanus* is the weight of the gastropod shell. The study observed a positive correlation (Figure 3c) between the weight of *C. africanus* and the weight of gastropod shell selected. This relationship was also found to be of statistical significance ($F = 27.540$, $df = 1$, $P < 0.005$).


4.3 Sediment characteristics and abundance of *C. africanus*.

In the study area, it was observed that sand and mud intermixed in varying proportions (Figure 4). The percentage of sand ranged from 45.7 to 99.4% while mud varied between 0.5 and 63.9%. In site 1, sand ranged between 45.7 and 87.4%, and mud between 5.7 and 63.9%. Lowest values observed for sand in sites 2 to 5 were 58.8%, 66.2%, 63.7% and 65.2% respectively, while those of mud for the same stations were 4.3%, 0.5%, 3%, and 1.5% respectively.


Variation in the percentage of sand fractions in the sediment of the study area was a major physical parameter that affected the abundance of *C. africanus*

recorded in the study sites. Figure 5 depicts a positive correlation between abundance of the animal and percentage sand in sediment. The overall results indicate that, although there were higher percentage sand fractions than mud in sediment of all the study sites, values for site 3 were relatively higher. Site 3 also recorded the highest number of individuals of *C. africanus*. The lowest mud content was recorded in site 1

3a



3b



3c




Figure 3a-c. Relationships between morphometric characteristic of *C. africanus* and those of the shells inhabited.




Figure 4 . Grain size composition of the study sites.




Figure 5. Relationship between percentage of sand in sediment and abundance of *C. africanus*

5. Discussion

The most striking feature of the result obtained in this study is the positive and significant relationships among morphometric qualities of *C. africanus* and that of the gastropod shell inhabited. This is a confirmation of the views of many Crustacean biologists that shell utilization in hermit crabs is based on the dimension of the crab (Hazlett, 1981, 1989, Koutsoubas *et al.*, 1993, Ates *et al.*, 2007, Mantelatto *et al.*, 2007, Nakin *et al.*, 2007).

Several factors have been adduced and hypotheses put forward to explain the discrepancy in gastropod shell occupancy in hermit crabs (Bertness, 1980; 1981a, b, 1982): 1) Gastropod life cycle; availability of different shell types (species) in nature is determined by the relative abundance of different live gastropods and their mortality rates (Meireles *et al.*, 2003), 2) Environmental conditions; differences in abiotic characteristics of the area in terms of water dynamics (wave activity, intensity of currents, food supply) are determinant of installation of some invertebrate species (Fransozo and Mantelatto, 1998), 3) Predation pressure; several combined actions from natural and artificial predators can act in different ways to reduce the diversity of gastropod shells in the region.

This study reveals that shell dimension constitutes mainly the determinant for *C. africanus* shell utilization. In hermit crabs a well-fitted shell is essential for maintaining low evaporation rates and carrying ample water (Angel, 2000). An appropriately sized shell in good condition allows for effective movement and provides competitive edge than ill – fitted shells. Hermit crabs with broken, ill-fitted shells are restricted to the coast and appear to be in relatively poor conditions (Koutsoubas *et al.*, 1993; Angel, 2000; Ates *et al.*, 2007).

Bertness (1980), reported that hermit crabs do not necessarily live in shells they prefer, the availability of different shell types and contact with competitors for empty shells influences shell occupation. This study revealed that shells of *Tympanotonus* spp provided a low specimen adequacy to *C. africanus*, this may be connected with the relatively higher abundance and availability of *Pachymelania* spp in the study area (Oyenekan, 1975, 1979, 1988; Brown, 2000; Uwadiae, 2009), which provided larger number of shells for *C. africanus*. Although, Yoshino *et al.* (1999), posited that there is a trade-off between shell size and species preference and that less preferred shell species are actively chosen when the more preferred shell species the crabs encounter frequently in the field are of a less suitable size, in this study, considering the great availability (in size and number) of shells of *Pachymelania* spp in the study area, we may infer that the assertion proposed by Yoshino *et al.* (1999) may not be applicable because availability of shell was considered as the overriding factor determining shell usage in this particular study. Shell morphology and morphometric qualities of *Pachymelania* and *Tympanotonus* spp would not have been significant in preferential utilization of shell by *C. africanus* since the two taxa bear similar qualities. Previous studies including Lively (1988) found that shells generally used by hermit crabs were found in the same frequency as the gastropod fauna, revealing a close relationship between shell use and availability of the resources.

The availability of shells in the environment plays a fundamental role in the population dynamics and distributional pattern of hermit crabs (Meireles *et al.* 2003). Although we did not evaluate the shell

availability to the hermit crab community in this present study, the availability of empty shells at any given place depends on the relative abundance of gastropods and hermit crabs matched for size. An equally important issue is the population of organisms that prey upon gastropods and leave the shells intact (Tricarico and Gherardi, 2006). The families Potamiididae and Melaniidae are major components of the benthic communities in the lagoons, estuaries and mangrove swamps in West Africa (Buchanan, 1954). The two families are euryhaline and adapts perfectly to freshwater and brackish water conditions, hence, their survival and continuous presence all year round, providing shells for *C. africanus*. The sedimentary characteristics of the study area have also enhanced the continued presence of these gastropods. Uwadiae *et al* (2009) reported that *P. aurita* preferred sedimentary conditions similar to those observed in this study

Most shells collected have their columela and callosity modified, this agrees with the report of Wolcott (1988), which observed that, shells are modified by hermit crab use, the new shells in some cases are small to accommodate big crabs, so relatively bigger crabs may inhabit shell whose internal volume have been modified by the occupation of another hermit crab. A similar observation has been made by Kinoshita and Okajima (1968) on shells of *Nerita striata* occupied by *Coenobita rugosus* from Japan.

Differences in gastropod shells utilization can occur as a function of the area of occurrence of the hermit crabs (Garcia and Mantelatto, 2000). *Clibanarius africanus* was found in places with small percentage of mud, and high concentrations of sand. The nature of the substratum influences the frequency and ability of the hermit crabs in burying themselves, in a way, that they rarely choose another substrate that is not sand (Oyeneke and Adediran, 1987). This fact corroborates our observations in the present study, the animal was registered principally in locations with the highest percentage of sand.

Besides the interactions with the abiotic factors, the animals share the environment with other organisms from the benthic community. The individuals of a community are in various ways interdependent, and some organisms thrive only in the presence of particular associated fauna. These inter-relations can interfere in the population and distribution of the organism (Pardo *et al.*, 2007). The populations of *C. africanus* like other benthic organisms are threatened by the intensive unregulated human activities in the Lagos lagoon. This suggests that the data gathered may be reflecting both

responses to biotic and abiotic environmental features. Further studies are encouraged to analyse the consistency of the patterns depicted here and explore the causal mechanisms.

References

1. Abrams, PA. Shell selection and utilization in a terrestrial hermit crab, *Coenobita compressus* (H. Milne Edwards). *Oecologia* 1978; 34: 239-253.
2. Angel, JE. Effects of shell fit on the biology of the hermit crab *Pagurus longicarpus* (Say). *J. Exp. Mar. Biol. Ecol.* 2000; 243: 169-184.
3. Ates, AS, Kataúan, T and Kocatas, A. Gastropod Shell Species Occupied by Hermit Crabs (Anomura: Decapoda) along the Turkish Coast of the Aegean Sea. *Turk J. Zool.* 2007; 31:13-18.
4. Ball, EE. Observations on the biology of the hermit crab, *Coenobita compressus* H. Milne Edwards (Decapoda; Anomura) on the west coast of the Americas. *Revista de Biología Tropical* 1972; 20(2): 265-273.
5. Barnes, DKA. Ecology of subtropical hermit crabs in SW Madagascar: short-range migrations. *Marine Biology* 2003; 142: 549-557.
6. Beach, J. *Crustacean Biol.*, 21: 393-406.
7. Bertness, MD. Shell preference and utilization patterns in littoral hermit crabs of the Bay of Panama. *J. Exp. Biol. Ecol.* 1980; 48: 1 - 16.
8. Bertness, MD. Conflicting advantages in resource utilization: The hermit crab housing dilemma. *Am. Nat.* 1981a; 118: 432 - 437.
9. Bertness, MD. The influence of shell-type on hermit crab growth rate and clutch size. *Crustaceana* 1981b; 40: 197 - 205.
10. Bertness, MD. Shell utilization, predation pressure and thermal stress in Panamanian hermit crabs: an interoceanic comparison. *J. Exp. Mar. Biol. Ecol.* 1982; 64: 159 - 187.
11. Biagi, R, Meireles, AL and Mantelatto, FL. Bio-ecological aspects of the hermit crab *Paguristes calliopsis* (Crustacea, Diogenidae) from Anchieta Island, Brazil. *Anais da Academia Brasileira de Ciências.* 2006; 78: 45-62.
12. Blackstone, NW. The effects of shell size and shape on growth and form in the hermit crab *Pagurus longicarpus*. *Biol. Bull.* 1985; 171: 379 - 390.
13. Blackstone, NW. Size, shell-living and carcinization in geographic populations of a hermit crab, *Pagurus hirsutiusculus*. *J. Zool.*, 1989; 217: 477-790.
14. Brown, CA. The diversity and density of macrobenthic fauna in the western part of the

- Lagos lagoon, Lagos, South-west, Nigeria. Ph.D. Thesis, University of Lagos 2000. 346pp
15. Buchanan, J.B. (1954). Marine molluscs of Gold Coast, West Africa. *J. West Africa Sci. Ass.*, **7**: 30 – 45.
 16. Chapple, WD. Mechanoreceptors innervating soft cuticle in the abdomen of the hermit crab, *Pagurus pollicarus*. *Journal of Comparative Physiology* 2002; 188(10): 753–766.
 17. Conover, M.. The importance of various shell characteristics to the shell-selection behavior of the hermit crabs. *Journal of Experimental Marine Biology and Ecology* 1978; 32:131 - 142.
 18. Edmunds, J. **Sea shells and molluscs found on West African Coasts and Estuaries**. Ghana University Press, Accra, 1978. 146pp.
 19. Fransozo, A and Mantelatto, FL. Population structure and reproductive period of the tropical hermit crab *Calcinus tibicen* (Decapoda, Diogenidae) in the Ubatuba Region, São Paulo, Brazil. *Journal of Crustacean Biology* 1998; 18(4): 738-745.
 20. Garcia, RB and Mantelatto, FL. Variability of shell occupation by intertidal and infralitoral *Calcinus tibicen* (Anomura: Diogenidae) populations. *Nauplius* 2000; 8(1): 99-105.
 21. Garcia, RB and Mantelatto, FL. Shell selection by the tropical hermit crab *Calcinus tibicen* (Anomura, Diogenidae) from southern Brazil. *Journal of Experimental Marine Biology and Ecology* 2001; 265: 1 - 14.
 22. Gherardi, F. Relative growth, population structure and shell utilization of the hermit crab *Clibanarius erythropus* in the Mediterranean. *Obelia* 1991; 17: 181 - 196.
 23. Gherardi, F and Vannini, M. Field observations on activity and clustering in two intertidal hermit crabs, *Clibanarius virescens* and *Calcinus laevimanus* (Decapoda, Anomura). *Mar. Behav. Physiol.* 1989; 14: 145 - 159.
 24. Hazlett, BA. The behavioral ecology of hermit crab. *Ann. Rev. Ecol. Syst.* 1981; 12: 1- 22.
 25. Hazlett, BA. Shell exchanges in the hermit crab *Calcinus tibicen*. *Animal Behav* 1989; 37: 104 - 111.
 26. Hazlett, BA. Shell exchange in Hawaiian hermit crabs. *Pacific Sci.* 1990; 44: 401 - 406.
 27. Hazlett, BA. The effect of past experience on the size of shells selected by hermit crabs. *Animal Behav.* 1992; 44: 203 - 205.
 28. Holme, NA and McIntyre, AD. **Methods for the study of marine benthos**. Blackwell Scientific Publications 1970. 334pp.
 29. Imazu, M and Asakura, A. Distribution, reproduction and shell utilization patterns in three species of intertidal hermit crabs on a rocky shore on the Pacific coast of Japan. *Journal of Experimental Marine Biology and Ecology* 1994; 184: 41-65.
 30. Ingle, R. **Hermit crabs of the Northeastern Atlantic Ocean and Mediterranean Sea**. An Illustrated key. Nat. Hist. Mus. Publ., Chapman and Hall. London 1993.
 31. Kellogg, CW. Gastropod shells: a potentially limiting resource for hermit crabs. *Journal of Experimental Marine Biology and Ecology* 1976; 22: 101-111.
 32. Kinosit, AH and Okajima, A. Analysis of shell-searching behavior of the land hermit-crab, *Coenobita rugosus* H. Shell occupation of land hermit crab *C. scaevola* 19 Milne Edwards. *Journal of Faculty of Science of University of Tokyo* 1968; 11: 293-358.
 33. Koutsoubas, D., Labadariou, N. and Koukouras, A. Gastropod shells inhabited by Anomura Decapoda in the North Aegean Sea. *Bios.* 1993; 1: 247-249.
 34. Leonard, M, Gainess, KH and Sandoval, CM. Gastropod shell distribution and factors affecting their utilization by marine hermit crabs in Bahia Kino, Sonora, Mexico, Aquatic Sciences Meeting, Albuquerque 2001.
 35. Lively, C.M. A graphical model for shell-species selection by hermit crabs. *Ecology* 1988; 69: 1233-1238.
 36. Mantelatto, FL and Garcia, RB. Shell utilization pattern of the hermit crab *Calcinus tibicen* (Anomura) (Diogenidae) from Southern Brazil. *Journal of Crustacean Biology* 2000; 20(3):460-467.
 37. Mantelatto, FL, Biagi1, R, Meireles, AL and Marcelo A. Scelzo, MA. Shell preference of the hermit crab *Pagurus exilis* (Anomura: Paguridae) from Brazil and Argentina: a comparative study. *Rev. Biol. Trop.* 2007; 55:153-162.
 38. Meireles, AL and Mantelatto, FL. Shell use by *Pagurus brevidactylus* (Anomura, Paguridae): a comparison between laboratory and field conditions. *Acta Zool. Sinica* 2005; 51: 813-820.
 39. Nakin, MD. V1 and Somers, MJ. Shell availability and use by the hermit crab *Clibanarius virescens* along the eastern Cape Coast, South Africa. *Acta Zoologica Academiae Scientiarum Hungaricae* 2007; 53 (2): 149–155.

40. Nybakken, JW. **Marine Biology. An ecological Approach.** Harper and Row Publisher, New York . 1988. 514pp.
41. Ohmori, H., Wada, S, Goshima, S and Nakao, S. Effects of body size and shell availability on the shell utilization pattern of the hermit crab *Pagurus filholi* (Anomura: Paguridae). *Crust. Res.* 1995; 24: 85-92.
42. Olaniyan, CIO **An introduction to West African Animal Ecology.** Heinemann Education Books Ltd., London 1975. 170pp.
43. Osorno, JJ, Fernandez-Casillas, L and Rodriguez-Juarez, C Are hermit crabs looking for light and large shells? Evidence from natural and field induced shell exchanges. *Journal of Experimental Marine Biology and Ecology* 1998; 222:163-173.
44. Oyekan, JA. A survey of the Lagos lagoon benthos (with particular reference to the mollusca). M.Sc. Dissertation, University of Lagos, Nigeria 1975. 137pp.
45. Oyekan, JA. The ecology of the genus *Pachymelania* in the Lagos lagoon. *Arch. Hydrobiol.* 1979; 86(4): 115—522.
46. Oyekan, JA. Macro-benthic invertebrates communities of Lagos lagoon, Nigeria. *Nigerian Journal of Sciences* 1988; 21: 45 – 51.
47. Oyekan, JA and Adediran, AI. Crab ecology in Lagos area, Nigeria. *Nig. J. Biol. Sci.* 1987; 1(12): 126-133.
48. Oyewo, EO. Industrial sources and distribution of heavy metals in Lagos lagoon and their biological effects on estuarine animals Ph.D. Thesis, University of Lagos, Nigeria 1998. 321pp.
49. Randi, DR, Jeffrey, RC and Sara, ML Social context of shell acquisition in *Coenobita clypeatus* hermit crabs. *Behavioral Ecology* 2010; 21 (3): 639–646.
50. Reese, ES. Behavioral adaptations of intertidal hermit crabs. *Am.Zool.*, 1969; 9: 343-355.
51. Rodrigues, LJ, Dunham, DW and Coates, DA. Shelter preferences in the endemic bermudian hermit crab *Calcinus verrilli* (Rathbun, 1901) (Decapoda, Anomura). *Crustaceana* 2000; 73: 737-750.
52. Scully, EP. The effects of gastropod shell availability and habitat characteristics on shell utilization by the intertidal hermit crab *Pagurus longicarpus* Say. *Jr. Exp. Biol. Ecol.*, 1979; 37: 139-152.
53. Tait, RV and Dipper, FA. **Elements marine ecology.** Butterworth-Heinemann, Oxford 1998. 459pp.
54. Tricarico, E and Gherardi, F Shell acquisition by hermit crabs: which tactic is more efficient? *Behavioral Ecology and Sociobiology* 2006; 60 (4): 492–500.
55. Uwadiae, RE. An ecological study on the macrobenthic invertebrate community of Epe lagoon, Lagos. **PhD.** Thesis University of Lagos, Akoka, Lagos, Nigeria 2009. 253pp.
56. Vance, RR. The role of shell adequacy in behavioral interactions involving hermit crabs. *Ecology* 1972; 53: 1062-1074.
57. Yoloye, VL. Basic invertebrates zoology. Codes and Quanta Nig. Ltd., Lagos 1994. 320pp
58. Yoshino, K, Goshima, S and Nakao, S. The interaction between shell size and shell species preferences of the hermit crab *Pagurus filholi*. *Benthos Res.* 1999; 54: 37-44.

12/15/2010