

Effect of Cadmium Pollution on Neuromorphology and Function of brain in Mice Offspring

*¹Hussein A. Kaoud and ²Mohey M. Mekawy

¹Department of Veterinary Hygiene and Environmental Pollution, Faculty of Veterinary Medicine, Cairo University, Egypt

² Department of Toxicology and Forensic Medicine, Faculty of Veterinary Medicine, Cairo University, Egypt
[*ka-oud@link.net](mailto:ka-oud@link.net)

Abstract: Cadmium chloride (CdCl₂) was given to Swiss-Webster pregnant female mice at the concentrations of (50mg/L) and (100mg/L) (w/v) respectively, in their drinking water. Treatment started few days before pregnancy and it continued until delivery the offspring. After the weaning (22 days of age), all male offspring were isolated and subjected to "Standard Opponent" test at the age of 25 days. The results of this test showed a significant and dose-dependent increase in the non-social behavior, such results showed a significant decline in the social behavior including naso-genital and naso-nasal contact, number of fights, rear, wall rear and displacement activities of the Cd exposed groups. Brain impairments due to the neurotoxic effect of Cd treated groups were indicated by histopathological investigation and neurochemical analysis. **Effect of Cadmium Pollution on Neuromorphology and Function of brain in Mice Offspring.** Nature and Science 2011; 9(4):28-35]. (ISSN: 1545-0740). <http://www.sciencepub.net>.

Key Words: Cadmium; Prenatal exposure; Brain impairment; Mice offspring; Behaviour.

1. Introduction:

Prenatal exposure to neurotoxicants may affect brain development and function. Neurotoxic agents present in the environment and especially in the food chain may reach the brain of the fetus or the newborn during critical periods of brain development. These agents may affect cerebral function and development resulting in long-lasting or permanent deficits in cerebral function that can be reflected at different ages (youngness, adulthood or elderly) as alterations in motor function or coordination and/or altered intellectual function with alterations in learning ability and/or memory. These neurological alterations would be consequence of alterations in the function of one or more of brain [1].

Heavy metals discharged as waste products from different industries are considered a major source of environmental pollution. Cadmium (Cd) which is one of these pollutants has taken considerable attention for its great different toxic effects on living individuals. Cadmium toxicity is increasing in incidence today for several reasons. One of the primary reasons is zinc deficiency in many commonly eaten foods. Zinc, which is protective against cadmium, is becoming increasingly deficient in the soil and consequently in food. Food processing and eating of refined food further reduces zinc intake.

Exposure to cadmium is also increasing due to

its use as in iron coating, steel and copper. It is also used in copper alloys, stabilizers in rubber and plastics, cigarette papers, fungicides and in many other products. Often these industries cause water, air and food pollution with this metal [2]. Cadmium used in industry finds its way into many water supplies. In addition, soft water is more dangerous since the calcium in hard water has a protective effect. Old galvanized pipes and new plastic (PVC) pipes are sources of cadmium in our drinking water. Cadmium is unique among other metals because of its toxicity at a very low dosage and long biologic half life and its low rate of excretion from the body [3]. It is listed by the U.S. Environmental Protection Agency as one of 126 priority pollutants.

Neurotoxicity appears in a variety of neurochemical and behavioral changes [4, 5]. This neurotoxicity was still not regarded to certain specific reason; Cd may exhibit several effects on neural level concerning neurochemical mediators [4, 6] Assembly of cell membrane proteins and phospholipids may also be affected under Cd toxicity [7, 8].

The aim of our present study was to assess the impact of cadmium -exposure present in the environment on some brain functions in mice offspring with special reference to neurobehavioral effects.

2. Materials and Methods

Experimental animals

Adult male and female Swiss-Webster mice were used (n=50, BW 25±5 g), the animals housed in opaque plastic cages (three females with one male in each cage) in the Animal House Unit in Faculty of Veterinary Medicine, Cairo University. Animals were kept under reversed lighting conditions with white light on from 22.30 to 10.30 hours local time. The ambient temperature was regulated between 18 and 22 °C. On pregnancy (appearance of vaginal plug was considered as day one of pregnancy), the males were removed from the cages and the females were subjected to the experiment. Treatment started few days before pregnancy and it continued until delivery. Food (Standard Diet) and water were available ad libitum.

Cadmium administration

Cadmium chloride (analytical grade, Riedel de Haen, Germany) was dissolved in deionized distilled water in the concentrations of (50 mg/L) and (100 mg/L) (w/v). Cadmium concentrations formed the sole drinking fluid source for the experimental group of dams during the prenatal period of the experiment. The drinking fluid containing cadmium concentrations were changed with fresh preparations every five days. The control groups were received deionized distilled water only. All pregnant mice were housed individually.

After weaning at the age of 22 days, 25 male offspring from each treatment category and control group were subjected to the tests at the age of 25 days. Puberty is around day 25-30 in mice, estrogen has been shown to affect non-reproductive behaviors in humans and rodents, including anxiety, fear, and activity levels [9], therefore "Standard opponent" tests were carried out on male offspring.

"Standard Opponent" Test

"Standard opponent" tests under dim red lighting (ca. 8 lux) as described by Brain *et al.*, 1989, [10]. The docile and age-matched male "standard opponents" were rendered anosmic by applying 25µl of 4% zinc sulphate solution to the nasal tract under ether anesthesia for three days prior to encounters [11]. The anosmic 'standard opponent' intruders were introduced in the home cages of 'test animals' and the "standard opponent" test of each 'test animal' was observed visually for 500 seconds. The opponents were used only once and the selected "elements" of behavior were studied [12].

Biochemical Analysis

All the mice were weighed and sacrificed by

cervical decapitation. Brains were cut into two sagittal pieces with surgical scalpel, and tissues of at least five animals were pooled to prepare enough samples for biochemical estimation. Samples were then diluted tenfold and the homogenate was spun at 10,000 rpm for 15 min and the supernatant was used for enzymatic assay. Biochemical estimation by standard methods was conducted for AchE and BchE [13], MDA [14] and ascorbic acid [15] in brain tissues of control and Cd-treated animals.

Cadmium Determination

Each sample was represented by one gram of tissues dissected from the brain of at least five animals were pooled to prepare enough samples for determination of Cd in brain tissues. The samples then placed in a clean screw-capped tube and digested according to the method described by Finerty *et al.*, 1999, [16]. The obtained solutions were then analyzed by using Air/Acetylene Flame Atomic Absorption Spectrophotometer (UNICAM 696 AA Spectrometer) for determination of cadmium levels in examined samples.

Histopathological examination

Histopathological examination in brain of control and Cd-treated animals. Tissue specimens from brain of all experimental mice were collected and fixed in neutral buffered formalin, processed by conventional method, embedded in paraffin, sectioned at 4-5 µm and stained by Haematoxylin and Eosin [17].

Statistical Analysis

Data of "Standard opponent" tests were compared within the experimental groups by ANOVA test and Mann-Whitney U test [18].

3. Results

"Standard Opponent" Test

Results of the various treatments are shown in Tables 1 and 2. The behavioral data (mean ± SE) in the "Standard Opponent" test of male offspring are given in (Tables 1 and 2). Almost all of the elements of behavior were affected by prenatal CdCl₂- exposure and these effects were statistically significant in a dose-dependent manner. The non-social and social behavior of the exposed offspring were significantly and dose-dependently affected showing an increase in the former and a decrease in the later, respectively. The elements of social behavior including attack, numbers of fights, naso-nasal and naso-genital contacts, wall rears and rears were decreased significantly in a dose-dependent manner. The latencies to threat and attack were also increased significantly and dose-

dependently. Overall, the results indicate that the social behavior is significantly decreased due to the prenatal CdCl₂-exposure in a dose- dependent manner. Conversely, the nonsocial behavior and its

elements like wall rears and rears, in young male adult offspring were increased significantly and dose- dependently.

Table 1 . Effect of the prenatal cadmium exposure on the social behavior of male laboratory mice offspring, (Number of seconds allocated to behaviors like)

<u>Parameter</u>	<u>Nonsocial Investigation</u>	<u>Social Investigation</u>	<u>Defense</u>	<u>Threat</u>	<u>Attack</u>	<u>Displacement</u>
<u>Group</u>						
Control	114.7±11.02	228.3±6.44	4.2±0.9	8.4± 1.20	120.4±11.0	24.9±4.20
W/v (50mg/L)	290.31±11*	136.8±12**	5.9±0.5	10.98±1.2	35.88±3.6*	18.7±1.3
W/v (100mg/L)	364.1±**	110.4±5.2**	3.5±1.8	13.4±2.2	4.0± .82**	6.5±0.68*

*and ** statistically significant at $P < 0.05$, and $P < 0.01$ respectively from the control by Mann-Whitney U-test. LD₅₀=110 mg/kg BW oral (cadmium chloride) WHO (1992). The average daily intake of cadmium was estimated to be 4487 µg/kg BW and 8974 for the groups receiving 50 and 100 mg/kg, respectively)

- WHO (1992) and Agency for Toxic Substances and Disease Registry (1999)

Acetylcholinesterase and Butyrylcholinesterase

Activities of AchE and BchE (Acetylcholinesterase and Butyrylcholinesterase) showed a concentration-dependent decrease in all the tissues studied, Tables 3.

Excessive intake of Cd in drinking water significantly reduces AchE and BchE activity in the brain tissues.

MDA

The significant decrease in free radical scavenging enzymes and a concomitant increase in lipid peroxidation (MDA) in brain tissues suggest an increase in oxidative stress after excessive Cd intake. Such increase in free radicals in neuronal cell bodies could be correlated with loss of neurons in brain and synaptic structures in neuromuscular junctions.

The impairments due to the neurotoxic effect of Cd in brain of mice treated with high dose (100 mg /L) were indicated by the histopathological investigation. The impairments included congestion

of blood vessels, necrosis of neurons (Fig.A), neuronphagia (Fig.B) and focal gliosis (Fig.C) as well as haemorrhage in Virchow space. Moreover, in hippocampus the pyramidal cells appeared atrophied and necrosed (Fig. D).

Ascorbic acid

A parallel increase in ascorbic acid content, Table 3, an important antioxidant in brain after Cd intake, is difficult to explain. An increase in ascorbic acid content suggests its role in amelioration of stress.

Concentrations of cadmium in the brain tissues

The concentrations of cadmium in the brain tissues of Cd-exposed groups were 10 µg/g and 65 µg/g of (50mg/L) and (100mg/L) (w/v) respectively, in their drinking water ($P < 0.05$ vs. control). While, the mean concentration of Cd in the brain tissues of control group was 2 µg/g.

Table 2. Effect of the prenatal cadmium exposure on the number of various acts and postures of social behavior of male mouse offspring in a 'standard opponent test'.

[Mean number and latency (mean±SE) of acts and postures]

<u>Parameter</u>	<u>Latency to Threat (sec)</u>	<u>Latency to Attack(sec)</u>	<u>Number of ghts</u>	<u>Number of Naso-Nasal</u>	<u>Number of Naso-Genital</u>	<u>Wall contact</u>	<u>Rears contact</u>
<u>Group</u>							
Control	20	96	20	28	29	23	11
W/v (50mg/L)	90*	140*	8 *	19 *	18*	10 *	7*
W/v % (100mg/L)	260**	280**	2 **	10**	8**	4**	2**

* and ** statistically significant at $P < 0.05$, and $P < 0.01$ respectively from the control by Mann-Whitney U-test.

Table 3. Effects of Cd in drinking water on various parameters in the brain tissues of Mice

<u>Group</u>	<u>Control</u>	<u>50 mg/L (A)</u>	<u>100 mg/L (B)</u>
<u>Parameter</u>			
AchE (Δ OD)	0.10± 0.01	0.059±0.02*	0.024±0.006**
BchE (Δ OD)	0.09± 0.007	0.018±0.013*	0.028±0.026**
MDA (nmoles/mL)	1.95± 0.42	1.31±0.38*	4.87±0.66**
Ascorbic acid (mg/100mL)	0.89± 0.32	1.73±0.19*	2.44±0.26**

$P < 0.01$; Comparison between: Control with A and ** $P < 0.05$; *Data represented as mean ±S.E. control with group B. AchE (Δ OD): Acetylcholinesterase Absorbance/min · mg protein. BchE (Δ OD).




Fig. (A): Microphotograph of brain of mice treated with cadmium showing necrosis of neurons (H & E stain X 200).

Fig. (B): Microphotograph of brain of mice treated with cadmium showing neuronophagia of pyknotic neurons (H & E stain X 200).

Fig. (C): Microphotograph of brain of mice treated with cadmium showing focal gliosis (H & E stain X 200).

Fig. (D): Microphotograph of brain of mice treated with cadmium showing atrophy and necrosis of pyramidal cells of the hippocampus (H & E stain X 200).

Fig. (E): Microphotograph of brain of control, untreated mice showing no histopathological changes (H & E stain X 200)

4. Discussion:

The present results clearly suggest that after the weaning period, various behavioural indices in young male offspring were affected in the "Standard Opponent" test. Thus, these results clearly emphasize that pre and postnatal cadmium exposure is extremely dangerous. A strong correlation exists between maternal and umbilical cord blood cadmium levels indicating prenatal transfer of cadmium from mother to developing fetus in utero [19]. Cadmium can enter into the brain parenchyma and neurons during the critical point of development [20] causing neurological alterations [21, 22]. Baranski, 2007, [23] described the toxic result of cadmium exposure in the form of morphological development, sensory motor reflexes, biochemical and behavioral outcomes in rodents. It is now well documented that significant quantities of compounds that are given to mothers during late pregnancies and during postnatal period, may be transmitted to the offspring in utero and/or via mother's milk during lactation [24-28,19]. It is known that a major portion of brain cells (70%) of the

closely related rats are formed after birth [29]. Furthermore, it has been established that the hippocampus in the brain may be most vulnerable to the neurotoxicity of neurotoxic materials in the very rapid growth period. The hippocampus and the cerebral cortex are the key structures of memory formation [30-33], because the hippocampus is especially indispensable in the integration of spatial information.

There were significant differences in hippocampus and the Opponent" test, in mice which received high dose of Cd (100 mg / L) and untreated group (control). The Opponent" test has been associated with hippocampal activity and cholinergic activity neurotransmission (learning ability and motor activities) Many investigators have reported that hippocampus is one of the most vulnerable regions in the AD brain [34] and hippocampal lesions in general produce changes in rat's activity levels [35] and impairment in spatial memory [36]. So, we could be stated that, hippocampus is the main target of CNS due to Cd exposure. The impairments due to the

neurotoxic effect of Cd in brain of mice treated with high dose (100 mg /L) were indicated by histopathological investigation. Figures A-E, showed congestion of blood vessels, necrosis of neurons, neuronphagia, focal gliosis as well as the pyramidal cells appeared atrophied and necrosed especially in hippocampus (Fig.D).

Histological changes were observed in rat organs exposed to 8 mixtures of metals [37]. Cadmium causes hemorrhages in the autonomic ganglia with cell necroses and damage to nerve cells and nerve fibers. De Castro 1996,[4] observed changes of drinking behavior in rats exposed to acute Cd intoxication. There were significant decreases ($P < 0.05$) in the activity of acetylcholinesterase. (AChE) enzyme in the brain of Cd administered groups, Table 3. Antonio and Leret, 2000, [6] observed enzymatic changes in rats exposed to lead intoxication.

Acetylcholinesterase (AChE) is an enzyme that responsible for hydrolyzing and so deactivating acetylcholine in the body. It is a good indicator of sublethal toxicity by heavy metals [38]. Brain contains 2 forms of AChE, membrane bound forms constitute 90% of the enzyme and soluble form represents the rest 10% [39, 40]. Level of the soluble form considered a simple and sufficient indicator of relative change of AChE in the brain [41, 42] which measures the turnover of Ach activity [43]. Alterations in this enzyme level are indicative to impairment of cholinergic function [44]. Our results revealed significant inhibitory effect on AChE activity in brain tissue of mice offspring which is in accordance with previous investigations of [45- 47] in rats.

Cadmium inhibits release of acetylcholine, probably by interfering with calcium metabolism [37]. Cadmium can enter into the brain parenchyma and neurons [20] causing neurological alterations in human [21] and animal models [22]. This decrease could be due to loss of neuron cell bodies in the brain, loss of synaptic structures, or inhibition of enzyme activity. These effects could be corroborated with cognitive dysfunctions observed in experimental animals.

This study suggested that, exposure to cadmium present in the environment and especially in the food chain may reach the brain of the fetus or the newborn during critical periods of brain development and could be produced cumulative developmental abnormalities in the brain prenatally, which is expressed in the young adult male offspring in disturbed form of long lasting social behavioral outcomes and learning ability. It seems that the

impairments of behaviors in relation to learning and memory are due to the disturbance of the hippocampal circuit and its vast connections through cortical and subcortical pathway.

Acknowledgements

This research was sponsored by Department of Animal Hygiene and Environmental Pollution (Faculty of Veterinary Medicine, Cairo University). We thank Dr. Mervat M.Kamel (Prof.of Animal Behavior Faculty of Veterinary Medicine, Cairo University) for technique assistance.

Corresponding author

Hussein A. Kaoud
Department of Veterinary Hygiene and Environmental Pollution, Faculty of Veterinary Medicine, Cairo University, Egypt
[*ka-oud@link.net](mailto:ka-oud@link.net)

5. References:

- [1] Piedrafita B, Monfort P, Cauli O, Llansola M, Rodrigo R, Montoliu C, El Mlili N, Felipe V Mechanisms of developmental neurotoxicity: Molecular and behavioral correlates. Abstracts / Toxicology Letters 2006; 164S, S1-S324.
- [2] Newsome WH, Davies D, Doucet J PCB and organochlorine pesticides in Canadian human milk. Chemosphere 1995; 30, 2143- 2153.
- [3] Jones MM, Cherian C Cadmium, a unique metal. Toxicology J; 1990: 62,1.
- [4] De Castro E, E Silva, H Ferreira, M Cunha, C Bulcao, C Sarmento, et al., Effect of central acute administration of cadmium on drinking behavior. Pharmacol. Biochem. Behav 1996; 53, 687-693.
- [5] Desi I, Nagymajtenyi L, Schuiz H Behavioural and neurotoxicological changes caused by cadmium treatment of rats during development. J. Appi Toxicol 1998; 18 (1), 63-70.
- [6] Antonio M T, Leret M L Study of the neurochemical alterations produced in discrete brain areas by perinatal low-level lead exposure. Life sciences 2000; 67(6):635-42.
- [7] Herak-Kramberger C M, Sabolic I The integrity of renal cortical brush-border and basolateral membrane vesicles is damaged in vitro by nephrotoxic heavy metals. Toxicology 2001; 156, 139-147.
- [8] M T, Laura Corredor L, Leret M L Study of the activity of several brain enzymes like markers of the neurotoxicity induced by perinatal exposure to lead and/or cadmium. Toxicology Letters 2003; 143, (3), 331-340.
- [9] Morgan MA, Pfaff DW Effects of Estrogen on

- Activity and Fear-Related Behaviors in Mice. *Hormones and Behavior* 2001; (40), 4, 472-482.
- [10] Brain P F, McAllister K H, Walmsley S V Drug effects on social behaviour: methods in ethopharmacology. In: Boulton, A.A., Baker, G.B. & Greenshaw, A.J. (Eds.) *Neuromethods* 1989; Vol. 13 (pp. 687-739). Humana press, Clifton NJ.
- [11] Brain P F, Benton D, Childs G, Parmigiani S The effect of the type of opponent in tests of murine aggression. *Behav Proc* 1981; 6,319 - 328.
- [12] Brain P F, Ajarem J S ,Petkov V V The utility of ethological assessments of murine agonistic interactions in behavioural teratology: the foetal alcohol syndrome 1987. In: Oliver, B., Mos, J. & Brain, P.F. (Eds.) *Ethopharmacology of agonistic behaviour in animals and humans* (pp. 110 -121). Martinus Nijhoff Publishers, Dordrecht.
- [13] Ellman G L , Courney U D , Andres Junior V , Featherstone R M A new and rapid colorimetric determination of AchE activity. *Biochem Pharmacol* 1961; 7, 88-95.
- [14] Netelson, S., 1971. *Techniques of clinical chemistry*, (3rd ed). Springfield, IL: Charles C Thomas.
- [15] Buege J A , Aust S D The thiobarbituric acid assay. *Methods of enzymol* 1978; 52,306-7.
- [16] Finerty M W, Madden J D, Feagly S E , Grodner R M Effect of environs and seasonality on metal residues in tissues of wild and pond raised Cray fish in Southern Louisiana. *Arch Environ. Contam Toxicol* 1990; 19:49.
- [17] Bancroft J D, Stevens A , Turner D R *Theory and Practice of Histological Techniques* 1996; (4th ed). New York, Churchill, Livingstone.
- [18] Sokal R R , Rohlf F J *Biometry: The principles and practice of statistics in biological research* 1981; W.H. Freeman, San Francisco.
- [19] Ajarem J S, Qasim M A, Ahmad M Effect of postnatal lead exposure on the development and behaviour of mice offspring. *Saudi J Biol Sci* 2003; 10, 1-14.
- [20] Nishimura Y, Yamaguchi J. Y , Kanada A, Horimoto K, Kanemaru K, Satoh M Increase in intracellular Cd(2+) concentration of rat cerebellar granule neurons incubated with cadmium chloride, cadmium cytotoxicity under external Ca (2+) – free condition. *Toxicol In Vitro* 2006; 20 (2) : 211- 6.
- [21] Rose C S , Heywood P G , Costanzo R M Olfactory impairment after chronic occupational cadmium exposure *J Occup Med* 1992; 34 (6), 600 -5.
- [22] Lukawski K, Nieradko B, Sieklucka- Dziuba M Effects of cadmium on memory processes in mice exposed to transient cerebral oligemia. *Neurotoxicol Teratol* 2005; 27 (4), 575-84.
- [23] Baranski B Effect of exposure of pregnant rats to cadmium on prenatal and postnatal development of the young. *Neurosci. Res* 2007; 58 (2) 149 – 55.
- [24] Draski L J , Burrig R G , Donovan P J The influence of prenatal and/or postnatal exposure to lead on behavior of preweaning mice. *Physiol Behav* 1989; 45, 711-715.
- [25] Ajarem J S , Ahmad M Behavioral and biochemical consequences of perinatal exposure of mice to instant coffee: a correlative evaluation. *Pharmacol. Biochem Behav* 1991; 40, 847-852.
- [26] Ajarem, J S, Ahmad, M Prenatal nicotine exposure modifies behaviour of mice through early development. *Pharmacol. Biochem. Behav.* 1998 a; 59,313-318.
- [27] Ajarem, J S, Ahmad, M Effects of prenatal exposure of mice to non-alcoholic malt beverage “beer” on the offspring. *Saudi J Bio Sci* 1998 b; 5, 78-92 .
- [28] Ajarem, J S , Brain, P F , Prenatal caffeine exposure modifies behavioral responses in mice. *Behav Pharmacol* 1993; 4,541-544.
- [29] Patel, A J Under nutrition and brain development *Trends Nat Sci* 1983; 6, 151-154.
- [30] Izquierdo, I , da Cunha, C , Rosat, R , Ferreira, M B C , Jerusalinsky, D , Medina, J H Neurotransmitter receptors involved in memory processing by amygdala, medial septum and hippocampus of rats. *Behavioral and Neurobiology* 1992; 58,16-25.
- [31] Thiel, W in: "Computational Thermochemistry" (Eds. K K Irikura and D J Frurip), ACS Symposium Series 677, American Chemical Society, Washington, DC, 1998; pp. 142-161.
- [32] Shirai, N , Suzuki, H Effect of dietary docasahexaenoic acid and catechins on maze behavior in mice. *Ann Nutr Metab* 2004; 48, 51-8
- [33] Leussis, M P , Bolivar, V J Habituation in rodents: a review of behaviour, neurobiology and genetics *Neurosci Biobeh Rev* 2006;30, 1045 – 1064.
- [34] Liu, R, Liu, I Y , Bi, X , Thompson, R F , Doctrow, S R , Malfroy, B , Baudry, M Reversal of age-related learning deficits and brain oxidative stress in mice with superoxide dismutase/catalase mimetics. *Proc Nati. Acad Sci USA* 2003 ; 100, 8526-8531.
- [35] Holland, P C , Lamouneux, J A , Han, J S , Gallagher, M Hippocampal lesions Interfere

- with Pavlovian negative occasion setting. *Hippocampus* 1999 ;9,143 – 157.
- [36] Deacan, R M J , Craucher, A ,Rawlins, JNP Hippocampal cytotoxic lesions effects on species –typical behaviours in mice. *Beh. Brain Res* 2002; 32, 203-213.
- [37] Jadhav, SH, Sarkar, SN, Patil, RD ,Tripathi, HC Effect of subchronic exposure via drinking water to a mixture of eight water –contaminating Metals: A Biochemical and histopathological study in male rats. *Arch Environ. Contam Toxicol* 2007; 53, 667-677.
- [38] Forget, J, Pavillon, JF, Beliaeff, B, Bocquene, G Joint action of pollutant combinations (Pesticides and metals) on survival (LC₅₀ values) and acetyl- cholinesterase activity of *Trigriopus Brevicornis* (Copepoda Harpacticoida). *Environ Toxicol Chem* 1999; 18,912-918.
- [39] Atack, JR, Perry, EK, Bonham, JR, Candy, JM, et al. Molecular forms of acetylcholinestrace and butyrylcholinesterase in the aged human central nervous system, *Journal of Neurochemistry* 1986; 47,263-277.
- [40] Mortensen, S R, Brimijoin, S, Hooper, M J, Padilla, S Comparison of the in vitro sensitivity of rat acetylcholinesterase to chlorpyrifosoxon: What do tissue LC50 values represent? *Toxicol Appl Pharmacol* 1998; 148, 46–49.
- [41] Muller, F, Dumez, Y ,Massoulie, J Molecular forms and solubility of acetylcholinesterase during the embryonic development of rat and human brain *Brain Res* 1985 ; 331, 295-302.
- [42] Zakut, H, Matzkel, A, Schejter, E, Avni, A ,Soreq, H Polymorphism of acetylcholinesterase in discrete regions of the developing human fetal brain. *Journal of Neurochemistry* 1985; 45,382-389.
- [43] Sastry, BVR, Janson, VE, Jaiswal, N ,Tayeb, OS Changes in enzymes of cholinergic system and acetylcholine release in the cerebra of aging male Fischer rats. *Pharmacology* 1983;26,61-72.
- [44] Slechta, DA, Pokora, MJ Lead induced changes in muscarinic cholinergic sensitivity, *Neurotoxicology* 1995; 16, 337-348.
- [45] Eck, PC, Wilson, B Cadmium toxicity. Copyright © 1989 - The Eck Institute of Applied Nutrition and Bioenergetics, Ltd.
- [46] Gupta, A, Gupta A, Murthy, RC, Chandra, SV Neurochemical changes in developing rat brain after pre-and postnatal cadmium exposure. *Bull. Environ Contam Toxicol* 1993; 51, 12-17.
- [47] Antonio, MT, Corredor, L, Leret, ML Study of the activity of several brain enzymes like markers of the neurotoxicity induced by perinatal exposure to lead and/or cadmium *Toxicol Lett* 2003; 143,331-340.