

## Response of Three Sweet Basil Cultivars to Inoculation with *Bacillus subtilis* and Arbuscular Mycorrhizal Fungi under Salt Stress Conditions

Sayed S.A. Abdel-Rahman, \* Atef A.S. Abdel-Kader \*\* and Soha E. Khalil \*\*\*

\*Department of Horticulture, Faculty of Agriculture, Assiut University, Assiut, Egypt

\*\*Department of Medicinal and Aromatic plants Research, Horticulture Research Institute, A.R.C, Giza, Egypt

\*\*\*Department of Water Relation and Field Irrigation, National Research Centre, Dokki, Egypt

Corresponding author: [soha\\_khalil2001@yahoo.com](mailto:soha_khalil2001@yahoo.com)

**Abstract:** A pot experiment was conducted to investigate the effect of inoculation with *Bacillus subtilis* and/or Arbuscular Mycorrhizal Fungus (AMF) or maintained as uninoculated controls on the growth, oil %, oil yield and nutrient uptake of three sweet basil cultivars (Local cultivar, Nano Compatt and Red Bordeaux) under different salt stress levels (0, 1000, 2000 and 4000 ppm). Results indicated that the high salinity level (4000 ppm) caused significantly reduction in plant height, fresh and dry weights of the herb, number of branches/plant, oil % and oil yield as well as contents of N, P and K in leaves of all studied sweet basil cultivars. Meanwhile, sodium Na<sup>+</sup> content in leaves were high, especially at high NaCl concentration. Red Bordeaux cultivar was more sensitive to salinity stress than Local and Nano Compatt cultivars. Inoculation with *Bacillus subtilis* and/or mycorrhizal fungi showed positive effects on growth, oil %, oil yield and nutrient uptake either with or without the salinization treatment. Mycorrhizal colonization showed generally more pronounced effects than *B. subtilis*. Dual inoculation with *B. subtilis* and mycorrhizae provided higher tolerance to salinity compared with the individual treatment. It could be concluded that inoculation of sweet basil cultivars with *B. subtilis* and mycorrhizal fungi may induce increases in tolerance to salinity of the three tested sweet basil cultivars.

[Sayed S.A. Abdel-Rahman, Atef A.S. Abdel-Kader and Soha E. Khalil. **Response of Three Sweet Basil Cultivars to Inoculation with *Bacillus subtilis* and Arbuscular Mycorrhizal Fungi under Salt Stress Conditions.** Nature and Science 2011;9(6):93-111]. (ISSN: 1545-0740). <http://www.sciencepub.net>.

**Key words:** Basil, growth, oil yield, nutrient uptake, *Bacillus subtilis*, mycorrhizae, salt stress

### 1. Introduction

Salinity is the main environmental factor accountable for restricting plant growth and decreasing crop productivity in many areas of the world, especially in arid and semi-arid regions (Apse *et al.*, 1999). About one billion hectares of the world's land area is not in use due to salinity stress (Jain *et al.*, 1989). Additionally, one-third of the world's arable land resources are affected by salinity (Qadir *et al.*, 2000). Soil salinity is mainly attributed to the soluble salts in irrigation water, fertilizers used in agriculture, high evaporation rate and insufficient leaching of ions especially in arid lands (Copeman *et al.*, 1996; and El-Saidi, 1997). Excessive salts in soil lower the availability of water, inhibit metabolic processes and affect nutrient composition, osmotic balance and hydraulic conductivity resulting in stunted growth and productivity of plants (Hopkins, 1999; and Al-Karaki *et al.*, 2001). The development of salt-tolerant crops or desalination of soil by leaching excessive salts, though successful is not economical for sustainable agriculture (Hamdy, 1990; and Cantrell and Linderman, 2001). In this respect, the use of a non-hazardous biological methods, such as Plant Growth-Promoting

Rhizobacteria (PGPR) and mycorrhizal applications to alleviate salt stress and use of moderately salt-tolerant plants are better options (Dixon *et al.*, 1993; and Mayak *et al.*, 2004). Many studies have demonstrated that inoculation with plant growth-promoting rhizobacteria (e.g., *Bacillus subtilis*) and arbuscular mycorrhizal fungi improve plant growth, yield and nutrient uptake under a variety of salinity stress conditions (Al-Karaki *et al.*, 2001 and Mayak *et al.*, 2004).

Plant Growth-Promoting Rhizobacteria (PGPR) have been reported to be key elements for plant establishment under nutrient-imbalance conditions (Egamberdiyeva and Höflich, 2004). PGPR may improve plant growth and yield by direct and indirect mechanisms (Noel *et al.*, 1996). Indirect mechanisms of plant growth stimulation include a variety of mechanisms by which the bacteria prevent phytopathogens from inhibiting plant growth and development (Glick and Bashan, 1997). Direct mechanisms may act on the plant itself and affect growth by providing plants with fixed nitrogen, phytohormones, iron and soluble phosphate (Kloepper and Schroth, 1978). PGPR can also protect plants from

the deleterious effects of some environmental stresses including flooding (Grichko and Glick, 2001), drought (Mayak *et al.*, 2004a), salt (Mayak *et al.*, 2004b) and phytopathogens (Harman and Björkman, 1998). *Bacillus subtilis* can induce plant resistance to stress and produces various plant hormones for growth improvement (Han and Lee, 2005). Many workers have showed that inoculation of plants with *B. subtilis* increased plant growth, yield and nutrient uptake, especially under salt stress conditions (Bochow *et al.*, 2001; Ashraf *et al.*, 2004; Saleh *et al.*, 2005), by influencing phytohormone production (e.g. auxin, cytokinin, or giberallin), and/or by enzymatic lowering of plant ethylene levels (Björkman *et al.*, 1998; Grichko and Glick, 2001).

Arbuscular Mycorrhizal Fungi (AMF) form symbiotic associations with the roots of most plant species (Al-Karaki and Al-Raddad, 1997). These symbiotic associations can enhance plant growth and nutrient uptake under various environmental stress conditions such as salinity, drought and low fertility (Al-Karaki and Al-Raddad, 1997; and Zuccarini and Okurowska, 2008). Also under conditions of low nutrient availability the hyphae of AMF can absorb nutrient from soil beyond the zone depleted by roots so they increase the effectiveness with which the soil volume is explored (Smith and Read, 1997). The beneficial effect of mycorrhizal fungi on plant growth was attributed to enhanced phosphorus uptake (Al-Karaki *et al.*, 2001). Some authors also point out how AM fungi can increase plant resistance to salt stress by influencing the hormonal balance of the host plant (Danneberg *et al.*, 1992) or by increasing water uptake (Ruiz-Lozano and Azcon, 1995).

Little available information in the literature about the interaction between plant growth-promoting rhizobacteria and arbuscular mycorrhizal fungi to alleviating salt stress. Thus, the present study was conducted to evaluate the response of three sweet basil cultivars to inoculation with *Bacillus subtilis* and / or arbuscular mycorrhizal fungi under salt stress.

## 2. Material and Methods

The present experiment was carried out twice at the Experimental Farm of Floriculture, Faculty of Agriculture, Assiut University, Assiut, Egypt during the two successive seasons of 2007 and 2008 to investigate the effect of inoculation with *Bacillus subtilis* and/or Arbuscular Mycorrhizal Fungus (AMF) on the growth, volatile oil %, yield and nutrient uptake of three sweet basil cultivars under salt stress conditions. The three basil cultivars were obtained from the Agriculture Research Center, Giza, Cairo, Egypt. Uniform rooted cuttings (ca. 3 weeks old) of three sweet basil cultivars (Local cultivar, Nano Compact and Red Bordeaux) were transplanted in earthenware pots 30 cm diameter and 40 cm height with perforated bottoms. All pots were filled with 10 Kg of clay loam soil, physical and chemical properties of the soil used were done according to the methods described by Jackson (1973) as shown in Table (1).

At the beginning of May one plantlet was planted in each pot in both seasons. One week after transplanting, plantlets were inoculated with either of *B. subtilis* and/or mycorrhizal fungi or maintained as uninoculated controls. Active strain of *Bacillus subtilis* ( $10^8$  CFU/ml) and arbuscular mycorrhizal fungi (*Glomus irradicans*) provided by the Unit of Biofertilizers, Faculty of Agriculture, Ain Shams University, Shobra El-Kheima, Egypt. The soil was inoculated with *B. subtilis* and/or AMF at three times (one week after transplanting, one month later and one week after the 1<sup>st</sup> cut). *B. subtilis* inoculation was applied at a rate of 10 ml/pot. The inoculation with AMF was placed in the pots at 25 spores/pot before the seeds were sown (Demir and Onogur, 1999). The combined treatment of both microorganisms was applied at 10 ml/pot broth culture of *B. subtilis* + 25 spores/pot of AMF. Also, uninoculated plants are involved as a control.

**Table (1): Some physical and chemical analysis of the soil used in the experiment before planting.**

Properties	Value	Properties	Value	
Texture analysis:	Clay %	Soluble cations meq/L. (soil paste), Ca <sup>++</sup>	2.03	
	Silt %		Mg <sup>++</sup>	0.31
	Sand %		Na <sup>+</sup>	3.06
Texture grade	Clay loam	K <sup>+</sup>	1.83	
Total Ca CO <sub>3</sub> (%)	1.60	Soluble anions meq/L. (soil paste), Cl <sup>-</sup>	2.01	
E.C. dS/m (1:5) soil extract	0.72		CO <sub>3</sub> <sup>-</sup>	-
			HCO <sub>3</sub>	3.79
			SO <sub>4</sub> <sup>-</sup>	1.48
pH (1:2.5 soil suspension)	8.5	Total nitrogen (%)	0.28	
Organic matter (%)	1.27	Total phosphorus (%)	0.164	
		Total potassium (%)	0.221	

Plantlets were irrigated regularly with tap water for two weeks after transplanting, then seedling were subjected to different salinity levels of different NaCl concentrations (0 “tap water”, 1000, 2000 and 4000 ppm). The irrigation whether with tap water or saline water must reach the level of 65% of total Water Holding Capacity (W.H.C.) of the soil by weighing every pot daily and the needed amount of water was added. The general principal stated by Boutraa and Sanders (2001) was used for the water treatment application.

The experiment including 48 treatments which were the combination between three sweet basil cultivars (Local cultivar, Nano Compatt and Red Bordaux), four salinity levels (0 “tap water”, 1000, 2000 and 4000 ppm NaCl) and three inoculation with beneficial microorganisms plus uninoculated control (control, *B. subtilis*, AMF and *B. subtilis* + AMF). The treatments arranged in a split-split-plot design, with three replicates. The three basil cultivars represented the main plots, while the four salinity levels and the four microorganisms inoculation represented in sub-plots and sub-sub plots, respectively.

Plant samples were collected for two cuts, the 1<sup>st</sup> at the beginning of August and the 2<sup>nd</sup> at the end of October to estimate different growth and yield parameters. For each cut three plants were selected randomly from three separated pots and the following growth parameters were recorded: plant height (cm), number of the branches per plant, fresh and dry weights of herbs per plant (g), essential oil percentage in fresh herb and oil yield per plant. The essential oil was extracted by water distillation according to the method described by Guenther (1961).

Samples were collected and dried for 48 h at 70 °C to determine the chemical constituents of leaves which taken at full blooming stage. Total nitrogen was determined by using semi-micro Kjeldahl method described by Black *et al.* (1965). Total phosphorus was determined using Spectrophotometer according to Jackson (1973). Leaf content of K were determined photometrically using a flame photometer according to the method of Jackson (1958). Na content was determined according to the method described by A.O.A.C method (A.O.A.C., 1990).

The collected data were subjected to statistical analysis of variance using the normal (F) test and the means separation were compared by using Least Significant Difference (LSD) at 5% level according to Snedecor and Cochran (1980).

### 3. Results

#### 3.1. Plant height and number of branches:

Obtained results in Table 2 presented a clear comparison between the three chosen cultivars irrespective to salinity and inoculation with microorganisms in both cuts and both seasons. Nano Compatt cultivar surpassed those of Red Bordaux and Local cultivars in plant height and number of branches and with significant difference, followed by Local cultivar.

High salinity badly affected studied growth parameters of basil plant during the two growing seasons and in both cuts (Table 2). It was clear also that low salinity levels of 1000 ppm significantly increased plant height and number of branches of the first cut compared with control plants, while in second cut different salinity levels induced a marked significant reduction in plant height and number of branches compared with control plants in both growing seasons.

Inoculation basil plants with beneficial microorganisms revealed that all treatments caused significant increase in plant height and number of branches compared with control plants. The presented data also showed that the highest significant increase in plant height and number of branches obtained when plants inoculated with *B. subtilis* + AMF treatment compared with control plants. Followed by single inoculation with AMF, while the lowest means obtained for inoculation with *B. subtilis* compared with control plants.

The interaction between tested cultivars and different salinity levels showed that the maximum significant increase in plant height and number of branches in the first cut was observed in Nano compatt cultivar grown under 1000 ppm in both growing seasons, and in control plants (without salinity) of Nano compatt cultivar in the second cut for both growing seasons compared with the other treatments.

Also, the interaction between tested cultivars and different beneficial microorganisms inoculation illustrated that pronounced results were obtained in Nano Compatt cultivar as a response to *B. subtilis* + AMF inoculation in both cuts and both seasons compared with the other treatments.

For the combined effect of different salinity levels and different beneficial microorganism treatments the highest records of the studied growth parameters observed under the combined effect of 1000 ppm and *B. subtilis* + AMF treatment in the first cut in both seasons, and control treatment (without salinity) combined with *B. subtilis* + AMF inoculation in the

second cut of both growing seasons compared with the other treatments.

The tri-interaction between cultivars, salinity and different beneficial microorganisms treatments on plant height and number of branches of the first cut illustrated that the highest records were observed under the combined effect between Nano Compatt

cultivar and 1000 ppm as response to inoculation with *B. subtilis* + AMF compared with the other treatments. While for the second cut, the highest means observed under the combined effect of control treatment (without salinity) of Nano Compatt cultivar and *B. subtilis* + AMF inoculation in both growing seasons.

**Table (2): Effect of inoculation with *Bacillus subtilis* and arbuscular mycorrhizal fungi (AMF) on plant height and number of branches of the three sweet basil cultivars under salt stress condition during 2007 and 2008 seasons.**

Charact.	Treat.	First season				Second season			
		Plant height		No of branches/plant		Plant height		No of branches/plant	
		1 <sup>st</sup> cut	2 <sup>nd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut
<b>Basil cultivars</b>									
Local cultivar		39.0	43.8	6.5	8.0	43.2	49.0	7.8	9.0
Nano Compatt		43.3	50.8	7.4	8.6	48.8	53.9	8.9	10.2
Red Bordeaux		34.3	28.2	5.7	3.7	38.0	31.2	6.6	5.2
LSD <sub>0.05</sub>		1.37	1.31	0.09	0.09	1.74	1.78	0.23	0.19
<b>Salinity levels</b>									
Control		41.8	49.0	7.4	8.4	47.0	53.4	9.2	10.3
1000		44.3	47.0	7.8	7.9	49.0	51.4	9.7	10.0
2000		39.1	40.3	6.1	6.6	42.7	45.5	7.3	7.8
4000		30.3	27.4	4.9	4.2	34.6	28.4	5.0	4.4
LSD <sub>0.05</sub>		1.09	0.91	0.16	0.14	0.60	1.39	0.29	0.11
<b>Beneficial microorganisms</b>									
control		35.5	38.4	5.6	6.0	37.8	39.8	5.6	6.9
<i>B. subtilis</i>		38.6	41.0	6.5	6.7	43.2	44.7	6.6	8.1
AMF		39.8	41.7	6.9	7.0	45.4	46.4	7.1	8.5
<i>B.</i> + AMF		41.6	42.7	7.1	7.3	47.0	47.8	7.3	9.0
LSD <sub>0.05</sub>		0.80	0.81	0.16	0.14	0.85	1.01	0.49	0.10
<b>Basil cultivars X Salinity levels</b>									
Local cultivar	Cont.	40.8	48.5	7.2	9.4	45.6	53.5	8.9	11.1
	1000	43.2	47.1	7.8	9.1	46.7	50.8	9.6	10.5
	2000	40.5	43.1	6.2	7.6	44.2	50.6	7.4	8.5
	4000	31.7	36.7	6.9	6.0	36.4	41.3	5.4	6.1
Nano Compatt	Cont.	47.2	56.3	8.5	10.0	53.9	60.1	10.5	12.3
	1000	48.7	53.8	8.8	9.6	55.1	57.8	11.2	11.9
	2000	43.9	47.7	6.7	8.2	47.9	53.8	8.5	9.5
	4000	33.4	45.3	5.7	6.6	38.4	43.8	5.5	7.0
Red Bordeaux	Cont.	37.5	42.4	6.5	5.7	42.5	46.7	6.2	7.7
	1000	41.0	40.2	6.9	4.9	45.3	45.7	6.5	7.5
	2000	32.9	30.2	5.3	4.0	35.2	32.2	4.5	5.5
	4000	25.6	00.0	4.1	0.0	29.0	00.0	0.0	0.0
LSD <sub>0.05</sub>		1.90	1.58	0.27	0.24	1.04	2.41	N.S	0.20
<b>Basil cultivars X Beneficial microorganisms</b>									
Local cultivar	Cont.	42.2	40.7	5.7	7.1	35.1	43.4	6.8	7.5
	<i>B.sub.</i>	40.1	44.0	6.5	8.0	38.8	49.1	7.8	9.1
	AMF	38.8	44.8	6.9	8.3	40.1	51.1	8.2	9.6
	<i>B.</i> + AMF	35.1	45.9	7.0	8.6	42.2	52.4	8.5	9.9
	AMF								
Nano Compatt	Cont.	39.0	47.7	6.3	7.6	39.0	49.3	7.5	8.8
	<i>B.Sub</i>	44.4	50.8	7.4	8.4	43.1	53.4	9.0	10.0
	AMF	43.1	51.7	7.8	9.1	44.4	55.4	9.5	10.6
	<i>B.</i> + AMF	46.7	52.9	8.2	9.3	46.7	57.3	9.7	11.3
	AMF								
Red Bordeaux	Cont.	34.9	26.8	5.0	3.4	32.4	26.7	5.6	4.3
	<i>B.Sub</i>	34.9	28.1	5.7	3.7	33.9	31.7	6.6	5.2
	AMF	33.9	28.8	6.0	3.7	34.9	32.7	7.1	5.5
	<i>B.</i> + AMF	32.4	29.3	6.2	3.9	35.9	33.7	7.3	5.7
	AMF								
LSD <sub>0.05</sub>		1.39	N.S	0.29	0.24	1.48	N.S	N.S	0.17

Continuous Table (2)

Treat.	Charact.	First season				Second season				
		Plant height		No of branches/plant		Plant height		No of branches/plant		
		1 <sup>st</sup> cut	2 <sup>nd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	
<b>Salinity levels X Beneficial microorganisms</b>										
Cont.	Cont.	40.2	46.6	6.6	7.5	46.6	48.0	7.9	8.7	
	<i>B.sub.</i>	41.8	49.0	7.3	8.4	48.9	53.6	9.2	10.4	
	AMF	42.1	49.8	7.7	8.7	49.8	54.3	9.6	10.9	
	<i>B.+AMF</i>	43.2	50.8	7.9	8.9	49.0	56.8	9.9	11.3	
1000	Cont.	40.9	45.0	6.8	5.5	45.0	46.6	8.2	8.3	
	<i>B.sub.</i>	43.6	46.6	7.6	7.1	46.6	51.1	9.7	10.0	
	AMF	44.9	47.6	8.4	7.7	47.6	53.3	10.2	10.5	
	<i>B.+AMF</i>	47.8	48.9	8.5	8.3	50.8	55.6	10.7	11.0	
2000	Cont.	35.1	37.4	5.2	5.9	37.4	38.8	6.3	6.6	
	<i>B.sub.</i>	39.2	40.7	6.1	6.6	40.7	45.9	7.2	7.7	
	AMF	40.6	41.2	6.3	6.8	41.2	47.6	7.8	8.1	
	<i>B.+AMF</i>	41.4	42.0	6.7	7.2	42.0	49.7	8.1	8.7	
4000	Cont.	25.7	24.6	4.0	3.7	24.6	25.7	4.0	3.7	
	<i>B.sub.</i>	29.8	27.7	5.0	4.1	27.7	28.3	5.1	4.3	
	AMF	31.9	28.2	5.1	4.4	28.2	29.5	5.7	4.6	
	<i>B.+AMF</i>	33.8	29.0	5.4	4.5	29.0	30.1	5.1	4.9	
LSD <sub>0.05</sub>		1.61	N.S	N.S	0.28	1.71	2.02	N.S	0.20	
<b>Basil cultivars X Salinity levels X Beneficial microorganisms</b>										
Local cultivar	Cont.	Cont.	38.4	46.0	6.7	8.2	40.4	47.3	7.8	9.0
		<i>B.sub.</i>	41.1	48.4	7.1	9.5	46.1	53.8	9.1	11.3
		AMF	41.1	49.0	7.3	9.7	47.5	56.5	9.3	11.9
		<i>B.+AMF</i>	42.5	50.4	7.5	10.0	48.3	56.2	9.5	12.0
	1000	Cont.	40.1	44.3	6.9	8.1	42.1	46.3	8.2	8.5
		<i>B.sub.</i>	43.1	46.7	7.5	9.3	46.8	50.7	9.7	10.7
		AMF	43.4	48.4	8.6	9.5	48.2	52.8	9.9	11.0
		<i>B.+AMF</i>	46.0	49.0	8.4	9.7	49.8	53.2	10.5	11.6
	2000	Cont.	35.5	39.3	5.3	6.7	38.2	44.0	6.5	6.9
		<i>B.sub.</i>	40.1	43.7	6.2	7.5	44.7	50.5	7.3	8.4
		AMF	42.4	43.9	6.4	7.9	46.1	52.2	7.8	9.0
		<i>B.+AMF</i>	44.1	45.5	6.2	8.3	47.6	55.5	8.0	9.5
	4000	Cont.	26.3	33.4	4.0	5.3	26.6	36.0	4.5	5.5
		<i>B.sub.</i>	30.9	37.2	5.0	5.8	36.1	41.5	5.3	6.1
		AMF	33.7	37.7	5.2	6.2	40.6	43.2	5.8	6.3
		<i>B.+AMF</i>	36.1	38.5	5.2	6.5	42.2	44.5	6.0	6.5
Nano Compatt	Cont.	Cont.	45.3	53.4	7.3	9.0	48.5	54.7	9.0	10.7
		<i>B.sub.</i>	46.8	56.1	8.4	9.9	53.5	59.3	10.5	12.2
		AMF	48.0	57.5	9.1	10.5	56.3	62.2	11.0	12.7
		<i>B.+AMF</i>	48.7	58.2	9.3	10.7	57.3	64.3	11.5	13.4
	1000	Cont.	44.0	51.8	7.5	8.6	49.5	52.0	9.3	10.2
		<i>B.sub.</i>	47.4	53.7	8.7	9.0	54.7	56.3	11.0	11.8
		AMF	49.0	54.1	9.3	10.3	57.0	60.0	12.0	12.6
		<i>B.+AMF</i>	54.4	55.7	9.5	10.5	59.0	62.7	12.6	13.0
	2000	Cont.	39.2	45.3	5.9	7.2	42.0	49.5	7.2	8.5
		<i>B.sub.</i>	45.1	47.9	6.7	8.2	48.0	54.8	8.5	9.3
		AMF	45.5	48.1	7.0	8.5	50.1	54.3	9.0	9.5
		<i>B.+AMF</i>	45.7	49.4	7.2	8.9	51.3	56.5	9.3	10.6
	4000	Cont.	27.4	40.2	4.3	5.7	30.5	40.9	4.5	5.7
		<i>B.sub.</i>	33.2	45.7	5.7	6.5	38.2	43.3	6.0	6.8
		AMF	35.2	46.9	5.8	7.0	42.1	54.8	6.8	7.5
		<i>B.+AMF</i>	37.8	48.4	6.8	7.0	42.7	54.3	4.7	8.1
Red Bordeaux	Cont.	Cont.	37.1	40.3	5.8	5.2	39.2	42.1	7.0	6.5
		<i>B.sub.</i>	37.4	42.4	6.5	5.8	42.0	47.8	8.0	7.7
		AMF	37.3	43.0	6.7	5.8	43.4	47.2	8.5	8.2
		<i>B.+AMF</i>	38.4	43.9	7.0	6.0	54.3	49.8	9.6	8.5
	1000	Cont.	38.7	39.1	6.0	4.5	40.5	41.5	7.2	6.2
		<i>B.sub.</i>	40.3	37.4	6.7	4.8	45.5	46.2	8.5	7.5
		AMF	42.2	40.4	7.3	5.0	47.0	47.0	8.7	8.0
		<i>B.+AMF</i>	43.1	42.0	7.5	5.3	48.2	48.0	9.0	8.3

	2000	Cont.	30.5	27.7	4.4	3.7	31.5	23.0	5.1	4.5
		<i>B.sub.</i>	32.5	30.5	5.3	4.0	34.5	32.5	5.9	5.5
		AMF	34.0	31.7	5.5	4.0	36.6	36.3	6.5	5.8
		<i>B.+AMF</i>	34.5	31.0	6.0	4.3	38.1	37.0	6.9	6.0
	4000	Cont.	23.5	00.0	3.6	0.0	24.1	00.0	3.0	0.0
		<i>B.sub.</i>	25.4	00.0	4.2	0.0	28.2	00.0	4.0	0.0
		AMF	26.2	00.0	4.3	0.0	30.2	00.0	4.5	0.0
		<i>B.+AMF</i>	27.5	00.0	4.3	0.0	33.5	00.0	4.7	0.0
LSD <sub>0.05</sub>			N.S	N.S	N.S	N.S	N.S	3.49	N.S	0.34

### 3.2. Fresh and Dry weights:

The data present in Table 3 proved that there was a significant difference observed between the three tested basil cultivars in the mean values of their fresh and dry weights during the two growing seasons and in both cuts, where Nano Compatt cultivar recorded higher values compared with that recorded by Red Bordeaux and Local cultivars and with significant differences, followed by Local cultivar.

It was noticed also from the obtained data in the same table that 1000 ppm salinity level showed the highest significant increase in fresh and dry weights of the first cut in both seasons compared with control plants, followed by significant decrease with further increase in salinity levels. While, the data of the second cut revealed that there was an inverse proportional relationship between increasing the severity of salinity on one hand and fresh and dry weights on the other hand in both growing seasons. The results also revealed that the highest salinity level (4000 ppm) revealed the lowest significant means in fresh and dry weights of the two cuts and in both growing seasons.

The use of different types of beneficial microorganisms proved significant increase in fresh and dry weights of basil plant compared with control one, where the highest significant means obtained in *B. subtilis* + AMF inoculation compared with control treatment, followed by single inoculation with AMF.

The interactive effect between the three chosen basil cultivars and different salinity levels deduced that during the two growing seasons and for the three tested cultivars of the first cut, the fresh and dry weights attained their highest values and with significant difference under 1000 ppm compared with the other treatments, moreover Nano Compatt cultivar suppressed that of Red Bordeaux and Local cultivar. While, in the second cut the highest means obtained in control plants (without salinity) of Nano Compatt cultivar.

The combined effect of the three basil cultivars and beneficial microorganisms inoculation revealed that the highest means for both cuts and both seasons were recorded by the interaction between

Nano Compatt cultivar and *B. subtilis* + AMF inoculation compared with the other treatments.

The combined effect of salinity and beneficial microorganisms indicated that the highest means of the first cut obtained under 1000 ppm salinity level combined with *B. subtilis* + AMF inoculation compared with the other treatments. While in the second cut, the highest means obtained when control treatment (without salinity) inoculated with *B. subtilis* + AMF. These results were true in both growing season

The effect of the tri-interaction between tested cultivars, salinity levels and different microorganisms inoculation indicated that the highest values of fresh and dry weights for the first cut in both seasons obtained when Nano Compatt cultivar grown under 1000 ppm and inoculated with *B. subtilis* + AMF. While in the second cut, the highest means obtained under control plants (without salinity) of Nano Compatt cultivar as response to *B. subtilis* + AMF inoculation compared with the other treatments.

### 3.3. Oil percent and yield:

Examination of the collected data in Table 4 revealed the superiority of Local cultivar in oil % and oil yield compared with the other two cultivars and with significant difference, followed by Nano compatt cultivar while the lowest means obtained by Red bordaux cultivar.

Oil % showed progressive significant increase with increasing NaCl salinity level up to 2000 ppm while further increase in salinity up to 4000 ppm revealed significant decrease in both cuts and in both growing seasons compared with control plants. For oil yield, the data of the first cut showed that increasing salinity up to 1000 ppm revealed significant increase in oil yield while further increase in salinity caused significant decrease in oil yield compared with control plants in both growing seasons. Furthermore, the data of the second cut revealed that different salinity level caused significant decrease in oil yield compared with control treatment in both growing seasons.

Pots treated with microorganisms showed greater oil % and oil yield than untreated pots and with significant difference compared with control treatment. The

greatest significant oil % and oil yield means were obtained as a response to *B. subtilis* + AMF inoculation, followed by single inoculation with AMF.

The interaction effect between tested cultivars and different salinity levels showed that during the two growing seasons for both cuts, the highest significant oil % values recorded under the combined effect between Local cultivar and 2000 ppm salinity level

compared with the other treatments. Moreover, the collected data for oil yield indicated in the first cut that the highest significant means obtained under the combined effect between Local cultivar and 1000 ppm salinity level compared with the other treatments in both seasons. while for the second cut, the highest records obtained in control plant of Local cultivar in both growing season.

**Table (3): Effect of inoculation with *Bacillus subtilis* and arbuscular mycorrhizal fungi (AMF) on plant fresh and dry weights of the three sweet basil cultivars under salt stress condition during 2007 and 2008 seasons**

Charact	First season				Second season				
	Fresh weight		Dry weight		Fresh weight		Dry weight		
	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	
<b>Basil cultivars</b>									
Local cultivar	108.1	140.6	21.7	30.2	114.9	170.7	25.6	36.6	
Nano Compatt	120.6	150.3	24.2	31.3	126.8	187.6	27.9	38.7	
Red Bordeaux	66.2 1.48	48.3 8.2	18.8	13.6	106.2	61.6	17.8	13.3	
LSD <sub>0.05</sub>			0.31	1.07	3.0	4.4	0.60	0.60	
<b>Salinity levels</b>									
Control	133.4	152.4	28.9	33.6	151.4	205.2	33.0	42.4	
1000	142.8	148.2	31.6	31.9	156.3	183.0	34.1	39.4	
2000	88.3	104.5	19.1	23.6	87.6	119.5	20.4	24.2	
4000	28.8 1.61	47.2 8.2	06.6	10.8	29.4	52.1	07.5	12.1	
LSD <sub>0.05</sub>			0.68	0.70	1.6	4.4	0.46	0.68	
<b>Beneficial microorganisms</b>									
control	77.9	93.8	17.7	20.9	81.6	115.7	18.8	24.8	
<i>B. subtilis</i>	95.6	112.3	21.0	25.1	105.3	136.3	23.8	28.6	
AMF	106.3	116.8	23.1	26.4	112.7	149.6	25.4	31.4	
<i>B.+ AMF</i>	113.4	129.5 7.4	24.4	27.5	125.1	158.1	27.0	33.3	
LSD <sub>0.05</sub>	1.56		0.47	0.56	1.1	3.9	0.64	0.47	
<b>Basil cultivars X Salinity levels</b>									
Local cultivar	Cont.	142.3	183.4	28.0	37.6	157.4	237.2	33.9	49.1
	1000	151.3	177.9	30.1	37.0	162.9	214.4	35.1	48.4
	2000	106.8	135.9	21.7	30.7	103.5	156.6	24.0	31.4
	4000	32.1	65.3	06.9	15.7	35.9	74.5	09.3	17.3
Nano Compatt	Cont.	153.5	192.9	29.7	39.0	170.7	265.1	36.6	52.5
	1000	166.6	184.0	32.4	37.6	176.4	236.8	37.7	51.0
	2000	117.3	148.2	24.8	31.3	120.3	166.7	27.5	32.5
	4000	45.1	76.3	09.8	16.8	39.9	81.7	09.8	18.9
Red Bordeaux	Cont.	104.5	89.7	28.9	24.2	126.2	113.3	28.6	25.5
	1000	110.4	73.9	32.3	21.2	129.5	97.7	29.6	18.9
	2000	40.7	29.6	10.9	08.9	39.1	35.1	09.6	08.8
	4000	09.2	00.0	03.2	00.0	12.4	00.0	03.3	00.0
LSD <sub>0.05</sub>		2.78	14.2	1.18	1.21	2.8	7.6	0.79	1.18
<b>Basil cultivars X Beneficial microorganisms</b>									
Local cultivar	Cont.	85.7	115.0	17.7	25.6	89.3	135.2	20.0	29.7
	<i>B.sub.</i>	104.1	135.8	20.8	30.3	112.0	166.4	25.6	35.2
	AMF	116.0	151.5	23.2	32.0	120.0	182.6	27.6	39.7
	<i>B.+ AMF</i>	126.8	160.5	24.9	33.1	138.4	198.5	29.1	41.7
Nano Compatt	Cont.	97.3	130.9	20.1	26.9	101.4	163.2	23.0	33.5
	<i>B.sub.</i>	118.2	152.6	23.8	31.1	125.1	182.4	27.6	37.6
	AMF	130.8	145.7	25.9	32.5	133.8	199.8	29.4	40.4
	<i>B.+ AMF</i>	136.2	172.2	27.0	34.2	146.9	204.9	31.5	43.5
Red Bordeaux	Cont.	50.7	35.4	15.2	20.9	54.1	48.7	13.3	11.2
	<i>B.sub.</i>	64.5	48.6	18.6	25.1	78.8	60.2	18.2	13.0
	AMF	72.2	53.1	20.3	26.4	84.3	66.4	19.3	14.2
	<i>B.+ AMF</i>	77.4	56.2	21.3	27.5	90.0	71.0	20.3	14.9
LSD <sub>0.05</sub>		2.70	12.8	N.S	0.97	1.8	6.8	N.S	0.82

Continuous Table (3)

Treat.		Charact.	First season				Second season			
			Fresh weight		Dry weight		Fresh weight		Dry weight	
			1 <sup>st</sup> cut	2 <sup>nd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut
<b>Salinity levels X Beneficial microorganisms</b>										
Cont.	Cont.		107.6	134.7	24.2	29.4	123.5	175.7	26.8	37.7
	<i>B.sub.</i>		127.3	153.3	27.7	33.6	148.3	197.3	32.6	41.3
	AMF		146.1	143.8	31.1	34.5	159.2	217.7	35.1	44.2
	<i>B.+ AMF</i>		152.7	177.6	32.6	36.8	174.7	230.0	37.6	46.3
1000	Cont.		120.4	124.9	27.4	27.6	124.8	159.7	27.3	33.2
	<i>B.sub.</i>		139.0	147.0	30.7	31.9	154.8	185.3	34.3	38.4
	AMF		151.3	157.4	33.5	33.7	165.0	194.7	36.5	42.1
	<i>B.+ AMF</i>		160.4	163.6	34.7	34.4	180.3	192.4	38.5	44.0
2000	Cont.		62.8	84.7	14.3	19.2	54.8	94.0	15.4	20.0
	<i>B.sub.</i>		87.9	104.7	19.2	24.5	89.3	115.4	20.9	23.7
	AMF		96.1	111.5	20.7	25.3	94.7	125.4	21.9	25.5
	<i>B.+ AMF</i>		106.2	117.3	22.3	25.5	111.7	143.0	23.3	27.7
4000	Cont.		20.8	30.8	04.8	07.3	23.2	33.4	05.7	08.2
	<i>B.sub.</i>		28.1	44.3	06.5	10.5	28.7	47.3	07.4	11.0
	AMF		31.8	54.3	07.3	12.2	32.0	60.6	08.2	13.9
	<i>B.+ AMF</i>		34.4	59.5	07.9	13.3	33.7	67.1	08.6	15.3
LSD <sub>0.05</sub>			3.11	N.S	0.93	N.S	2.1	7.8	1.28	0.95
<b>Basil cultivars X Salinity levels X Beneficial microorganisms</b>										
Local cultivar	Cont.	Cont.	118.4	158.5	24.2	34.0	132.0	200.7	26.7	43.5
		<i>B.sub.</i>	134.6	155.6	26.1	37.4	148.0	227.5	32.8	47.8
		AMF	150.0	173.0	29.5	38.2	164.0	250.0	36.8	51.0
		<i>B.+AMF</i>	166.0	184.6	32.3	40.7	185.5	270.5	39.4	54.1
	1000	Cont.	127.6	155.6	25.8	34.0	136.5	183.8	38.1	41.1
		<i>B.sub.</i>	145.1	173.0	29.0	36.8	156.0	216.3	34.9	46.4
		AMF	158.0	184.6	31.6	38.0	170.0	223.8	37.6	52.3
		<i>B.+AMF</i>	174.6	198.3	33.9	39.2	189.0	233.8	39.8	53.9
	2000	Cont.	75.1	108.2	16.1	23.9	59.0	118.3	18.1	25.4
		<i>B.sub.</i>	106.4	135.2	21.5	31.7	108.0	155.0	25.2	31.0
		AMF	120.0	147.2	24.1	33.5	109.0	165.0	26.1	33.7
		<i>B.+AMF</i>	125.8	153.0	25.0	33.6	138.0	188.0	26.5	35.4
	4000	Cont.	21.6	37.8	4.8	10.5	36.0	38.0	7.1	8.9
		<i>B.sub.</i>	30.1	61.0	6.5	15.1	36.0	66.7	9.3	15.5
		AMF	35.9	79.3	7.7	18.3	37.0	91.7	10.1	21.7
		<i>B.+AMF</i>	40.6	83.2	8.6	19.0	41.0	101.5	10.8	23.2
Nano Compatt	Cont.	Cont.	126.8	176.1	25.6	34.7	145.0	239.0	31.8	47.1
		<i>B.sub.</i>	143.9	198.6	27.4	38.5	165.0	252.5	35.4	50.9
		AMF	169.8	141.6	32.3	39.9	175.2	279.8	37.6	54.7
		<i>B.+AMF</i>	173.4	219.8	33.6	42.7	197.5	289.0	41.4	57.4
	1000	Cont.	144.5	167.3	29.2	33.7	152.5	214.2	31.7	43.0
		<i>B.sub.</i>	161.5	192.0	31.1	37.2	168.0	243.0	36.9	50.7
		AMF	177.3	204.0	34.0	39.4	180.0	255.2	39.4	53.4
		<i>B.+AMF</i>	183.2	208.3	35.2	40.1	205.0	235.0	42.8	56.8
	2000	Cont.	84.6	125.6	18.5	27.5	77.0	137.5	20.9	28.2
		<i>B.sub.</i>	121.8	148.0	26.4	32.3	129.0	159.2	28.5	31.2
		AMF	127.2	153.7	26.9	32.5	135.0	174.2	29.7	33.5
		<i>B.+AMF</i>	135.5	165.3	27.3	33.0	140.0	196.0	30.9	37.0
	4000	Cont.	33.2	54.6	7.0	11.5	31.0	62.0	7.8	15.7
		<i>B.sub.</i>	45.5	71.8	10.1	16.3	38.5	75.0	9.6	17.4
		AMF	49.0	83.6	10.5	18.2	45.0	90.0	10.8	20.0
		<i>B.+AMF</i>	52.6	95.2	11.7	21.0	45.0	99.8	11.0	22.0
Red Bordaax	Cont.	Cont.	77.5	69.5	22.7	19.5	93.5	87.5	21.9	22.5
		<i>B.sub.</i>	103.5	87.2	29.6	24.9	132.0	112.0	29.6	25.2
		AMF	118.5	95.1	31.5	25.4	138.3	123.3	30.0	26.8
		<i>B.+AMF</i>	118.6	106.9	32.0	27.0	141.0	130.5	30.9	27.4
	1000	Cont.	89.1	51.8	27.2	15.2	85.5	81.0	22.1	15.6
		<i>B.sub.</i>	110.3	76.0	32.0	21.7	140.5	96.5	30.0	18.0
		AMF	118.7	83.7	34.8	23.8	145.0	105.0	32.5	20.5
		<i>B.+AMF</i>	123.4	84.1	35.1	24.0	147.0	108.4	32.8	21.4
	2000	Cont.	28.6	20.2	8.4	6.2	28.5	26.3	7.1	6.5
		<i>B.sub.</i>	35.6	31.0	9.7	9.4	31.0	32.1	8.9	8.8

		AMF	41.1	33.5	11.0	9.8	40.0	37.0	10.0	9.4
		B.+AMF	57.4	33.6	14.5	10.0	57.0	45.0	12.4	10.6
	4000	Cont.	7.5	00.0	2.5	00.0	09.0	00.0	2.2	00.0
		B.sub.	8.7	00.0	3.0	00.0	11.5	00.0	3.3	00.0
		AMF	10.5	00.0	3.8	00.0	14.0	00.0	3.7	00.0
		B.+AMF	10.0	00.0	3.5	00.0	15.0	00.0	3.9	00.0
LSD <sub>0.05</sub>			5.39	N.S	1.62	N.S	3.7	13.5	2.22	1.64

**Table (4): Effect of inoculation with *Bacillus subtilis* and arbuscular mycorrhizal fungi (AMF) on oil percentage and oil yield in fresh herb of the three sweet basil cultivars under salt stress condition during 2007 and 2008 seasons.**

Charact.	Treat.	First season				Second season				
		Oil percentage		Oil yield		Oil percentage		Oil yield		
		1 <sup>st</sup> cut	2 <sup>nd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	
<b>Basil cultivars</b>										
	Local cultivar	0.19	0.22	0.214	0.317	0.21	0.30	0.242	0.517	
	Nano Compatt	0.13	0.14	0.170	0.225	0.12	0.14	0.156	0.269	
	Red Bordeaux	0.12	0.11	0.079	0.070	0.12	0.12	0.087	0.099	
LSD <sub>0.05</sub>		0.009	0.015	0.005	0.004	0.005	0.010	0.002	0.005	
<b>Salinity levels</b>										
	Cont.	0.14	0.17	0.191	0.274	0.13	0.19	0.206	0.405	
	1000	0.15	0.18	0.220	0.270	0.15	0.20	0.234	0.392	
	2000	0.16	0.18	0.153	0.201	0.17	0.21	0.158	0.275	
	4000	0.12	0.10	0.036	0.071	0.16	0.14	0.047	0.108	
LSD <sub>0.05</sub>		0.003	0.004	0.002	0.001	0.002	0.004	0.002	0.002	
<b>Beneficial microorganisms</b>										
	control	0.12	0.13	0.100	0.138	0.13	0.16	0.106	0.213	
	<i>B. subtilis</i>	0.14	0.15	0.141	0.195	0.15	0.18	0.153	0.274	
	AMF	0.15	0.17	0.169	0.230	0.15	0.19	0.171	0.319	
	<i>B.+ AMF</i>	0.16	0.17	0.190	0.253	0.17	0.21	0.216	0.374	
LSD <sub>0.05</sub>		0.003	0.004	0.002	0.002	0.004	0.004	0.001	0.002	
<b>Basil cultivars X Salinity levels</b>										
	Local cultivar	Cont.	0.19	0.22	0.267	0.409	0.19	0.29	0.300	0.692
		1000	0.20	0.23	0.307	0.405	0.20	0.30	0.335	0.651
		2000	0.22	0.25	0.235	0.333	0.23	0.32	0.250	0.507
		4000	0.15	0.18	0.047	0.120	0.23	0.28	0.082	0.218
	Nano Compatt	Cont.	0.12	0.15	0.185	0.291	0.11	0.13	0.193	0.364
		1000	0.13	0.15	0.218	0.286	0.12	0.15	0.213	0.347
		2000	0.15	0.16	0.172	0.231	0.14	0.15	0.170	0.258
		4000	0.11	0.12	0.051	0.092	0.12	0.13	0.048	0.106
	Red Bordeaux	Cont.	0.11	0.14	0.119	0.125	0.10	0.15	0.124	0.167
		1000	0.12	0.15	0.136	0.114	0.12	0.16	0.155	0.161
		2000	0.13	0.13	0.052	0.039	0.14	0.16	0.056	0.059
		4000	0.10	0.00	0.009	0.000	0.12	0.00	0.013	0.000
LSD <sub>0.05</sub>			0.005	0.007	0.004	0.003	0.004	0.007	0.004	0.003
<b>Basil cultivars X Beneficial microorganisms</b>										
	Local cultivar	Cont.	0.16	0.17	0.136	0.201	0.18	0.27	0.155	0.372
		<i>B.sub.</i>	0.18	0.22	0.197	0.307	0.20	0.30	0.220	0.490
		AMF	0.20	0.23	0.241	0.361	0.21	0.31	0.246	0.566
		<i>B.+ AMF</i>	0.21	0.24	0.282	0.399	0.25	0.32	0.345	0.640
	Nano Compatt	Cont.	0.11	0.13	0.110	0.170	0.11	0.12	0.111	0.303
		<i>B.sub.</i>	0.13	0.14	0.151	0.214	0.12	0.13	0.153	0.243
		AMF	0.13	0.15	0.176	0.247	0.13	0.14	0.167	0.281
		<i>B.+ AMF</i>	0.14	0.15	0.190	0.270	0.13	0.16	0.193	0.348
	Red Bordeaux	Cont.	0.10	0.10	0.053	0.043	0.11	0.10	0.052	0.065
		<i>B.sub.</i>	0.11	0.10	0.074	0.065	0.12	0.11	0.086	0.089
		AMF	0.12	0.12	0.092	0.081	0.13	0.13	0.100	0.110
		<i>B.+ AMF</i>	0.13	0.12	0.097	0.090	0.13	0.14	0.110	0.133
LSD <sub>0.05</sub>			0.005	0.007	0.003	0.003	0.007	0.007	0.003	0.003

**Continuous Table (4):**

Charact.		First season				Second season				
		Oil percentage		Oil yield		Oil percentage		Oil yield		
		1 <sup>st</sup> cut	2 <sup>nd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	
<b>Salinity levels X Beneficial microorganisms</b>										
Cont.	Cont.	0.12	0.14	0.132	0.188	0.12	0.17	0.147	0.314	
	<i>B.sub.</i>	0.14	0.16	0.176	0.258	0.13	0.18	0.193	0.366	
	AMF	0.15	0.18	0.215	0.305	0.13	0.20	0.215	0.437	
	<i>B.+ AMF</i>	0.15	0.19	0.239	0.343	0.15	0.21	0.268	0.503	
1000	Cont.	0.13	0.15	0.154	0.188	0.13	0.18	0.162	0.302	
	<i>B.sub.</i>	0.15	0.17	0.210	0.258	0.14	0.20	0.218	0.378	
	AMF	0.16	0.19	0.245	0.301	0.15	0.21	0.247	0.419	
	<i>B.+ AMF</i>	0.17	0.19	0.273	0.333	0.17	0.23	0.310	0.470	
2000	Cont.	0.14	0.16	0.092	0.138	0.15	0.18	0.085	0.185	
	<i>B.sub.</i>	0.15	0.18	0.142	0.200	0.17	0.21	0.154	0.257	
	AMF	0.17	0.19	0.175	0.225	0.17	0.22	0.171	0.290	
	<i>B.+ AMF</i>	0.18	0.20	0.202	0.242	0.19	0.24	0.223	0.367	
4000	Cont.	0.11	0.08	0.023	0.038	0.14	0.11	0.030	0.052	
	<i>B.sub.</i>	0.12	0.10	0.034	0.066	0.15	0.14	0.046	0.095	
	AMF	0.12	0.11	0.043	0.087	0.16	0.14	0.051	0.131	
	<i>B.+ AMF</i>	0.13	0.11	0.045	0.094	0.17	0.15	0.062	0.155	
LSD <sub>0.05</sub>		0.006	0.008	0.004	0.003	N.S	0.008	0.003	0.003	
<b>Basil cultivars X Salinity levels X Beneficial microorganisms</b>										
Local cultivar	Cont.	Cont.	0.15	0.16	0.178	0.254	0.16	0.27	0.211	0.542
		<i>B.sub.</i>	0.18	0.22	0.242	0.383	0.18	0.28	0.266	0.637
		AMF	0.20	0.24	0.300	0.468	0.18	0.30	0.295	0.750
		<i>B.+AMF</i>	0.21	0.25	0.249	0.516	0.23	0.31	0.427	0.843
	1000	Cont.	0.16	0.18	0.204	0.280	0.17	0.28	0.230	0.515
		<i>B.sub.</i>	0.20	0.23	0.290	0.398	0.19	0.30	0.296	0.649
		AMF	0.21	0.24	0.332	0.443	0.20	0.31	0.340	0.694
		<i>B.+AMF</i>	0.23	0.26	0.402	0.516	0.25	0.32	0.473	0.748
	2000	Cont.	0.18	0.22	0.135	0.215	0.20	0.29	0.118	0.839
		<i>B.sub.</i>	0.20	0.25	0.213	0.338	0.22	0.31	0.238	0.481
		AMF	0.23	0.26	0.276	0.383	0.24	0.33	0.269	0.545
		<i>B.+AMF</i>	0.25	0.26	0.315	0.398	0.27	0.35	0.273	0.658
	4000	Cont.	0.13	0.14	0.028	0.053	0.20	0.23	0.059	0.087
		<i>B.sub.</i>	0.14	0.18	0.042	0.110	0.22	0.29	0.079	0.193
		AMF	0.16	0.19	0.057	0.151	0.22	0.30	0.081	0.275
		<i>B.+AMF</i>	0.15	0.20	0.061	0.166	0.26	0.31	0.107	0.315
Nano Compatt	Cont.	Cont.	0.11	0.13	0.139	0.229	0.10	0.12	0.145	0.287
		<i>B.sub.</i>	0.12	0.14	0.173	0.278	0.11	0.12	0.182	0.303
		AMF	0.12	0.15	0.204	0.305	0.12	0.13	0.210	0.364
		<i>B.+AMF</i>	0.13	0.16	0.225	0.352	0.12	0.15	0.237	0.434
	1000	Cont.	0.11	0.13	0.159	0.217	0.11	0.13	0.168	0.278
		<i>B.sub.</i>	0.13	0.14	0.209	0.269	0.12	0.14	0.202	0.340
		AMF	0.14	0.16	0.248	0.326	0.12	0.15	0.216	0.383
		<i>B.+AMF</i>	0.14	0.16	0.256	0.333	0.13	0.17	0.267	0.456
	2000	Cont.	0.13	0.14	0.110	0.176	0.13	0.13	0.100	0.179
		<i>B.sub.</i>	0.14	0.15	0.171	0.222	0.14	0.15	0.181	0.239
		AMF	0.15	0.16	0.191	0.246	0.14	0.15	0.189	0.261
		<i>B.+AMF</i>	0.16	0.17	0.217	0.281	0.15	0.18	0.210	0.353
	4000	Cont.	0.10	0.10	0.033	0.060	0.10	0.11	0.031	0.068
		<i>B.sub.</i>	0.11	0.12	0.050	0.086	0.12	0.12	0.046	0.090
		AMF	0.12	0.13	0.059	0.109	0.12	0.13	0.054	0.117
		<i>B.+AMF</i>	0.12	0.12	0.063	0.114	0.13	0.15	0.059	0.150
Red Bordeaux	Cont.	Cont.	0.10	0.12	0.078	0.082	0.09	0.13	0.084	0.114
		<i>B.sub.</i>	0.11	0.13	0.114	0.113	0.10	0.14	0.132	0.157
		AMF	0.12	0.15	0.142	0.143	0.10	0.16	0.139	0.197
		<i>B.+AMF</i>	0.12	0.15	0.142	0.160	0.10	0.18	0.141	0.235
	1000	Cont.	0.11	0.13	0.098	0.067	0.10	0.14	0.086	0.113
		<i>B.sub.</i>	0.12	0.14	0.132	0.106	0.11	0.15	0.155	0.145
		AMF	0.13	0.16	0.154	0.134	0.13	0.17	0.186	0.179
		<i>B.+AMF</i>	0.13	0.18	0.160	0.151	0.13	0.19	0.191	0.206

	2000	Cont.	0.11	0.12	0.031	0.024	0.13	0.12	0.037	0.032
		<i>B.sub.</i>	0.12	0.13	0.043	0.040	0.14	0.16	0.043	0.051
		<i>B.+AMF</i>	0.14	0.14	0.058	0.047	0.14	0.17	.056	0.063
			0.13	0.14	0.075	0.047	0.15	0.20	0.086	0.090
	4000	Cont.	0.09	0.00	0.007	0.00	0.11	0.00	0.001	0.00
		<i>B.sub.</i>	0.10	0.00	0.009	0.00	0.12	0.00	0.014	0.00
		AMF	0.11	0.00	0.012	0.00	0.13	0.00	0.018	0.00
		<i>B.+AMF</i>	0.10	0.00	0.010	0.00	0.13	0.00	0.020	0.00
LSD <sub>0.05</sub>			0.011	0.014	0.007	0.006	N.S	0.013	0.005	0.006

Regarding the interaction between the three chosen basil cultivars and different microorganisms inoculation in both cuts and in the two growing seasons the data showed that the highest significant means in both oil % and oil yield obtained in Local cultivar as response to inoculation with *B. subtilis* + AMF compared with the other treatments, followed by single inoculation with AMF in the same cultivar.

Oil % revealed mostly significant increase under the combined effect of *B. subtilis* + AMF treatment and 2000 ppm in the both cuts and in both seasons compared with the other treatments. For the oil yield the data showed that *B. subtilis* + AMF inoculation revealed the highest significant means in both seasons and both cuts combined with 1000 ppm in the first cut and with control plants (without salinity) in the second cut compared with the other treatments.

The tri-interaction between the three studied factors showed that inoculated Local cultivar with *B. subtilis* + AMF under 2000 ppm salinity level revealed mostly the highest significant means of oil % in both cuts and seasons. Furthermore, inoculated Local cultivar with *B. subtilis* + AMF under 1000 ppm showed the highest significant records for oil yield in the first cut of both seasons, and *B. subtilis* + AMF combined with control plants (without salinity) of Local cultivar in the second cut of both growing

### 3.4. Minerals content:

The obtained results pointed out in both growing seasons that there was a significant difference observed between the three tested basil cultivars in the mean values of their leaf mineral content where the highest values for N, P and K % were observed in Nano Compatt cultivar in the two cuts, followed by Local cultivar, while the lowest means obtained in Red Bordeaux cultivar (Tables 5&6). For Na % the data revealed that the highest Na % obtained in Red Bordeaux cultivar, while the lowest values obtained in Nano Compatt cultivar in both cuts for both growing seasons (Table 6).

Leaf N, P and K % revealed an increase in their contents in the first cut of both growing seasons with increase salinity concentrations up to 1000 ppm followed by significant decrease with further increase

in salinity levels compared with control plants. The data of the second cut showed that increasing salinity badly affected minerals content compared with control plants during the two growing seasons (Tables 5 & 6). Na<sup>+</sup> content showed in the first cut of both growing seasons significant increase with increasing salinity level compared with control plants, similar trend was obtained in the second cut but with decrease in Na% under the highest salinity level (4000 ppm), this result were true for both seasons (Table 6). All biological inoculations increased N, P and K % significantly compared to control treatment in absence of salinity and cultivars in both cuts and both seasons. Minerals content showed that the highest significant increases in their content obtained as a response to *B. subtilis* + AMF treatment compared with control plants, followed by single inoculation with AMF in both P and K % and followed by *B. subtilis* in case of N %, similar results obtained in the second cut and second season (Tables 5 & 6). While, reversed trend obtained in Na % where different inoculations with microorganisms revealed gradual significant decrease in Na % compared with control plants in both cuts of both seasons, the lowest significant means obtained in *B. subtilis* + AMF treatment followed by AMF treatment (Table 6).

Nano Compatt cultivar grown under 1000 ppm showed the highest values in N, P and K % compared with the other treatments of both cuts and both seasons. Salt stress increased the concentration of Na<sup>+</sup> in leaves of the three basil cultivars where the highest values for Na% obtained under the highest salinity levels 2000 & 4000 ppm in the three basil cultivars, also it was clear from the results that Red Bordeaux cultivar was more sensitive to salinity than the other two basil cultivars.

Nano Compatt cultivar inoculated with *B. subtilis* + AMF treatment revealed generally the highest values of N, P and K % in both cuts and during the two growing seasons. Different inoculations revealed decrease in Na % in both seasons and both cuts, where the lowest means obtained as a response to *B. subtilis* + AMF treatment for the three basil cultivars.

Different microorganisms inoculation showed marked increase in N, P and K concentrations under different salinity levels compared with control plants, where the highest concentrations of N, P and K in the

first cut found under 1000 ppm combined with *B. subtilis* + AMF inoculation in both growing seasons. While, in the second cut the highest concentrations obtained in control treatment (without salinity) as a response to *B. subtilis* + AMF inoculation. For Na % the present data revealed that different microorganisms inoculation caused decrease in Na concentration under different salinity levels compared with control plants, where the highest concentrations Na% of the first cut found in uninoculated treatment under 4000 ppm and in uninoculated treatment under 2000 ppm in the second cut, these results were true for both growing seasons.

The interactions between the three studied factors indicated in the first cut of the two growing seasons that the highest records of N, P and K % obtained in Nano Compatt cultivar grown under 1000 ppm as a response to *B. subtilis* + AMF inoculation. While, the data of the second cut showed that the highest records obtained in control plants (without salinity) of Nano Compatt cultivar as a response to *B. subtilis* + AMF inoculation in both seasons. Furthermore, the means of Na % showed that Red Bordeaux cultivar was highly sensitive for the high concentrations of Na content (4000 ppm) than the other two cultivars even with different inoculations with microorganisms.

**Table (5): Effect of inoculation with *Bacillus subtilis* and arbuscular mycorrhizal fungi (AMF) on Nitrogen and Phosphorus % in dry leaves of the three sweet basil cultivars under salt stress condition during 2007 and 2008 seasons.**

Charact.		First season				Second season			
		N %		P %		N %		P %	
		1 <sup>st</sup> cut	2 <sup>nd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut
<b>Basil cultivars</b>									
Local cultivar		1.93	1.96	0.206	0.200	2.09	2.17	0.211	0.219
Nano Compatt		1.97	1.99	0.211	0.213	2.11	2.20	0.217	0.226
Red Bordeaux		1.85	1.45	0.194	0.147	2.07	1.65	0.207	0.160
LSD <sub>0.05</sub>		0.05	0.03	0.004	0.008	0.02	0.04	0.001	0.001
<b>Salinity levels</b>									
Cont.		1.96	2.01	0.212	0.216	2.23	2.32	0.223	0.237
1000		2.00	2.01	0.226	0.215	2.27	2.29	0.231	0.234
2000		1.89	1.92	0.196	0.193	1.99	2.13	0.205	0.208
4000		1.83	1.26	0.180	0.121	1.86	1.30	0.187	0.128
LSD <sub>0.05</sub>		0.03	0.03	0.003	0.005	0.02	0.02	0.002	0.002
<b>Beneficial microorganisms</b>									
control		1.80	1.69	0.187	0.170	1.91	1.83	0.192	0.183
<i>B. subtilis</i>		1.96	1.84	0.200	0.183	2.16	2.07	0.207	0.212
AMF		1.92	1.80	0.211	0.195	2.11	2.03	0.224	0.237
<i>B.+ AMF</i>		1.99	1.86	0.216	0.197	2.17	2.11	0.224	0.237
LSD <sub>0.05</sub>		0.02	0.02	0.002	0.002	0.01	0.02	0.002	0.002
<b>Basil cultivars X Salinity levels</b>									
Local cultivar	Cont.	1.97	2.02	0.211	0.217	2.23	2.32	0.222	0.237
	1000	2.01	2.03	0.229	0.217	2.25	2.26	0.230	0.237
	2000	1.88	1.90	0.198	0.190	2.01	2.18	0.206	0.212
	4000	1.85	1.88	0.184	0.174	1.88	1.94	0.185	0.190
Nano Compatt	Cont.	2.02	2.05	0.222	0.228	2.24	2.33	0.231	0.242
	1000	2.06	2.06	0.232	0.229	2.31	2.33	0.238	0.247
	2000	1.95	1.96	0.204	0.207	1.99	2.19	0.209	0.219
	4000	1.87	1.89	0.186	0.188	1.88	1.96	0.190	0.193
Red Bordeaux	Cont.	1.89	1.97	0.203	0.204	2.23	2.30	0.217	0.227
	1000	1.91	1.94	0.217	0.200	2.24	2.27	0.225	0.221
	2000	1.82	1.89	0.186	0.181	1.98	2.04	0.200	0.192
	4000	1.76	0.00	0.170	0.000	1.82	0.00	0.187	0.000
LSD <sub>0.05</sub>		N.S	N.S	N.S	0.008	0.03	0.03	0.004	0.004
<b>Basil cultivars X Beneficial microorganisms</b>									
Local cultivar	Cont.	1.83	1.87	0.187	0.181	1.91	1.98	0.190	0.198
	<i>B.sub.</i>	1.95	2.00	0.203	0.193	2.18	2.25	0.204	0.212
	AMF	1.94	1.96	0.215	0.209	2.12	2.19	0.226	0.234
	<i>B.+ AMF</i>	1.98	2.01	0.217	0.215	2.16	2.28	0.222	0.232
Nano Compatt	Cont.	1.86	1.88	0.196	0.197	1.97	2.03	0.197	0.204
	<i>B.sub.</i>	2.02	2.04	0.207	0.212	2.17	2.25	0.215	0.222
	AMF	1.97	1.98	0.217	0.220	2.12	2.22	0.227	0.238
	<i>B.+ AMF</i>	2.05	2.07	0.225	0.224	2.17	2.32	0.229	0.238

Red Bordeaux	Cont.	1.71	1.33	0.179	0.133	1.86	1.48	0.189	0.148
	<i>B.sub.</i>	1.90	1.50	0.190	0.143	2.15	1.71	0.202	0.159
	AMF	1.85	1.47	0.203	0.153	2.11	1.68	0.219	0.168
	<i>B.+ AMF</i>	1.93	1.52	0.204	0.156	2.15	1.74	0.219	0.166
LSD <sub>0.05</sub>		N.S	N.S	0.003	0.004	0.02	N.S	0.003	0.003

Continuous Table(5):

Charact.		First season				Second season				
		N %		P %		N %		P %		
		1 <sup>st</sup> cut	2 <sup>nd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	
<b>Salinity levels X Beneficial microorganisms</b>										
Cont.	Cont.	1.76	1.86	0.197	0.202	1.93	2.10	0.204	0.216	
	<i>B.sub.</i>	2.04	2.06	0.206	0.211	2.33	2.38	0.216	0.236	
	AMF	1.98	2.02	0.218	0.223	2.32	2.33	0.234	0.246	
	<i>B.+ AMF</i>	2.05	2.11	0.227	0.229	2.35	2.45	0.239	0.250	
1000	Cont.	1.85	1.86	0.204	0.196	2.03	2.05	0.210	0.214	
	<i>B.sub.</i>	2.04	2.06	0.221	0.211	2.35	2.37	0.227	0.227	
	AMF	2.00	2.02	0.234	0.227	2.31	2.33	0.242	0.243	
	<i>B.+ AMF</i>	2.09	2.09	0.246	0.227	2.37	2.40	0.245	0.249	
2000	Cont.	1.82	1.83	0.183	0.173	1.90	1.93	0.183	0.183	
	<i>B.sub.</i>	1.91	1.96	0.195	0.191	2.08	2.19	0.199	0.204	
	AMF	1.86	1.91	0.205	0.205	1.97	2.15	0.220	0.224	
	<i>B.+ AMF</i>	1.95	1.96	0.202	0.202	2.02	2.60	0.218	0.221	
4000	Cont.	1.76	1.22	0.166	0.111	1.78	1.23	0.172	0.120	
	<i>B.sub.</i>	1.86	1.28	0.179	0.118	1.91	1.34	0.185	0.125	
	AMF	1.82	1.25	0.188	0.127	1.85	1.30	0.197	0.134	
	<i>B.+ AMF</i>	1.86	1.29	0.188	0.129	1.90	1.34	0.195	0.134	
LSD <sub>0.05</sub>		N.S	N.S	0.004	0.004	0.03	0.03	0.003	0.004	
<b>Basil cultivars X Salinity levels X Beneficial microorganisms</b>										
Local cultivar	Cont.	Cont.	1.77	1.89	0.197	0.207	1.92	2.16	0.204	0.216
		<i>B.sub.</i>	2.04	2.06	0.204	0.214	2.32	2.36	0.211	0.234
		AMF	2.00	2.02	0.217	0.221	2.32	2.32	0.238	0.248
		<i>B.+AMF</i>	2.06	2.11	0.227	0.227	2.34	2.44	0.234	0.248
	1000	Cont.	1.89	1.90	0.204	0.200	1.98	1.98	0.207	0.217
		<i>B.sub.</i>	2.06	2.08	0.227	0.207	2.34	2.36	0.227	0.227
		AMF	2.02	2.04	0.238	0.227	2.32	2.30	0.248	0.256
		<i>B.+AMF</i>	2.08	2.09	0.248	0.234	2.36	2.40	0.238	0.248
	2000	Cont.	1.84	1.86	0.183	0.166	1.90	1.94	0.180	0.183
		<i>B.sub.</i>	1.90	1.94	0.197	0.186	2.16	2.26	0.197	0.204
		AMF	1.87	1.89	0.208	0.200	1.96	2.20	0.221	0.234
		<i>B.+AMF</i>	1.92	1.90	0.204	0.207	2.02	2.30	0.227	0.227
	4000	Cont.	1.80	1.82	0.165	0.152	1.82	1.82	0.170	0.176
		<i>B.sub.</i>	1.87	1.90	0.183	0.166	1.90	2.00	0.183	0.183
		AMF	1.85	1.87	0.197	0.186	1.87	1.94	0.197	0.197
		<i>B.+AMF</i>	1.87	1.92	0.190	0.193	1.92	1.98	0.190	0.204
Nano Compatt	Cont.	Cont.	1.80	1.89	0.207	0.214	1.96	2.13	0.211	0.226
		<i>B.sub.</i>	2.11	2.11	0.217	0.227	2.34	2.38	0.227	0.248
		AMF	2.04	2.05	0.227	0.234	2.31	2.34	0.238	0.255
		<i>B.+AMF</i>	2.11	2.17	0.238	0.248	2.36	2.48	0.248	0.261
	1000	Cont.	1.92	1.92	0.211	0.207	2.18	2.19	0.216	0.217
		<i>B.sub.</i>	2.11	2.09	0.224	0.227	2.36	2.36	0.238	0.237
		AMF	2.06	2.07	0.238	0.238	2.30	2.34	0.248	0.261
		<i>B.+AMF</i>	2.16	2.16	0.255	0.234	2.38	2.44	0.251	0.254
	2000	Cont.	1.89	1.89	0.190	0.186	1.92	1.96	0.186	0.190
		<i>B.sub.</i>	1.96	2.00	0.204	0.207	2.02	2.24	0.204	0.214
		AMF	1.92	1.94	0.211	0.221	2.00	2.24	0.225	0.234
		<i>B.+AMF</i>	2.04	2.01	0.211	0.214	2.02	2.32	0.221	0.238
	4000	Cont.	1.82	1.82	0.176	0.180	1.80	1.87	0.176	0.183
		<i>B.sub.</i>	1.90	1.94	0.183	0.186	1.96	2.02	0.190	0.190
		AMF	1.84	1.87	0.190	0.193	1.87	1.94	0.197	0.204
		<i>B.+AMF</i>	1.90	1.94	0.197	0.193	1.90	2.02	0.197	0.197
Cont.	Cont.	1.72	1.80	0.186	0.186	1.90	2.02	0.197	0.207	
	<i>B.sub.</i>	1.96	2.02	0.197	0.193	2.34	2.40	0.211	0.227	
	AMF	1.89	1.99	0.211	0.213	2.32	2.34	0.227	0.234	
	<i>B.+AMF</i>	1.99	2.07	0.217	0.222	2.34	2.42	0.234	0.238	

Red Bordeaux	1000	Cont.	1.75	1.77	0.197	0.180	1.94	1.99	0.207	0.207
		<i>B.sub.</i>	1.94	2.01	0.211	0.200	2.34	2.38	0.217	0.217
		AMF	1.91	1.96	0.227	0.207	2.32	2.34	0.238	0.234
		<i>B.+AMF</i>	2.04	2.02	0.234	0.214	2.36	2.36	0.238	0.227
	2000	Cont.	1.72	1.73	0.176	0.166	1.87	1.90	0.183	0.176
		<i>B.sub.</i>	1.87	1.95	0.183	0.180	2.06	2.06	0.197	0.193
		AMF	1.80	1.91	0.197	0.193	1.96	2.02	0.214	0.204
		<i>B.+AMF</i>	1.89	1.97	0.190	0.186	2.02	2.16	0.207	0.197
	4000	Cont.	1.65	0.00	0.156	0.00	1.72	0.00	0.170	0.00
		<i>B.sub.</i>	1.82	0.00	0.170	0.00	1.87	0.00	0.183	0.00
		AMF	1.78	0.00	0.176	0.00	1.82	0.00	0.197	0.00
		<i>B.+AMF</i>	1.80	0.00	0.176	0.00	1.87	0.00	0.197	0.00
LSD <sub>0.05</sub>			N.S	N.S	N.S	0.007	0.05	0.06	0.006	0.006

**Table (6): Effect of inoculation with *Bacillus subtilis* and arbuscular mycorrhizal fungi (AMF) on potassium and sodium % in dry leaves of the three sweet basil cultivars under salt stress condition during 2007 and 2008 seasons.**

Charact.	Treat.	First season				Second season				
		K %		Na %		K %		Na %		
		1 <sup>st</sup> cut	2 <sup>nd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	
Basil varieties										
	Local cultivar	1.14	1.27	1.03	1.01	1.20	1.32	1.04	1.03	
	Nano Compatt	1.15	1.27	1.00	0.75	1.21	1.32	1.03	0.75	
	Red Bordeaux	1.02	0.90	1.04	1.05	1.09	1.20	1.04	1.04	
	LSD <sub>0.05</sub>	0.001	0.005	0.004	0.002	0.003	0.003	0.003	0.005	
Salinity levels										
	Cont.	1.14	1.28	0.96	0.95	1.22	1.36	0.96	0.96	
	1000	1.14	1.27	0.99	0.99	1.23	1.33	1.00	1.00	
	2000	1.08	1.23	1.06	1.06	1.16	1.29	1.06	1.07	
	4000	1.03	0.81	1.09	0.74	1.07	0.84	1.11	0.73	
	LSD <sub>0.05</sub>	0.001	0.003	0.002	0.004	0.002	0.003	0.004	0.003	
Beneficial microorganisms										
	control	1.05	1.10	1.07	0.98	1.11	1.13	1.09	0.98	
	<i>B. subtilis</i>	1.10	1.14	1.03	0.94	1.17	1.19	1.07	0.95	
	AMF	1.12	1.17	1.00	0.92	1.19	1.24	1.00	0.92	
	<i>B.+AMF</i>	1.13	1.17	0.99	0.90	1.21	1.25	1.00	0.91	
	LSD <sub>0.05</sub>	0.001	0.001	0.002	0.003	0.003	0.003	0.003	0.001	
Basil cultivars X Salinity levels										
	Local cultivar	Cont.	1.15	1.29	0.95	0.94	1.23	1.40	0.96	0.94
		1000	1.18	1.29	0.99	0.99	1.25	1.35	1.00	1.01
		2000	1.13	1.26	1.06	1.06	1.19	1.30	1.06	1.06
		4000	1.09	1.23	1.10	1.10	1.15	1.23	1.12	1.11
	Nano Compatt	Cont.	1.19	1.31	0.93	0.93	1.25	1.35	0.96	0.96
		1000	1.20	1.33	0.97	0.96	1.27	1.37	0.99	0.96
		2000	1.11	1.25	1.04	1.04	1.18	1.30	1.05	1.05
		4000	1.08	1.20	1.08	1.09	1.14	1.28	1.10	1.09
	Red Bordeaux	Cont.	1.09	1.23	1.00	0.99	1.18	1.31	0.97	0.98
		1000	1.06	1.20	1.00	1.03	1.15	1.28	1.01	1.02
		2000	1.01	1.16	1.07	1.08	1.11	1.27	1.07	1.10
		4000	0.92	0.00	1.10	1.12	0.92	0.00	1.12	0.00
	LSD <sub>0.05</sub>		0.002	0.005	0.003	0.007	0.004	0.005	0.006	0.005
Basil cultivars X Beneficial microorganisms										
	Local cultivar	Cont.	1.08	1.21	1.07	1.07	1.15	1.24	1.10	1.06
		<i>B.sub.</i>	1.12	1.25	1.04	1.02	1.22	1.31	1.05	1.04
		AMF	1.18	1.31	1.01	0.98	1.22	1.36	1.01	1.00
		<i>B.+AMF</i>	1.18	1.29	0.99	0.96	1.25	1.36	1.01	1.00
	Nano Compatt	Cont.	1.11	1.24	1.05	0.79	1.17	1.24	1.10	0.78
		<i>B.sub.</i>	1.16	1.28	1.01	0.76	1.22	1.32	1.04	0.75
		AMF	1.15	1.29	0.98	0.73	1.22	1.36	0.98	0.74
		<i>B.+AMF</i>	1.18	1.31	0.98	0.72	1.25	1.37	0.98	0.73
	Red Bordeaux	Cont.	0.95	1.10	1.10	1.09	1.02	0.92	1.09	1.09
		<i>B.sub.</i>	1.03	1.14	1.05	1.05	1.10	0.94	1.05	1.06

	AMF	1.04	1.17	1.02	1.04	1.12	1.00	1.01	1.03
	B.+	1.05	1.17	1.00	1.02	1.12	1.00	1.01	0.99
	AMF								
LSD <sub>0.05</sub>		0.002	0.002	0.004	0.005	0.005	0.006	0.006	0.003

**Continuous Table(6):**

Charact.		First season				Second season				
		K %		Na %		K %		Na %		
		1 <sup>st</sup> cut	2 <sup>nd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	
<b>Salinity levels X Beneficial microorganisms</b>										
Cont.	Cont.	1.10	1.25	1.03	1.02	1.17	1.27	1.04	1.01	
	<i>B.sub.</i>	1.13	1.27	0.97	0.96	1.21	1.35	0.99	0.98	
	AMF	1.16	1.30	0.93	0.92	1.24	1.40	0.93	0.94	
	B.+ AMF	1.16	1.32	0.91	0.91	1.26	1.41	0.90	0.91	
1000	Cont.	1.10	1.21	1.04	1.04	1.17	1.24	1.08	1.04	
	<i>B.sub.</i>	1.14	1.27	1.00	1.00	1.22	1.32	1.02	1.00	
	AMF	1.17	1.30	0.96	0.98	1.25	1.36	0.95	0.98	
	B.+ AMF	1.17	1.29	0.96	0.96	1.27	1.39	0.97	0.97	
2000	Cont.	1.03	1.18	1.10	1.11	1.11	1.22	1.12	1.11	
	<i>B.sub.</i>	1.09	1.24	1.07	1.07	1.15	1.27	1.07	1.07	
	AMF	1.10	1.24	1.04	1.04	1.18	1.34	1.02	1.05	
	B.+ AMF	1.11	1.25	1.02	1.01	1.19	1.33	1.05	1.04	
4000	Cont.	0.95	0.78	1.12	0.76	0.99	0.80	1.14	0.76	
	<i>B.sub.</i>	1.05	0.80	1.10	0.74	1.09	0.82	1.12	0.74	
	AMF	1.06	0.82	1.07	0.73	1.10	0.86	1.10	0.72	
	B.+ AMF	1.06	0.83	1.08	0.72	1.11	0.86	1.09	0.71	
LSD <sub>0.05</sub>		0.002	0.002	0.005	0.006	0.005	0.006	0.007	0.003	
<b>Basil cultivars X Salinity levels X Beneficial microorganisms</b>										
Local cultivar	Cont.	Cont.	1.09	1.24	1.01	1.04	1.18	1.31	1.01	1.04
		<i>B.sub.</i>	1.13	1.27	0.97	1.01	1.20	1.39	0.97	1.01
		AMF	1.20	1.33	0.93	0.97	1.25	1.43	0.95	0.95
		B.+AMF	1.20	1.33	0.91	0.93	1.27	1.47	0.91	0.91
	1000	Cont.	1.11	1.22	1.04	1.06	1.20	1.24	1.10	1.06
		<i>B.sub.</i>	1.15	1.27	1.01	1.01	1.22	1.33	1.01	1.04
		AMF	1.22	1.35	0.95	1.04	1.29	1.43	0.97	1.01
		B.+AMF	1.22	1.33	0.97	1.01	1.31	1.41	0.97	0.97
	2000	Cont.	1.07	1.20	1.10	1.10	1.13	1.22	1.12	1.12
		<i>B.sub.</i>	1.11	1.25	1.08	1.06	1.18	1.31	1.10	1.06
		AMF	1.15	1.29	1.06	1.06	1.20	1.34	1.01	1.04
		B.+AMF	1.17	1.31	1.01	1.04	1.24	1.33	1.06	1.01
	4000	Cont.	1.05	1.18	1.12	1.15	1.09	1.20	1.15	1.15
		<i>B.sub.</i>	1.07	1.22	1.10	1.12	1.15	1.22	1.12	1.12
		AMF	1.13	1.25	1.10	1.10	1.18	1.24	1.12	1.10
		B.+AMF	1.11	1.27	1.08	1.10	1.18	1.25	1.10	1.08
Nano Compatt	Cont.	Cont.	1.15	1.31	0.97	1.01	1.20	1.27	1.04	1.01
		<i>B.sub.</i>	1.20	1.33	0.93	0.95	1.26	1.39	1.01	0.97
		AMF	1.20	1.33	0.91	0.91	1.25	1.41	0.91	0.95
		B.+AMF	1.22	1.35	0.91	0.91	1.29	1.43	0.88	0.91
	1000	Cont.	1.18	1.25	1.01	1.04	1.22	1.25	1.08	1.04
		<i>B.sub.</i>	1.20	1.31	0.97	0.94	1.27	1.37	1.01	1.01
		AMF	1.22	1.35	0.95	0.93	1.29	1.35	0.91	0.97
		B.+AMF	1.22	1.33	0.93	0.91	1.31	1.41	0.97	1.01
	2000	Cont.	1.07	1.22	1.10	1.10	1.13	1.24	1.12	1.08
		<i>B.sub.</i>	1.13	1.27	1.04	1.06	1.18	1.26	1.04	1.06
		AMF	1.11	1.25	1.01	1.01	1.20	1.35	1.02	1.04
		B.+AMF	1.13	1.27	1.01	0.97	1.22	1.35	1.04	1.04
	4000	Cont.	1.03	1.18	1.12	1.12	1.11	1.22	1.15	1.12
		<i>B.sub.</i>	1.11	1.19	1.10	1.10	1.15	1.25	1.10	1.10
		AMF	1.07	1.22	1.04	1.08	1.13	1.33	1.08	1.06
		B.+AMF	1.11	1.22	1.06	1.05	1.18	1.31	1.06	1.06
	Cont.	Cont.	1.05	1.20	1.10	1.01	1.13	1.24	1.06	0.97
		<i>B.sub.</i>	1.07	1.22	1.01	0.93	1.17	1.27	0.97	0.95
		AMF	1.09	1.24	0.97	0.88	1.20	1.37	0.95	0.93
		B.+AMF	1.13	1.27	0.91	0.91	1.22	1.35	0.91	0.91
	Cont.	1.03	1.16	1.06	1.01	1.11	1.22	1.06	1.01	

Red Bordeaux	1000	<i>B.sub.</i>	1.07	1.22	1.01	1.01	1.15	1.25	1.02	0.95
		AMF	1.07	1.20	0.97	0.97	1.18	1.31	0.97	0.95
		<i>B.+AMF</i>	1.07	1.22	0.97	0.95	1.18	1.35	0.96	0.93
	2000	Cont.	0.96	1.12	1.10	1.12	1.07	1.22	1.10	1.12
		<i>B.sub.</i>	1.03	1.18	1.08	1.10	1.11	1.24	1.08	1.10
		AMF	1.03	1.18	1.06	1.06	1.13	1.33	1.01	1.08
	4000	<i>B.+AMF</i>	1.03	1.18	1.04	1.04	1.20	1.31	1.06	1.08
		Cont.	0.77	0.00	1.12	0.00	0.77	0.00	1.14	0.00
		<i>B.sub.</i>	0.96	0.00	1.10	0.00	0.96	0.00	1.12	0.00
		AMF	0.99	0.00	1.08	0.00	0.99	0.00	1.10	0.00
		<i>B.+AMF</i>	0.96	0.00	1.10	0.00	0.96	0.00	1.10	0.00
		LSD <sub>0.05</sub>		0.003	0.004	0.008	0.011	0.009	1.131	0.012

#### 4. Discussion

It was clearly seen from the results that Nano Compatt cultivar surpassed those of Red Bordeaux and Local cultivars in plant height, number of branches, fresh and dry weights, N, P and K % and showed lower Na %. While, Local cultivar surpassed the other two cultivars in oil % and oil yield. Moreover, Red Bordeaux cultivar was more sensitive to salinity stress than the other two basil cultivars. These variations in the response of the three cultivars to water deficit was attributed to the genetic ability of the resistant cultivar to undergo certain modifications in their metabolic pathway which causes increase in osmotic potential thus increasing cell turgor and eventually growth (Abdalla and Khoshiban, 2007).

Saline soils and saline irrigations constitute a serious production problem for most crops as saline conditions are known to suppress plant growth. The present study demonstrates that high salinity level (4000 ppm) adversely affected the growth attributes of the three sweet basil cultivars. These results are in agreement with those obtained by Shannon and Grieve (1999); Han and Lee (2005); Al-Karaki (2006); Soha (2006); Yildirim *et al.* (2006) and Turkmen *et al.* (2008). The observed reduction in growth criteria in basil plants under high salinity levels may be attributed to that salinity reduced cell size and the number of cells per unit area (Storganov; 1962 and Greenway and Munns; 1980), also high salt concentration adversely affects enzymatic processes through some interaction of salt and some organic substances of the cell (Oertel, 1966). In addition, Waisel (1969) suggested that the decrease in dry matter production of plants under stress conditions was due to reduction in photosynthesis and increased respiration. Ghazi (1976) added that growth depression under salinity conditions might be due to decrease of root growth and shortage of water absorption. Moreover, the increase in external ion concentration may lead also to reduction of both cell development and cell turgor as well as inhibition of enzyme activity and photosynthesis. In addition, the

suppressive effect of salinity may be caused by the disturbance in mineral uptake (Khadr *et al.*, 1980), protein synthesis (Tseniv *et al.*, 1983) or photosynthetic and carbohydrates metabolism (Patil *et al.*, 1983). Salinity also affects soil ions such as Na and Cl, decreasing water potential and disturbing ion balance concentration; therefore the uptake, transportation and usage of plant nutrients are negatively affected by salinity, for example an increase in soil Na concentration reduces the K uptake of plants (Turkmen *et al.*, 2008). The stimulatory effect of low salinity levels on growth of some plants in this study were recorded by several authors as Maraim (1990) who recorded growth stimulation at intermediate salinity, associated with ions accumulation and increased shoot succulence occurred in *Sporobolus virginicus*. Naidoo *et al.* (1995) recorded stimulatory effect of moderate salinity on growth of some plants, these may be due to improve shoot osmotic status as a result of increasing ions uptake. Similar results were reported by Ashraf and Sharif (1998) whom recorded good growth with slightly increase in shoot dry weight under moderate saline conditions, due to their lower accumulation of Na, Cl and Ca in their leaves.

To alleviate the negative effect of soil salinity on basil physiological responses we co-inoculated the three studied cultivars with two beneficial microorganisms, Arbuscular Mycorrhizal Fungus (AMF) and/or *B. subtilis*. Results of the measurement of growth response, oil %, oil yield and minerals uptake are given in Tables 2, 3, 4, 5 & 6 which showed that all previously mentioned characters with or without stalinization treatments were significantly increased by all beneficial microorganisms treatments, and these biological treatments ameliorated the deleterious effect of salinity. These results are in greeet accordance with those obtained by Glick *et al.* (1997); Mayak *et al.* (2004b); Yildirim and Taylor (2005); Barassi *et al.* (2006); and Wei Liu *et al.* (2010). Mycorrhizal colonization showed generally more pronounced effects than *B. subtilis*, dual inoculation

with *B. subtilis* and mycorrhizae provided higher tolerance to salinity compared with the individual treatment and untreated ones, where pots treated with *B. subtilis*+ AMF showed greater plant height, number of branches, fresh and dry weight, oil % and yield as well as N, P, K % and lower Na % compared to the other treatments under saline conditions. This result was in great accordance with those obtained by Al-Karaki *et al.*, 2001; Mayak *et al.*, 2004; Saleh *et al.*, 2005; Al-Karaki, 2006; and Zuccarini and Okurowska, 2008. These results may be due to mycorrhizal symbiosis is an important factor in helping plants to cope with adverse environmental conditions, the major reason for increasing growth can be attributed to the ability of plants in associations with AMF to uptake some nutrients efficiently (Smith *et al.*, 1992), AMF in saline conditions could have partly to increase the uptake of P, N, K, Ca, Zn, Cu, this may be due to the soil pores that can be penetrated by AMF hyphae are perhaps an order of magnitude smaller than those available to roots (Smith and Read, 1997), the effect of AMF on plant Na content was clearly observed that AMF decreased the Na uptake of plants, AMF could protect plants from Na toxicity either by regulating Na uptake from the soil or by accumulating it in root (Rabie and Almadini, 2006), this is important because the lower the Na uptake the higher the salinity tolerance. The increase in nutrient uptake proposed to be due to increasing affinity to a particular ion and lowering the threshold concentration for absorption (Bolan *et al.*, 1987) and by exploring greater soil volume and increasing root surface area (Rhodas and Gerdemann, 1980). The induction of plant resistance to salt stress is provided through a discriminated absorption of the ions present in the circulating solution, so that sodium  $\text{Na}^+$  and chloride  $\text{Cl}^-$  uptake is kept at tolerable levels (Al-Karaki and Hammad, 2001), and through better balance of mineral nutrient uptake (Graham, 1986). Also *B. subtilis* enhanced the stability of the cell membrane, raised the root vigor of plant under salt stress, improved photosynthesis under salt stress by increasing the net photosynthetic rate and the stomatic conductance. PGPR strains such as *B. subtilis* can produce bacterial exopolysaccharides (EPSs) that bind cations, including  $\text{Na}^+$  (Geddie and Sutherland, 1993), it may be envisaged that increasing the population density of EPS-producing bacteria in the root zone would decrease the content of  $\text{Na}^+$  available for plant uptake and thus help alleviating salt stress in plants growing in saline environments (Ashraf *et al.*, 2004). So the application of *Bacillus subtilis* is a safe and promising way to relieve salt stress in crop

production (Bochow *et al.*, 2001; Ashraf *et al.*, 2004; Saleh *et al.*, 2005 and Wei Liu *et al.*, 2010).

## 5. References

1. **A.O.A.C., 1990.** Official methods of analysis. Association of official analytical chemists. 15<sup>th</sup> ed. Washington D.C.,USA.
2. **Abdalla, M.M and N.H. El-Khoshiban, 2007.** The influence of water stress on growth, relative water content, photosynthetic pigments, some metabolic and hormonal contents of two *Triticum aestivum* cultivars. *Journal of Applied Sciences Research*, 3(12): 2062-2074.
3. **Al-Karaki, G.N. and A. Al-Raddad, 1997.** Effect of arbuscular mycorrhizal fungi and drought stress on growth and nutrient uptake of two wheat genotypes differing in drought resistance. *Mycorrhiza*, 7:87-88.
4. **Al-Karaki, G.N. and R. Hammad, 2001.** Mycorrhizal influence on fruit yield and mineral content of tomato grown under salt stress. *J. Plant Nutrition*, 24(8):1311-1323.
5. **Al-Karaki, G.N., 2006.** Nursery inoculation of tomato with arbuscular mycorrhizal fungi and subsequent performance under irrigation with saline water. *Scientia Horticulturae*, 109:1-7.
6. **Al-Karaki, G.N., R. Hammad and M. Rusan, 2001.** Response of two tomato cultivars differing in salt tolerance to inoculation with mycorrhizal fungi under salt stress. *Mycorrhiza*, 11:43-47.
7. **Apse, M.P., G.S. Dharon, W.A. Snedden and E. Bumerold, 1999.** Salt tolerance conferred by overexpression of a vacuolar  $\text{Na}^+/\text{H}^+$  antiport in Arabidopsis. *Science*, 285:1256-1258.
8. **Ashraf, M. and R. Sharif, 1998.** Does salt tolerance vary in a potential oil seed crop *Brassica carinata* at different growth stages. *Journal of Agronomy and crop science*, 181: 2, 103-115.
9. **Ashraf, M., S. Hasnain, O. Berge and T. Mahmood, 2004.** Inoculation wheat seedlings with exopolysaccharide-producing bacteria restricts sodium uptake and stimulates plant growth under salt stress. *Biol. Fertil. Soils*, 40:157-162.
10. **Barassi, C.A., G. Ayrault, C.M. Creus, R.J. Sueldo, and M.T. Sobrero, 2006.** Seed inoculation with *Azospirillum mitigates* NaCl effects on lettuce. *Sci. Hortic.* 109, 8–14.
11. **BjÖrkman, T., L.M. Blanchard, and G.E. Harman, 1998.** Growth enhancement of shrunken-2 (sh2) sweet corn by *Trichoderma harzianum* 1295-22: effect of environmental stress. *J. Am. Soc. Hortic. Sci.* 123, 35–40.
12. **Black, C.A., D.D. Evans and L.E. Ensminger, 1965.** Methods of soil analysis. *Agronomy, J. Amer. Soc. Agron. Inc.* Publ., Madison, Wisconsin, U.S.A.
13. **Bochow, H., S.F. El-Sayed, H. Junge, A. Stavropoulou and G. Schmiedeknecht, 2001.** Use of *Bacillus subtilis* as biocontrol agent. IV. Salt-stress tolerance induction by *Bacillus subtilis* FZB24 seed treatment in tropical vegetable field crops, and its mode

- of action. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz*, 108 (1):21-30.
14. **Bolan, N.S., A.D. Robson and N.J. Barrow, 1987.** Effect of VAM on the availability of iron phosphates to plants. *Plant Soil*, 99:401-410.
  15. **Boutraa, A., and F.E. Sander, 2001.** Influence of water stress on grain yield and vegetative growth of two cultivars of bean (*Phaseolus vulgaris* L.). *Journal of Agronomy and Crop Science*, 187 (4) page: 251.
  16. **Cantrell, I.C. and R.G. Linderman, 2001.** Preinoculation of lettuce and onion with VA mycorrhizal fungi reduces deleterious effects of soil salinity. *Plant Soil*, 233:269-281.
  17. **Copeman, R.H., C.A. Martin and J.C. Stutz, 1996.** Tomato growth in response to salinity and mycorrhizal fungi from saline or non-saline soils. *Hortic Sci.*, 31(3):341-344.
  18. **Danneberg, G., C. Latus, W. Zimmer, B. Hundeshagen, H.J. Schneider-Poetsch and H. Bothe, 1992.** Influence of vesicular-arbuscular mycorrhiza on phytohormone balances in maize (*Zea mays* L.). *J. Plant Physiol.*, 141:33-39.
  19. **Demir S, and E. Onour, 1999.** Glomus intraradices Schenck&Smith: A hopeful vesicular-arbuscular mycorrhizal fungus (VAM) determined in soils of Türkiye. *J. Turk. Phytopat.* 28: 33-34.
  20. **Dixon, R.K., V.K. Garg and M.V. Rao, 1993.** Inoculation of Leucaena and prosopis seedlings with Glomus and Rhizobium species in saline soil: rhizosphere relations and seedlings growth. *Arid Soil Res. Rehabil.* 7:133-144.
  21. **Egamberdiyeva, D. and G. H flich, 2004.** Effect of plant growth-promoting bacteria on growth and nutrient uptake of cotton and pea in a semi-arid region of Uzbekistan. *J. Arid Environments*, 56:293-301.
  22. **El-Saidi, M.T., 1997.** Salinity and its effect on growth , yield and some physiological processes of crop plants. In: Jaiwal PK, Singh R., Gulati A. (eds) Strategies for improving salt tolerance in higher plants. Scientific Publications, Jodhpur, pp. 40-55.
  23. **Geddie, J.L. and I.W. Sutherland, 1993.** Uptake of metals by bacterial polysaccharides. *J. App. Bacteriol.*, 74: 467-472.
  24. **Ghazi, S.M., 1976.** Physiological studies of cycocel and alar in relation to salt tolerance on *Vicia faba* plants. Ph. D. Thesis, Ain Shams Univ., Egypt.
  25. **Glick, B.R. and Y. Bashan, 1997.** Genetic manipulation of plant growth-promoting bacteria to enhance biocontrol of fungal phytopathogens. *Biotechnol. Adv.* 15:353-378.
  26. **Glick, B.R., C. Liu, S. Ghosh, and E.B. Dumbrof, 1997.** Early development of canola seedlings in the presence of the plant growth-promoting rhizobacterium *Pseudomonas putida* GR12-2. *Soil Biol. Biochem.* 29 (8),1233–1239.
  27. **Graham, J.H., 1986.** Citrus mycorrhizae: Potential benefits and interactions with pathogens. *Hort .Science*, 21:1302-1306.
  28. **Greenway, H. and S.R. Munns, 1980.** Mechanisms of salt tolerance in non halophytes. *Annals Review of Plant Physiology*, 31: 149-159.
  29. **Grichko, V.P., and B.R. Glick, 2001.** Amelioration of flooding stress by ACC deaminase-containing plant growth-promoting bacteria. *Plant Physiol.Biochem.* 39, 11–17.
  30. **Guenther, E., 1961.** The Essential Oils. D. Van Nostrand Co., Vol. III. 4<sup>th</sup> ed. New York , London .
  31. **Hamdy, A., 1990.** Management practices under saline water irrigation. *Acta Horticulturae*, 278:745-754.
  32. **Han, H.S. and K.D. Lee, 2005.** Physiological responses of soybean-inoculation of *Bradyrhizobium japonicum* with PGPR in saline soil conditions. *Res. J. Agric. and Biol. Sci.*, 1(3): 216-221.
  33. **Harman, G.E., and T. Bjo'rkman, 1998.** Potential and existing uses of Trichoderma and Gliocladium for plant disease control and plant growth enhancement. In: Harman, G.E., Kubicek, C.P. (Eds.), Trichoderma and Gliocladium, vol. 2. Taylor & Francis, London, United Kingdom, pp. 229–265.
  34. **Hopkins, W.G., 1999.** Introduction to plant physiology. Wiley, New York.
  35. **Jackson, M.L., 1958.** Soil chemical analysis. Prentice Hall Inc., Englewood Cliffs. N. J.
  36. **Jackson, M.L., 1973.** Soil Chemical Analysis. Prentice-Hall, Inc. Englewood Cliffs, N.J. New Delhi, India.
  37. **Jain, R.K., K. Paliwal, R.K. Dixon and D.H. Gjerstad, 1989.** Improving productivity of multipurpose trees growing on substandard soils of India. *J. For.*, 87:38-42.
  38. **Khadr, A.A., N. Nawagy and N.M. Hussein, 1980.** Differential tolerance of wheat, barley, broad bean and pea seedlings to various levels of salinity. *Bull. Nat. Res. Cent. Egypt*, (5): 283-289.
  39. **Kloepper, J.W. and M.N. Schroth, 1978.** Plant growth-promoting rhizobacteria on radishes. IV. International Conference on Plant Pathogenic Bacteria. Angers France, 2:879-882.
  40. **Maraim, K.B., 1990.** Physiological parameters of salinity tolerance in C4 tufgrasses. Dissertation Abst. International, B. Sci. and Engineerings, 51: 2, 4848.
  41. **Mayak, S., T. Tirosh and B.R. Glick, 2004.** Plant growth-promoting bacteria confer resistance in tomato plants to salt stress. *Plant Physiology and Biochemistry*, 42:565-572.
  42. **Mayak, S., T. Tirosh, and B.R. Glick, 2004a.** Plant growth-promoting bacteria that confer resistance to water stress in tomatoes and pepper. *Plant Sci.* 166, 525–530.
  43. **Mayak, S., T. Tirosh, and B.R. Glick, 2004b.** Plant growth-promoting bacteria confer resistance in tomato plants to salt stress. *Plant Physiol. Biochem.* 42, 565–572.
  44. **Naidoo, Y., J. Jahnke and D.J. Von Willert, 1995.** Gas exchange responses of the C4 grass *Sporobolus virginicus* (Poaceae) to salinity stress. In Khan, AM and

- Ungar, I.A. (Eds), Biology of salt tolerant plants, Pf. 121-130 Karachi, University of Karachi.
45. **Noel, T.C., C. Sheng, C.K. Yost, R.P. Pharis and M.F. Hynes, 1996.** *Rhizobium leguminosarum* as a plant growth promoting rhizobacterium: direct growth promotion of canola and lettuce. *Can. J. Microbiol.*, 42: 279-283.
  46. **Oertil, J.J., 1966.** Effect of external salt concentration on water relations in plants. *Soil Sci.*, 102: 258-263.
  47. **Patil, T.M., P.B. Mirojkar, B.A. Hedge and G. Joshi, 1983.** Influence of salinity on morphology, rate of CO<sub>2</sub> assimilation, photosynthesis products and enzyme activities in sorghum hybrid CSH-S. *Indian J. of Plant Physiol.*, 26 (2): 153-162.
  48. **Qadir, M., A. Ghafoor and G. Murtaza, 2000.** Amelioration strategies for saline soils: a review. *Land Degrad Dev.*, 11:501-521.
  49. **Rabie, G.H., and A.M. Amadini, 2006.** Role of bioinoculant in development of salt-tolerance of *Vicia faba* plants under salinity stress. *Afr. J. Biotech.*, 4(3):210-222.
  50. **Rhoads, L.H. and J.W. Gerdemann, 1980.** Nutrient translocation in VAM. In Cellular Interactions in Symbiosis and Parasitism: Cooks, C.B., Pappas, P.W., Rudolph, E.D., Eds., Ohio State University Press: Columbus, OH, pp. 173-195.
  51. **Ruiz-Lozano, J.M. and R. Azcon, 1995.** Hyphal contribution to water uptake in mycorrhizal plants as affected by the fungal species and water status. *Physiologia Plantarum*, 95: 472-478.
  52. **Saleh, S.A., H. Heuberger and W.H. Schnitzler, 2005.** Alleviation of salinity effect on artichoke productivity by *Bacillus subtilis* FZB24, supplemental Ca and micronutrients. *J. App. Bot. Food Qual.*, 79: 24-32.
  53. **Shannon, M.C., and C.M Grieve, 1999.** Tolerance of vegetables to salinity. *Sci. Hort.* 78, 5-38.
  54. **Smith, S.E., A.D. Robson and L. k. Abott, 1992.** The involvement of Mycorrhizas in assessment of genetically dependent efficiency of nutrient uptake and use. *Plant Soil*, 146: 169-179.
  55. **Smith, S.E., and D.J. Read, 1997.** Mycorrhizal symbiosis. London: Academic Press.
  56. **Snedecor, G.W. and W.G. Cochran, 1980.** "Statistical Methods" 7th ed. Iowa state Univ., Iowa, USA.
  57. **Soha, E.K., 2006.** Physiological studies on sesame plant grown under saline water irrigation conditions. Ph.D Thesis, Botany Department, Cairo University.
  58. **Strogonov, B.P., 1962.** Physiological basis of salt tolerance of plants under different types of soil salinization. Izd. Akad. Nauk. USSR, Inst. Plant Physiol. Moscow, 364. Eng. Translation by Israel Program for Sci., Translation (1964).
  59. **Tseniv, E., V.V. Kabanov and B.P. Strogonov, 1983.** Effect of NaCl on the content and synthesis of different fractions of nucleic acid in pea shoot species. *Fixiologiya Rastenii USSR*, 30, 523-547.
  60. **Turkmen, O., S. Demir, and C. Erdinc, 2008.** Effect of two different AMF species on growth and nutrient content of pepper seedling grown under moderate salt stress. *African Journal of Biotechnology* 7(4):392-396.
  61. **Waisel, Y., G.A. Borger and T.T. Kozlowski, 1969.** Effects of PMA on stomatal movement and transpiration of excised *Betula papyrifera* march leaves. *Plant Physiol.*, 44: 685-690.
  62. **Wei Liu, Diankui Chen, Hanwen Yin, Min Song, Shirong Guo, 2010.** *Bacillus subtilis* for salt stress relief in vegetable. ISHS Acta Horticulturae 856: International Symposium on Vegetable Safety and Human Health. URL. [www.actahort.org](http://www.actahort.org)
  63. **Yildirim, E., A.G. Taylor, and T.D. Spittler, 2006.** Ameliorative effects of biological treatments on growth of squash plants under salt stress. *Scientia Horticulturae* 111, 1-6.
  64. **Yildirim, E., and A.G. Taylor, 2005.** Effect of biological treatments on growth of bean Plants under Salt Stress. *Annu. Rep. Bean Improv. Coop.* 48, 176-177
  65. **Zuccarini, P. and P. Okurowska, 2008.** Effects of mycorrhizal colonization and fertilization on growth and photosynthesis of sweet basil under salt stress. *Journal of Plant Nutrition*, 31(3): 497-513.

5/6/2011