

## Improving Nitrogen Utilization Efficiency by Potato (*Solanum tuberosum* L.)

### A. Influence of Nitrification Inhibitors in Combination with Different Nitrogen Sources on Reducing Nitrogen Losses, Improving Productivity and Chemical Composition

Abd El-Badea S. Ezzat<sup>1\*</sup>, Adel M. Abd El-Hameed<sup>2</sup>, Hamdino M. I. Ahmed<sup>1\*</sup> and Aml A. El-Awady<sup>1</sup>  
1 Vegetable Research Department, Horticulture Research Institute, Agric. Res. Center, Ministry Agric., EGYPT  
2 Plant Nutrition Department, Soil, Water and Environment Research Institute, Agric. Res. Center, Ministry Agric., EGYPT

\* Corresponding authors: [as\\_ezzataf@yahoo.com](mailto:as_ezzataf@yahoo.com); [hamdino@yahoo.com](mailto:hamdino@yahoo.com)

**ABSTRACT:** Two field experiments were conducted at Baramoon Research Station, Mansoura, Dakahlia Governorate, Egypt (+ 7m altitude, 30° 11' latitude and 28° 26' longitude), during Nili seasons of 2007/08 and 2008/09, to study the effect of soluble-N (ammonium nitrate; AN, ammonium sulphate; AS and urea; U) and/or slow-N (compost and nitroform) fertilizers with or without nitrification inhibitor (guanylthiourea, GTU) on reducing nitrogen loss, productivity, and chemical composition of potato cv. Cara. The obtained results indicate that GTU with compost<sub>50%</sub> and AS<sub>50%</sub> led to significant increases in all traits, except NO<sub>3</sub><sup>-</sup> and NO<sub>2</sub><sup>-</sup> accumulation, which was significantly decreased in potato tubers. Application of compost<sub>50%</sub> and AS<sub>50%</sub> with GTU had significant effect of most vegetative growth, quality, yield parameters and chemical composition of potato tubers in both season of the investigation. This treatment led to significant increase in plant height, plant dry weight, total and marketable of tuber yield and significant decrease in unmarketable tuber yield in both season of study. Application of compost<sub>50%</sub> and AS<sub>50%</sub> with GTU gave rise to a significant increase in tuber dry matter, starch and specific gravity and decrease nitrate and nitrite content in tubers in comparison with other treatments. The NPK uptake of potato tubers and nitrogen efficiency ratio in treatment amended with compost<sub>50%</sub> + AS<sub>50%</sub> and GTU was higher than the other treatments in two seasons. The highest value of residual NH<sub>4</sub>-N in soil was obtained from compost treatment alone followed by nitroform, whereas, AN gave the highest residual NO<sub>3</sub>-N compared with other treatments, in both seasons of study. It could be concluded that, application of nitrogen fertilizer in the form compost at the rate of 9 ton fed<sup>-1</sup> and ammonium sulphate at the rate of 90 kg fed<sup>-1</sup> with GTU (nitrification inhibitor) in potato fields were the most effective treatment for improvement nitrogen use efficiency with reducing the pollution of environment.

[Abd El-Badea S. Ezzat, Adel M. Abd El-Hameed, Hamdino M. I. Ahmed and Aml A. El-Awady (2011). Improving Nitrogen Utilization Efficiency by Potato (*Solanum tuberosum* L.). A. Influence of Nitrification Inhibitors in Combination with Different Nitrogen Sources on Reducing Nitrogen Losses, Improving Productivity and Chemical Composition. Nature and Science 2011; 9 (7): 26-33]. (ISSN: 1545-0740). <http://www.sciencepub.net>.

**Key words:** potato; nitrogen; nitrification inhibitors; guanylthiourea

### INTRODUCTION

Modern agricultural practices require a new concept of N-fertilizer management in order to optimize N-utilization and avoid N-losses. Nitrification inhibitors or "N-stabilizers" fit very well into this conception.

Nitrification inhibitors are compounds that delay bacterial oxidation of the ammonium-ion (NH<sub>4</sub><sup>+</sup>) by depressing over a certain period of time the activities of *Nitrosomonas* bacteria in the soil. They are responsible for the transformation of ammonium into nitrite (NO<sub>2</sub><sup>-</sup>) which is further changed into nitrate (NO<sub>3</sub><sup>-</sup>) by *Nitrobacter* and *Nitrosolobus* bacteria. The objective of using nitrification inhibitors is, therefore, to control leaching of nitrate by keeping nitrogen in the ammonia form longer, to prevent denitrification of nitrate-N and to increase the efficiency of nitrogen applied (Trenkel, 1997).

Nitrification inhibitors may reduce loss of fertilizer N from the root zone by reducing leaching and denitrification. This reduced N loss should be reflected in increased crop yields (Martin, *et al.*, 1993).

Guanylthiourea (GTU) is an efficient nitrification inhibitor and blocks the first step of nitrification for 1–3 months (depending on temperature). GTU is a non-toxic, water soluble compound and will be degraded to CO<sub>2</sub>, NH<sub>3</sub> and H<sub>2</sub>O without any residues. There are various possibilities to use GTU: addition to liquid manure temporarily prevents oxidation of ammonium nitrogen e.g. of slurry or waste water from potato starch production (Amberger and Germann-Bauer, 1990).

Several studies emphasized that treating ammonium fertilizers and organic manure with nitrification inhibitors helped in delaying nitrification of ammonium based fertilizers. By

preventing rapid formation of nitrate in the soil, leaching and denitrification losses of nitrogen are limited, thus increasing the efficiency of fertilizers. Lower concentration of nitrate in soil should result in less nitrate contamination of the ground water as well as reduced emission of nitrous oxide from denitrification (Laskshmanan and Prasad, 2004; Di and Cameron, 2004). Moreover, nitrification inhibitor not only decrease nitrate leaching and nitrous oxide emission as reported previously, but also decrease the leaching loss of cation nutrient such as  $\text{Ca}^{2+}$ ,  $\text{K}^+$  and  $\text{Mg}^{2+}$  (Di and Cameron, 2004).

Dachler (1993) found that potatoes showed clear positive effects in yield, tuber size and starch-yield and economically higher proceeds with the use of ammonium-sulfate-nitrate (ASN) + nitrification inhibitor (DCD) compared with ammonium-nitrate-lime (ANL) with or without DCD. Amberger (1989) mentioned that nitrification inhibitor, dicyandiamide (DCD), reduced nitrate leaching and increased yields and N uptake of potato plants.

Shoji *et al.* (2001) found that use of controlled release fertilizer (polyolefin coated urea) and/or nitrification inhibitor (dicyandiamide) to conserve air and water quality are basically due to maximizing nitrogen use efficiency (NUE), reducing the N fertilization rate and gave maximum tuber yields under center-pivot irrigated potato grown in a sandy field.

In field trials were conducted under various soil-climatic conditions in west and south Europe, in order to assess the effects of N-fertilizers with the new nitrification inhibitor DMPP (3,4-dimethylpyrazole phosphate) on yield and quality of various agricultural and horticultural crops, Pasda *et al.* (2002) showed that DMPP may increase the mean crop yield (grain yield: winter wheat +0.24 t ha<sup>-1</sup>; wetland rice +0.43 t ha<sup>-1</sup>; grain maize +0.24 t ha<sup>-1</sup>; tuber yield: potatoes +1.9 t ha<sup>-1</sup>, corrected sugar yield: sugar beets +0.24 t ha<sup>-1</sup>; biomass: carrots +1.9 t ha<sup>-1</sup>; lettuce +2.6 t ha<sup>-1</sup>, onions +1.0 t ha<sup>-1</sup>, radish +4.6 t ha<sup>-1</sup>; cauliflower +2.3 t ha<sup>-1</sup>; leek +3.1 t ha<sup>-1</sup>, and celeriac +1.9 t ha<sup>-1</sup>).

Vallejo *et al.* (2006) reported that nitrification inhibitor dicyandiamide (DCD) inhibited nitrification rates and reduced N<sub>2</sub>O and NO emissions from pig slurry by at least 83% and 77%, respectively. Similar finding were reported by Watanabe (2006). In the wheat growth experiment, Khalil *et al.* (2009) reported that the N<sub>2</sub>O losses were generally smaller, ranging from 0.16% to 0.27% of the total fertilization, than in the pot experiment, and the application of the urease inhibitor and the combined urease plus nitrification

inhibitors decreased N<sub>2</sub>O emissions by 23% to 59%.

The objective of this study was to estimate the productivity, quality and chemical composition of potato fertilized with different sources of N-fertilizers in sole or combined applications with or without nitrification inhibitor. It was also aimed to reduce nitrogen loss in soil and nitrate and nitrite contents in potato tubers.

## MATERIALS AND METHODS

Two field experiments were carried out at Baramoon Research Station, Mansoura, Dakahlia Governorate, Egypt (+ 7m altitude, 30° 11' latitude and 28° 26' longitude), during two successive winter growing seasons of 2007/08 and 2008/09. Potato (*Solanum tuberosum* L.) Cara cultivar was used in this study. Seed tubers were planted on 15<sup>th</sup> of October in both seasons of study. Plot area was 11.25 m<sup>2</sup>; consisted of 3 ridges; 5 m long; 75 cm wide, and 25 cm apart. The experimental soil was analyzed, using the methods described by Page *et al.* (1982), for the physical and chemical properties and the obtained data are shown in Table (1).

The following treatments have been tested: (1) Ammonium sulphate (20.5% N) (AS), (2) ammonium nitrate (33.5% N) (AN), (3) Urea (46.0% N) (U), (4) AS + Guanylthiourea (GTU), (5) AN + GTU, (6) Urea + GTU, (7) AS<sub>50%</sub> + AN<sub>50%</sub> + GTU, (8) AS<sub>50%</sub> + Urea<sub>50%</sub> + GTU, (9) Nitroform (38% N), (10) Compost (1.2% N), (11) Compost<sub>50%</sub> + AS<sub>50%</sub> + GTU, and (12) Compost<sub>50%</sub> + AN<sub>50%</sub> + GTU. The amount of added fertilizers was adjusted to a total N supply of 180 kg/feddan (feddan=4200 m<sup>2</sup>) for potato production. Guanylthiourea (GTU) as nitrification inhibitor was mixed with the fertilizers at the rate of 5% of added nitrogen dose. A complete randomized blocks design with three replicates was used in this respect.

Ammonium sulphate, ammonium nitrate, and urea were used as a soluble N-fertilizer, while, compost and nitroform were used as a slow release N-fertilizers.

The slow release-N was added to experimental soil before planting, whereas, soluble form of fertilizers was added at two equal doses, i. e. the first after emergence, and second dose was applied with 2<sup>nd</sup> irrigation. Single superphosphate (15.5% P<sub>2</sub>O<sub>5</sub>) was applied before planting at the rate of 75 kg P<sub>2</sub>O<sub>5</sub> fed<sup>-1</sup>. Potassium sulphate (48% K<sub>2</sub>O) was used as a source of potassium at the rate of 96 kg K<sub>2</sub>O fed<sup>-1</sup> and was added in two equal doses with the 2<sup>nd</sup> and 3<sup>rd</sup> irrigation. Other agricultural practices were conducted according to recommendations.

**Table 1: The main physical and chemical properties of the experimental site during the two growing seasons.**

| Some Physical properties | Values                 |                        | Some Chemical Properties | Values                 |                        |
|--------------------------|------------------------|------------------------|--------------------------|------------------------|------------------------|
|                          | 1 <sup>st</sup> season | 2 <sup>nd</sup> season |                          | 1 <sup>st</sup> season | 2 <sup>nd</sup> season |
| Sand (%)                 | 28.1                   | 27.9                   | pH* value                | 8.0                    | 7.9                    |
| Silt (%)                 | 31.8                   | 31.6                   | EC dSm <sup>-1</sup>     | 0.9                    | 0.9                    |
| Clay (%)                 | 40.1                   | 40.5                   | Total N (%)              | 0.03                   | 0.04                   |
| Texture class            | Clay-loam              | Clay-loam              | Available N (ppm)        |                        |                        |
|                          |                        |                        | NH <sub>4</sub> -N       | 23.37                  | 23.00                  |
|                          |                        |                        | NO <sub>2</sub> -N       | 0.162                  | 0.126                  |
|                          |                        |                        | NO <sub>3</sub> -N       | 13.21                  | 13.12                  |
| CaCO <sub>3</sub> (%)    | 3.2                    | 3.0                    | Available P (ppm)        | 13.3                   | 12.6                   |
| Organic matter (%)       | 1.8                    | 1.6                    | Available K (ppm)        | 304                    | 302                    |

\*pH: (1: 2.5 soil extract).

At 70 days after planting (DAP), a random sample of four plants was taken from each experimental unit to determine the growth parameters of potato plants (plant height and dry weight/plant). At the harvesting time (130 DAP), the total tuber yield, marketable and unmarketable yield per feddan was recorded. A representative sample of 10 to 15 healthy tubers from each experimental plot was selected from the largest sizes to obtain quality data (dry matter, specific gravity, starch, and nitrate and nitrite content) according to the methods described by (AOAC, 1990). Nitrogen, phosphorus and potassium accumulation in tubers (based on tuber dry weight and element percentage in tubers) were determined using the methods described by Cottenie *et al.*, (1982). For calculation of nitrogen efficiency ratio (NER), total tuber yield (kg fed<sup>-1</sup>) was divided by the amount of nitrogen in kg fed<sup>-1</sup>(=180 kg fed<sup>-1</sup>) (Aujla *et al.*, 1982). The soil samples were taken out from plots for residual available nitrogen at harvesting according to Black (1965).

Data obtained were subjected to statistical analysis by the technique of analysis of variance (ANOVA) according to Snedecor and Cochran (1982). The treatments mean were compared using Duncan multiple range test at 5 % level of probability as described by Steel and Torrie (1980).

## RESULTS AND DISCUSSION

### 1. Vegetative growth and tuber yield parameters:

Data presented in Table 2 demonstrate the effect of various treatments of slow release-N and soluble-N fertilizers with nitrification inhibitors (GTU) on vegetative growth parameters of potato plants and tuber yield characters. Significant effects on plant height, dry weight/plant, total, and marketable yields were obtained under the treatment where Compost<sub>50%</sub> + AS<sub>50%</sub> + GTU was applied in comparison to other treatments, in both seasons of study. On the other hand, application of

urea significantly increases in unmarketable yield, in both seasons.

It is quite obvious that dry matter accumulation and tuber yield were always much higher whenever organic manure was added. This trend being clearer with two sources of soluble N. On the other hand, a sole of slow or soluble fertilizers did not materially increase the parameters. In general the presence of nitrification inhibitor tended to increases in all studied parameters. Such result could be explained on the basis the efficiency of this material in decreasing nitrification of nitrogen, either added or produced through mineralization of organic compounds, and thus minimize its loss by leaching or volatilization (Amberger and Germann-Bauer, 1990; Martin, *et al.*, 1993; Vallejo *et al.*, 2006; Watanabe, 2006). These results are in agreement with those reported by Pasda *et al.* (2002) who showed that use of nitrification inhibitor increase the tuber yield of potatoes by 1.9 t ha<sup>-1</sup>.

### 2. Tuber quality characters:

Tuber quality as affected by N-source and nitrification inhibitor is given in Table 3. Results reveal that the application of GTU associated with AS<sub>50%</sub> or AN<sub>50%</sub> plus compost<sub>50%</sub> caused significant increase in tuber dry matter, specific gravity and starch content in tuber. In contrast, NO<sub>3</sub><sup>-</sup> and NO<sub>2</sub><sup>-</sup> accumulation was markedly decreased. These results were true in both seasons.

The pronounced positive effect on potato tuber quality may be attributed to decreasing N-losses (delaying the nitrification process) and increasing the N-use efficiency with nitrification inhibitor (Laskshmanan and Prasad, 2004; Di and Cameron, 2004), and consequently, increase the plant chance to absorb nitrogen and other nutrients (Table 4), thereby, produce good quality, especially where soils are poor in nitrogen and organic matter (Table 1). The negative effect of GTU associated with AS<sub>50%</sub> plus compost<sub>50%</sub> on NO<sub>3</sub><sup>-</sup> and NO<sub>2</sub><sup>-</sup> accumulation may be attributed to the role of GTU and compost in reducing NO<sub>3</sub><sup>-</sup> concentration in soil,

subsequently, gives the chance for plant to absorb more  $\text{NH}_4\text{-N}$ , thereby reduced  $\text{NO}_3^-$  accumulation

in plant (Bakr and Gawish, 1997).

**Table 2: Vegetative growth and tuber yield characters of potato as affected by nitrogen sources and nitrification inhibitors in 2007/08 and 2008/09 seasons.**

| Treatments | Plant height (cm) |          | Dry weight/plant (g) |         | Tuber yield (ton fed <sup>-1</sup> ) |         |            |         |              |         |
|------------|-------------------|----------|----------------------|---------|--------------------------------------|---------|------------|---------|--------------|---------|
|            | 2007/08           | 2008/09  | 2007/08              | 2008/09 | Total                                |         | Marketable |         | Unmarketable |         |
|            |                   |          |                      |         | 2007/08                              | 2008/09 | 2007/08    | 2008/09 | 2007/08      | 2008/09 |
| T1         | 47.33fgh          | 47.00de  | 35.71de              | 33.10 e | 11.373e                              | 11.920d | 10.740f    | 11.280d | 0.633 g      | 0.640 f |
| T2         | 48.17efg          | 49.00 c  | 34.18 f              | 31.18 f | 11.187e                              | 11.767d | 10.540f    | 11.100d | 0.647fg      | 0.667 f |
| T3         | 46.33 gh          | 52.00 a  | 30.72 h              | 34.24 d | 10.880f                              | 10.867f | 9.907 g    | 9.893 f | 0.973 a      | 0.973 a |
| T4         | 50.55cde          | 46.00 ef | 36.78 d              | 37.18 b | 13.267c                              | 12.874b | 12.553cd   | 12.161c | 0.713 e      | 0.713 e |
| T5         | 49.33def          | 46.33 ef | 35.00 ef             | 29.20 h | 12.573d                              | 12.233c | 11.833e    | 11.387d | 0.740 d      | 0.847 c |
| T6         | 46.00 h           | 43.00 g  | 28.65 i              | 29.57gh | 10.273g                              | 10.653f | 9.447 h    | 9.693fg | 0.826 b      | 0.960 a |
| T7         | 51.33bcd          | 50.00bc  | 38.76 c              | 36.18 c | 9.893 h                              | 10.193g | 9.227 h    | 9.420 g | 0.667 f      | 0.773d  |
| T8         | 45.33 h           | 45.00 f  | 32.40 g              | 30.28fg | 9.707 h                              | 9.886 g | 8.926 i    | 9.006 h | 0.780 c      | 0.880b  |
| T9         | 49.00 ef          | 48.67cd  | 34.12 f              | 32.67 e | 13.247c                              | 11.380e | 12.760c    | 10.773e | 0.487 h      | 0.607g  |
| T10        | 51.67abc          | 45.67 ef | 32.28 g              | 30.65 f | 12.767d                              | 11.200e | 12.360d    | 10.700e | 0.406 i      | 0.500h  |
| T11        | 53.67 a           | 52.00 a  | 42.40 a              | 39.52 a | 14.127a                              | 13.740a | 13.813a    | 13.373a | 0.313 k      | 0.367 j |
| T12        | 52.67ab           | 51.00ab  | 40.08 b              | 37.10bc | 13.573b                              | 13.180b | 13.207b    | 12.773b | 0.367 j      | 0.407 i |

Means followed by the same letter (s) within each column do not significantly differed using Duncan's Multiple Range Test at the level of 5%. T1: AS<sub>20.5%</sub> N; T2: AN<sub>33.5%</sub> N; T3: Urea<sub>46.0%</sub> N; T4: AS + GTU; T5: AN + GTU; T6: Urea + GTU; T7: AS<sub>50%</sub> + AN<sub>50%</sub> + GTU; T8: AS<sub>50%</sub> + Urea<sub>50%</sub> + GTU; T9: Nitroform<sub>38%</sub> N; T10: Compost<sub>1.2%</sub> N; T11: Compost<sub>50%</sub>+AS<sub>50%</sub>+GTU, and T12: Compost<sub>50%</sub>+AN<sub>50%</sub>+GTU. AS: Ammonium sulphate; AN: Ammonium nitrate; GTU: Guanylthiourea (nitrification inhibitors).

**Table 3: Tuber quality characters of potato as affected by nitrogen sources and nitrification inhibitors in 2007/08 and 2008/09 seasons.**

| Treatments | Tuber dry matter (%) |          | Specific gravity of tuber |         | Starch (%) |         | Nitrate accumulation (ppm) |         | Nitrite accumulation (ppm) |         |
|------------|----------------------|----------|---------------------------|---------|------------|---------|----------------------------|---------|----------------------------|---------|
|            | 2007/08              | 2008/09  | 2007/08                   | 2008/09 | 2007/08    | 2008/09 | 2007/08                    | 2008/09 | 2007/08                    | 2008/09 |
| T1         | 21.41efg             | 21.48bcd | 1.082de                   | 1.085ef | 14.16cd    | 14.38cd | 49.28 f                    | 48.18fg | 0.40 h                     | 0.42 e  |
| T2         | 21.14 fg             | 22.08abc | 1.081de                   | 1.084fg | 14.00de    | 14.21de | 67.32 a                    | 65.38 a | 0.64 a                     | 0.58 a  |
| T3         | 20.965 g             | 20.82cde | 1.079ef                   | 1.083 g | 13.66ef    | 13.80ef | 63.00bc                    | 61.22bc | 0.58 c                     | 0.53 b  |
| T4         | 22.08 cd             | 21.47bcd | 1.087 b                   | 1.092 c | 14.35cd    | 14.10de | 51.74 f                    | 50.00ef | 0.46 g                     | 0.38 f  |
| T5         | 21.99cde             | 21.08b-e | 1.085bc                   | 1.089 d | 14.26cd    | 14.00de | 65.40ab                    | 62.23ab | 0.62 b                     | 0.57 a  |
| T6         | 20.88 g              | 20.53 de | 1.078 f                   | 1.081 h | 13.40 f    | 13.43 f | 62.12 c                    | 58.80 c | 0.55 d                     | 0.51 bc |
| T7         | 22.32 bc             | 21.82abc | 1.087 b                   | 1.0792i | 14.52bc    | 14.40cd | 58.72 d                    | 55.34 d | 0.51 e                     | 0.48 cd |
| T8         | 20.14 h              | 19.89 e  | 1.077 f                   | 1.0782i | 12.94 g    | 12.80 g | 55.38 e                    | 52.30de | 0.48 f                     | 0.45 de |
| T9         | 21.62def             | 21.64a-d | 1.083cd                   | 1.086e  | 12.90 g    | 15.38 b | 44.17 g                    | 45.13gh | 0.38 j                     | 0.35 fg |
| T10        | 22.22 c              | 22.10abc | 1.088 b                   | 1.096 b | 14.82 b    | 14.80 c | 38.33 h                    | 38.71 i | 0.33 j                     | 0.28 h  |
| T11        | 22.93 a              | 22.84 a  | 1.097 a                   | 1.098 a | 15.78 a    | 15.91 a | 36.18 h                    | 35.82 i | 0.28 k                     | 0.25 h  |
| T12        | 55.83 ab             | 22.32 ab | 1.095 a                   | 1.095 b | 15.40 a    | 15.73ab | 41.70 g                    | 44.23 h | 0.37 i                     | 0.32 g  |

Means followed by the same letter (s) within each column do not significantly differed using Duncan's Multiple Range Test at the level of 5%. T1: AS<sub>20.5%</sub> N; T2: AN<sub>33.5%</sub> N; T3: Urea<sub>46.0%</sub> N; T4: AS + GTU; T5: AN + GTU; T6: Urea + GTU; T7: AS<sub>50%</sub> + AN<sub>50%</sub> + GTU; T8: AS<sub>50%</sub> + Urea<sub>50%</sub> + GTU; T9: Nitroform<sub>38%</sub> N; T10: Compost<sub>1.2%</sub> N; T11: Compost<sub>50%</sub>+AS<sub>50%</sub>+GTU, and T12: Compost<sub>50%</sub>+AN<sub>50%</sub>+GTU. AS: Ammonium sulphate; AN: Ammonium nitrate; GTU: Guanylthiourea (nitrification inhibitors).

### 3. Chemical composition and nitrogen efficiency ratio:

Data presented in Table 4 show that, the differences in means of N, P and K-uptake as well as nitrogen efficiency ratio due to various application sources and/or nitrification inhibitor were differed significantly, in both season of study. The highest values of these traits were obtained from potato plants receiving Compost 50% + AS 50%

+ GTU, while the lowest values were recorded with sole soluble form of nitrogen (AS or AN). The positive effect of GTU on N, P and K-uptake may be due to the efficiency of nitrification inhibitor in keeping nitrogen for longer time in the form of  $\text{NH}_4^+$  which helps in modification of nutrient uptake by plant (Laskshmanan and Prasad, 2004; Di and Cameron, 2004). Moreover, Tisdale *et al.* (1985) reported that the addition of nitrogen in combination with adequate phosphorus tended to

increase K-uptake by plants. They added also that, potassium concentration may be as high in the  $\text{NH}_4^+$ -nourished plants as it absorbed by soil colloids, so, it does not leach out of soil and still reliable for plants, generally such case may give the plant more chance for absorbing N, and consequently, the other nutrients for building dry

matter. Shoji *et al.* (2001) discussed that contributions of controlled-release fertilizer and nitrification inhibitor to conserve air and water quality are basically due to maximizing NUE and reducing the N fertilization rate.

**Table 4: Chemical composition of potato tuber and nitrogen efficiency ratio as affected by nitrogen sources and nitrification inhibitors in 2007/08 and 2008/09 seasons.**

| Treatments | N-uptake (mg/plant tuber) |            | P-uptake (mg/plant tuber) |           | K-uptake (mg/plant tuber) |           | Nitrogen efficiency ratio (NER) |         |
|------------|---------------------------|------------|---------------------------|-----------|---------------------------|-----------|---------------------------------|---------|
|            | 2007/08                   | 2008/09    | 2007/08                   | 2008/09   | 2007/08                   | 2008/09   | 2007/08                         | 2008/09 |
| T1         | 4312.48 ef                | 4122.88 e  | 375.12 e                  | 485.12 e  | 4486.02 ef                | 4183.16gh | 63.18                           | 66.22   |
| T2         | 4354.22 ef                | 3518.78 f  | 370.01 e                  | 452.23 f  | 4452.91ef                 | 4172.52 h | 62.15                           | 65.37   |
| T3         | 3437.70 h                 | 2701.42 h  | 312.65 e                  | 372.65 h  | 3584.39 h                 | 3520.01 k | 60.44                           | 60.37   |
| T4         | 4826.16 cd                | 4523.27 d  | 494.80 d                  | 504.20 e  | 4943.45 d                 | 4612.04 f | 73.71                           | 71.52   |
| T5         | 4587.23 de                | 4307.34 e  | 478.96 d                  | 490.96 e  | 4693.22 e                 | 4311.44 g | 69.85                           | 67.96   |
| T6         | 3961.12 fg                | 3342.56 fg | 359.47 e                  | 420.73 g  | 4271.78 f                 | 3927.70 i | 57.07                           | 59.18   |
| T7         | 4924.64 cd                | 4892.04 c  | 615.42 b                  | 602.23 c  | 5400.31 c                 | 5050.21 d | 54.96                           | 56.63   |
| T8         | 3629.20gh                 | 3172.36 g  | 336.99 e                  | 398.10 gh | 3875.12 g                 | 3729.10 j | 53.93                           | 54.92   |
| T9         | 4875.82 cd                | 4712.21cd  | 540.49 cd                 | 540.28 d  | 5137.93 d                 | 4823.28 e | 73.59                           | 63.22   |
| T10        | 5176.46 bc                | 4900.32 c  | 597.32 bc                 | 680.04 a  | 5530.75 c                 | 5337.11 c | 70.92                           | 62.22   |
| T11        | 5765.08 a                 | 5369.28 a  | 710.37 a                  | 642.16 b  | 6270.50 a                 | 5922.34 a | 78.48                           | 76.33   |
| T12        | 5432.15 ab                | 5115.50 b  | 650.54 ab                 | 580.47 c  | 5842.04 b                 | 5729.20 b | 75.40                           | 73.22   |

Means followed by the same letter (s) within each column do not significantly differed using Duncan's Multiple Range Test at the level of 5%. T1: AS<sub>20.5%</sub> N; T2: AN<sub>33.5%</sub> N; T3: Urea<sub>46.0%</sub> N; T4: AS + GTU; T5: AN + GTU; T6: Urea + GTU; T7: AS<sub>50%</sub> + AN<sub>50%</sub> + GTU; T8: AS<sub>50%</sub> + Urea<sub>50%</sub> + GTU; T9: Nitroform<sub>38%</sub> N; T10: Compost<sub>1.2%</sub> N; T11: Compost<sub>50%</sub>+AS<sub>50%</sub>+GTU, and T12: Compost<sub>50%</sub>+AN<sub>50%</sub>+GTU. AS: Ammonium sulphate; AN: Ammonium nitrate; GTU: Guanylthiourea (nitrification inhibitors).

#### 4. Residual $\text{NH}_4$ and $\text{NO}_3$ in soil:

Concerning the residual ammonium and nitrate nitrogen in soil after plants harvesting. Data in Figures 1&2 indicate the highest residual available of  $\text{NH}_4^+$  -N was obtained in the treatment of compost and nitroform, while, a soluble form of AN or U gave the highest residual  $\text{NO}_3^-$  -N compared with other treatments, in both seasons of study.  $\text{NO}_3^-$ -N leaching loss decreased in the leachates with Compost, Compost<sub>50%</sub> + AS<sub>50%</sub> or AN<sub>50%</sub> + GTU compared to soluble form of

nitrogen. In the case of compost combined with GTU or nitroform treatment increase yield of  $\text{NH}_4^+$  -N and a reduction in  $\text{NO}_3^-$  -N compared with the amount of  $\text{NH}_4^+$  -N formed from other treatments. This result may be attributed to the effect of GTU or Compost or coated fertilizer on delaying the release of nitrogen as indicated by Vallejo *et al.* (2006) and Khalil *et al.* (2009). The application of GTU as a nitrification inhibitor regulate the release of  $\text{NH}_4^+$  -N out of compost treatments and it can also retard the nitrification process which produce  $\text{NH}_3^-$  -N in that easily leachable (Dahadouh, *et al.* 2004).






Fig. 1: Residual NH<sub>4</sub>-N in soil at harvesting as affected by nitrogen sources and nitrification inhibitors in 2007/08 and 2008/09 seasons.




Fig. 2: Residual NO<sub>3</sub>-N in soil at harvesting as affected by nitrogen sources and nitrification inhibitors in 2007/08 and 2008/09 seasons.

## REFERENCES

- [1] Amberger, A. 1989. Research on dicyandiamide as a nitrification inhibitor and future outlook. *Comm. Soil Sci. Plant Anal.*, 20 (19&20): 1933 -1955.
- [2] Amberger, A. and M. P. Germann-Bauer. 1990. Effect of the nitrification inhibitors 1-amidino-2-thiourea and dicyandiamide in combination with urea and ammonium sulfate. *Nutrient Cycling in Agroecosystems*, 21 (3): 179-183.
- [3] AOAC (Association of Official Analytical Chemists) .1990. *Official Methods of Analysis*. 15<sup>th</sup> Ed., Washington, DC, USA.
- [4] Aujla, M. S.; H. S. Thind, and G. S. Buttar. 2007. Fruit yield and water use efficiency of eggplant (*Solanum melongena* L.) as influenced by different quantities of nitrogen and water applied through drip and furrow irrigation. *Scientia Horticulturae*, 112: 142-148.
- [5] Bakr, A. A. and R. A. Gawish. 1997. Trials to reduce nitrate and oxalate content in some leafy vegetable. II. Interactive effects of the manipulating of the soil nutrient supply, different blanching media and preservation methods followed by cooking process. *J. Sci. Food Agric.*, 73 (2): 169-178.
- [6] Black, C. A. 1965. *Methods of Soil Analysis*. Part 1 and 2. Amer. Soc. Agron. Inc., Madison, Wisconsin, USA.
- [7] Cottenie, A.; M. Verloo; L. Kiekens, and G. Velghe. 1982. *Biological and Analytical Aspects of Soil Pollution Hand Book*. Gent, Belgium.
- [8] Dachler, M. 1993. The effect of dicyandiamide-containing nitrogen fertilizers on root crops. 2. Communication: The effect on grain-malze and potatoes. *Die Bodenkultur - Journal for Land Management, Food and Environment*, 44 (2): 541-546.

- [9] Dahadouh, F. S. M.; A. A. Osman, and M. O. El-Azab. 2004. Nitrogen loss from sandy soils treated with manure and N fertilizers with and without nitrification inhibitor. *Egypt J. Soil Sci.*, 44 (4): 531-546.
- [10] Di, H. J. and K. C. Cameron. 2004. Effects of the nitrification inhibitor dicyandiamide on potassium, magnesium and calcium leaching in grazed grassland. *Soil Use and Management*, 20 (1): 2-7.
- [11] Khalil, M. I.; R. Gutser, and U. Schmidhalter. 2009. Effects of urease and nitrification inhibitors added to urea on nitrous oxide emissions from a loess soil. *J. Plant Nutr. Soil Sci.*, 172 (5): 651-660.
- [12] Laskshmanan, R. and R. Prasad .2004. Influence of moisture regime on inhibition of nitrification by DCD. *Archives of Agronomy and Soil Science*, 50 (4-5): 495-500.
- [13] Martin, H. W.; D. A. Graetz; S. J. Locascio, and D. R. Hensel. 1993. Nitrification inhibitor influences on potato. *Agron. J.*, 85 (3): 651-655.
- [14] Page, A. L. 1982. *Methods of Soil Analysis*. 2<sup>nd</sup> Ed., Part 1, Soil Sci. Soc. Amer., Madison, Wisc., USA.
- [15] Pasda, G.; R. Hähndel, and W. Zerulla. 2002. The new nitrification inhibitor DMPP (ENTEC®): Effects on yield and quality of agricultural and horticultural crops. *Dev. Plant Soil Sci.*, 92 (10): 758-759.
- [16] Tisdale, S. L.; W. L. Nelson, and J. D. Beaton. 1985. *Soil Fertility and Fertilizers*. 4<sup>th</sup> Ed. MacMillan Publ., New York.
- [17] Trenkel, M. E. 1997. *Improving Fertilizer Use Efficiency: Controlled-Release and Stabilized Fertilizers in Agriculture*. International Fertilizer Industry Association, Paris, 156 pp.
- [18] Shoji, S.; J. Delgado; A. Mosier, and Y. Miura. 2001. Use of controlled release fertilizers and nitrification inhibitors to increase nitrogen use efficiency and to conserve air and water quality. *Comm. Soil Sci. Plant Anal.*, 32 (7&8): 1051-1070.
- [19] Snedecor, G. W. and W. G. Cochran. 1982. *Statistical Methods*. 7<sup>th</sup> Ed. 2<sup>nd</sup> Printing, Iowa State. Univ. Press, Ame., USA, pp 507.
- [20] Steel, R. G. D. and Torrie, J. H. 1980. *Principles and Procedures of Statistics. A Biometrical Approach*. 2<sup>nd</sup> Ed. McGraw-Hill Publishing Co., New York, USA.
- [21] Vallejo A.; U. M. Skiba; L. García-Torres; A. Arce; S. López-Fernández, and L. Sánchez-Martín. 2006. Nitrogen oxides emission from soils bearing a potato crop as influenced by fertilization with treated pig slurries and composts. *Soil Biol. Bioch.*, 38 (9): 2782-2793.
- [22] Watanabe, T. 2006. Influence of 2-chloro-6 (trichloromethyl) pyridine and dicyandiamide on nitrous oxide emission under different soil conditions. *Soil Sci. Plant Nutr.*, 52 (2): 226-232.

**Correspondence to:**

Dr. Abd El-Badea S. Ezzat  
Horticulture Research Institute,  
9 Gamaa Street,  
Giza 12619, Giza,  
EGYPT  
Cell Phone: 002 012 1078703  
Office Phone: 002 050 226 4976  
Fax: 00 2 02 3572 1628  
E-mail [as\\_ezzataf@yahoo.com](mailto:as_ezzataf@yahoo.com)

Dr. Hamdino M.I. Ahmed  
Horticulture Research Institute,  
9 Gamaa Street,  
Giza 12619, Giza,  
EGYPT  
Cell Phone: 002 018 1483434 & 002 010 6215073  
Office Phone: 002 050 226 4976  
Fax: 00 2 02 3572 1628  
E-mail [hamdino@yahoo.com](mailto:hamdino@yahoo.com);  
[hamdino.ahmed@gmail.com](mailto:hamdino.ahmed@gmail.com)

Date of submission 05/03/2011

تحسين كفاءة النيتروجين المستفادة بواسطة نباتات البطاطس  
أ. تأثير مثبتات النترتة مع مصادر مختلفة للنيتروجين على تقليل فقد النيتروجين وتحسين الإنتاجية والتركيبة الكيماوي

عبد البديع صالح عزت<sup>1</sup>، عادل محمد عبد الحميد<sup>2</sup>، حمدينو محمد إبراهيم أحمد<sup>1</sup> و أمل أبو الفتوح العوضى<sup>1</sup>  
١ قسم بحوث الخضار- معهد بحوث البساتين- مركز البحوث الزراعية- وزارة الزراعة- مصر  
٢ قسم بحوث تغذية النبات- معهد بحوث الأراضي والمياه والبيئة- مركز البحوث الزراعية- وزارة الزراعة- مصر

تحدث أكسدة بيولوجية للأمونيا  $NH_4^+$  (المضافة للتربة من خلال التسميد المعدني أو نتيجة تحلل المادة العضوية) إلى نترات من خلال عملية النترية nitrification، والتي تقوم بها بكتيريا متخصصة، وقد يكون من المفضل استخدام أي تقنية يؤدي إلى تأخير عملية النترية، وبالتالي تقليل فقد النتروجين في صورة نترات من خلال عملية الرش.

يهدف هذا البحث إلى رفع كفاءة استخدام الأسمدة النيتروجينية عن طريق استخدام مثبتات النترية وتأثير ذلك على تقليل فقد النتروجين وزيادة الإنتاجية والتركيب الكيماوي في البطاطس صنف كارا. كما يهدف إلى تقليل الأسمدة الأزوتية المضافة وتقليل التلوث البيئي والمحافظة على صحة الإنسان.

ولتحقيق هذه الأغراض أجريت تجربتان حقليةتان في المزرعة البحثية بالبرامون- المنصورة- محافظة الدقهلية خلال العروة الشتوية لموسمي ٢٠٠٧/٢٠٠٨ و ٢٠٠٨/٢٠٠٩ لدراسة تأثير الصور الدائبة من الأسمدة الأزوتية (نترات نشادر- سلفات نشادر- يوريا)، مع/أو الصور بطيئة الأمداد (الكمبوست- نتروفورم)، مع/أو بدون إضافة مثبت النترية جوانيل ثيوربا (guanylthiourea, GTU) وتأثير ذلك على تحقيق الأهداف السابقة. استخدم تصميم القطاعات الكاملة العشوائية في ثلاث مكررات. وكانت أهم النتائج المتحصل عليها ما يلي:

- بصفة عامة. أدى إضافة مثبت النترية GTU مع الكمبوست (٥٠% من المعدل الموصى به) و سلفات نشادر (٥٠% من المعدل الموصى به) إلى حدوث زيادة معنوية في كل الصفات محل الدراسة، ما عدا تركيز كل من النترات والنيتريت في الدرنات.
- أدى إضافة الكمبوست (٥٠%) و سلفات نشادر (٥٠%) مع GTU إلى حدوث زيادة معنوية في صفات النمو الخضري (متمثلة في طول النبات والوزن الجاف) و صفات المحصول (المحصول الكلي والمحصول القابل للتسويق) والجودة (المادة الجافة في الدرنات والكثافة النوعية والنشا) و التركيب الكيماوي لدرنات البطاطس (المحتوي الغذائي من ن، ف، بو)، بينما أدت هذه المعاملة إلى حدوث نقص معنوي في صفات المحصول الغير قابل للتسويق، وتركيز النترات والنيتريت في الدرنات، وذلك في موسمي الدراسة.
- أعطت معاملة الكمبوست وسماد النتروفورم أعلى القيم بالنسبة للنيتروجين المتبقي في التربة في صورة  $NH_4^+$ ، بينما سجلت معاملة سماد نترات النشادر أعلى القيم بالنسبة للنيتروجين المتبقي في التربة في صورة  $NO_3^-$  وذلك مقارنة بباقي المعاملات خلال موسم الدراسة.

وبناء عليه توصي هذه الدراسة باستخدام سماد الكمبوست بمعدل ٩ طن/فدان مع سلفات النشادر بمعدل ٩٠ كجم (٢/١ المعدل الموصى به) مع إضافة مثبت التآزت GTU إلى حقول البطاطس للحصول على أفضل النتائج بالنسبة للمحصول وتحقيق أقصى استفادة من النتروجين المضاف للتربة، مع خفض التلوث البيئي.