

## Effect of GA<sub>3</sub> and Potassium Nitrate in Different Dates on Fruit Set, Yield and Splitting of Washington Navel Orange

\*Abd El-Rahman, G.F.; Hoda, M.Mohamed and Ensherah, A.H.Tayh

Citrus Department, Hort. Res. Instit., Agric. Res. Center, Giza, Egypt  
\*gamalelashmanty@yahoo.com

**Abstract:** Two concentrations of GA<sub>3</sub> were used alone or with 0.5% urea i.e. (50 and 100 ppm) and three levels of potassium nitrate (KNO<sub>3</sub>, 2, 4 and 6% were tested as foliar sprays at different times i.e. first time application at full bloom stage, the second time application was at fruit diameter from 1.5-2.0 cm and (first and second time of application) for investigation their effects on fruit set, yield, fruit quality and splitting of 40-years-old Washington navel orange (*Citrus Sinensis*), trees budded on Sour Orange (*Citrus aurantium*, L.Obseck) rootstock during 2008 and 2009 seasons. Data indicated that, all treatments increased fruit set, yield and fruit quality and decreased fruit splitting as compared with control treatment. Data also revealed that, foliar sprays of trees by (GA<sub>3</sub> at 50 ppm) with or without 0.5% urea were superior for inducing the highest increase of fruit set and yield, in addition KNO<sub>3</sub> at 4% comparing with 2% and 6%. Also, KNO<sub>3</sub> treatments increased fruit size, peel thickness and juice acidity especially with high concentrations; on the other hand, the use of GA<sub>3</sub> at 50 or 100 ppm alone or with 0.5% urea increased fruit T.S.S and reduced nitrite and nitrate contents in fruit juice as compared with KNO<sub>3</sub> treatments. Moreover, it is noticed that trees sprayed at first application or first and second time of application gave the best results, while second application was the best for reducing fruit splitting. It could be recommended the best treatment for increasing yield and gave high fruit quality is GA<sub>3</sub> at 50 ppm with or without 0.5% urea at full bloom stage especially in respect with reducing nitrite and nitrate in fruit juice, and use KNO<sub>3</sub> 4% at the second time of application to reduce fruit splitting.

[Abd El-Rahman, G.F.; Hoda, M.Mohamed and Ensherah, A.H.Tayh **Effect of GA<sub>3</sub> and Potassium Nitrate in Different Dates on Fruit Set, Yield and Splitting of Washington Navel Orange**] Nature and Science 2011; 9(12): 196-204]. (ISSN: 1545-0740). <http://www.sciencepub.net> 27

**Key word:** GA<sub>3</sub>, Navel Orange, KNO<sub>3</sub>, Fruit Set, Yield

### 1. Introduction

Navel orange is a popular fresh fruit due to its seedless, large size, characteristic flavor and aroma (Wardowski, *et al.*, 1985). Also, navels orange are in important source of early season income for citrus growers in some commercial citrus areas of the world. Yields are erratic and usually low in many areas due to lack functional pollen, rarely produce viable ovules and in addition, are weakly parthenocarpic (Kretdorn, 1965). Flower and fruit drop of navel orange occurred in three phases and amount to a total 91%, giving a fruit set of 9% (Villafane, *et al.*, 1989). Gibberellic acid (GA<sub>3</sub>) and potassium nitrate have a board range of uses in citriculture; Gibberellins have been used in citrus production with several objectives including bloom reduction, increased fruit setting, improvement of fruit quality and improved maturation control (Agusti and Almela, 1991). The application of gibberellic acid soon after flowering at doses between 10 and 15 ppm can result in delayed abscission and increased fruit set, mainly in Clementine tangerines (Fornes *et al.*, 1992; El-Otmani, 1992). However, the increase in fixing and productivity does not happen frequently and depends on factors including variety, plant status and time of application (Davies, 1987; Talón *et al.*,

1997). Also, GA<sub>3</sub> increased fruit set in navels orange (Smith, 1993; Babu and Lavania, 1985), in tangerine hybrids (Brosh and Monselise, 1977) and increased the yield either as number or weight of fruits per tree of Washington navel orange (Ibrahim, *et al.*, 1994). Positive results have been obtained with potassium nitrate supplementation even with good plant nutritional status, which is possibly related to the lower ability to mobilize mineral reserves necessary for the period (Ruiz *et al.*, 2001). On the other hand, potassium decreases the loss of fruit from splitting (Lavon, *et al.*, 1992); also, GA<sub>3</sub> reduces fruit splitting when applied shortly after the end of the June drop in "Nova" hybrid mandarin (Garcia, *et al.* 1994). GA<sub>3</sub> application improved fruit quality (Davies, *et al.*, 1999 and 2001), while KNO<sub>3</sub> increased fruit size and juice acidity for Shamuti and Valencia oranges (Erner *et al.* 1993).

The purpose of this work is to study the effect of Gibberellic acid (GA<sub>3</sub>) and potassium nitrate (KNO<sub>3</sub>) sprays at different times and concentrations on fruit set, yield, fruit splitting and fruit quality on Washington navel orange trees. Hence, to identify the best treatments for achieve the highest return for the growers.

## 2. Material and Methods

The experiment was carried out in the private orchard belonged for Mr. Hassan Marie in Benha, El-Kalubiah Governorate, Egypt, in 2008 and 2009 seasons on forty-years-old of "Washington navel" (*Citrus Sinensis*) orange trees grafted on Sour orange rootstock (*Citrus aurantium*, L.Obseck) and were planted at 5×5 m spacing in clay soil. Ninety-six "Washington navel" orange trees were divided into twenty four groups according to vigor and number of flowers and used for data collection. The design was a randomized block with 4 single-tree replicates and 8 treatments with 3 time intervals of spray all treatments. The experiment involved the following eight treatments. GA<sub>3</sub> at 50 ppm and 100 ppm plus the same concentrations of GA<sub>3</sub> with 0.5% urea; potassium nitrate treatments was used at three levels 2, 4 and 6%; the control trees untreated. Trees were sprayed on full bloom stage (first application), fruit diameter from 1.5-2.0 cm (second application) and both (first and second time of application). The total number of flowers was counted before treatments and commercially available urea containing 46% N and Berlex (containing 92% GA<sub>3</sub> and 8% of other gibberellins) were used in the trials. Triton B at 0.1% was used as a wetting agent for all spraying solutions. In addition foliar spray of potassium treated as KNO<sub>3</sub> 98%. All trees generally received adequate organic and inorganic fertilization. Irrigation was given at intervals of about 10-15 days in summer and 15-21 days in winter. However, a balanced foliar fertilization of all microelements was adopted three times yearly (February, May and August). The following parameters of the studied treatments were carried out.

### Percentage of fruit set.

Fruitlets and fruits drop were expressed as an accumulative percentage per tree. Percentage of fruit set = number of fruit set during a given interval × 100/ number of fruit at the beginning of the interval.

### Yield

Weight (kg) and number of mature fruits per tree were recorded at late November of each season.

### Fruit splitting percentage

Determination dates of beginning primary symptoms of splitting for each treatment were observed and recorded. Total number of fruits was counted for each tree at the end of July to end of October 2008 and 2009. Percentage of fruit splitting was calculated using the following equation. Fruit splitting percentage = number of splitting fruits × 100/ Total number of fruits.

### Nitrite and nitrate contents in fruit juice

A sample of 10 ml of fruit juice was taken from each replicate to determine nitrite and nitrate according to the methods outlined by (Sen and Donaldson, 1978).

### Physical properties of fruits

Fruit weight, fruit size, fruit length and diameter and peel thickness for each fruit were measured by using the varinier caliper. Juice volume, also Juice weight percentage was calculated by using the following equation. Juice weight % = average juice weight (gm.) / fruit × 100 / average fruit weight (gm.).

### Chemical properties of fruits

Total soluble solids (T.S.S %) was determined by using Zeiss hand refractometer, total acidity was determined according (A.O.A.C., 1995), also T.S.S/Acid was calculated.

### Statistical analysis

The experiment was designed in completely randomized block design with four replicates for each treatment and each replicate was represented by one tree. The obtained data of both seasons were subjected to analysis of variance according to (Clark and Kempson, 1977) and the means were differentiated using Duncan multiple range test at 5% level (Duncan, 1955).

## 3. Results and Discussion

### Fruit set percentage

It is obvious from Tables (1&2) that the difference between the use of gibberellic acid and potassium nitrate on fruit set of Washington navel orange were significant. The results revealed that, trees spraying with both GA<sub>3</sub> and KNO<sub>3</sub> enhanced fruit set, whereas trees treated by GA<sub>3</sub> 50ppm + 0.5% urea gave significantly the highest fruit set percentage at (30, 60,165 and 205 days) after full blooming followed by trees sprayed with GA<sub>3</sub> 50ppm alone, while the lowest values of fruit set percentage were obtained by trees treated by KNO<sub>3</sub> 2% and control treatment meanwhile, the other treatments gave the intermediate values in this regard during the two seasons. On the other hand, concerning the time intervals of spray for all treatments, data revealed that, trees sprayed at ( first application+ second application) had the highest significant values for fruit set followed by (first application) while (second application) gave the lowest significant values during two seasons. Several scientific trials show that the gibberellins increase the cell wall flexibility by stimulating the synthesis of new cellulose polymers (Richards et al., 2001). The present results are a

general in harmony with (Davies, 1987) who stated that, gibberellins could reduce the senescence of leaves and fruits. Also; (Brosh, *et al.*, 1975 and Tominaga, 1998) reported that, GA<sub>3</sub> and KNO<sub>3</sub> decreased abscission of fruitlets in navel oranges and grapefruit. In addition, potassium nitrate (KNO<sub>3</sub>) enhanced fruit set and increased yield of orange and mandarin trees (El-Deeb, 1989; El-Fangary, 1998 and Mostafa, *et al.*, 2005).

#### **Fruit weight (gm)**

Results in Table (3) showed that no constant trend during two studied seasons for fruit weight due to different treatments was noticed on Washington navel orange. Whereas, foliar sprays of trees by KNO<sub>3</sub> 2% scored the highest significant values for fruit weight at the (first and second time of application) in the first season, while trees sprayed with GA<sub>3</sub> 50ppm gave the maximum values at full bloom stage in the second season. On the other hand, trees treated by GA<sub>3</sub> 100 ppm gave the lowest significant values for fruit weight at (second time of application) and (first and second time of application) in the first season. Meanwhile, the lowest values were on KNO<sub>3</sub> 4% treatment at full bloom stage in the second season. Anyhow, the differences between all treatments were high to be significant.

#### **Fruit number/tree**

Data in the Table (3) revealed that, all treatments increased fruit number with some fluctuations as compared with control treatment. On the other hand, it was noticed that the maximum fruit number were obtained during foliar sprays of trees at full bloom stage (first application) and (first and second time of application) while (second application) gave the lowest significant values of fruit number in the first and second season.

#### **Yield**

Yield as weight (Kg/tree or ton/fed.) are shown in Table (4) yield was significantly increased by all treatments either with spraying gibberellic acid or with KNO<sub>3</sub> at any concentrations as compare to the control treatment. However, yield was gradually increased by spraying of KNO<sub>3</sub> 2% than the control then the range of increase was higher when trees treated by KNO<sub>3</sub> at 6% then 4 % concentrations and reached the maximum with gibberellic acid (GA<sub>3</sub>) treatments especially (GA<sub>3</sub> 50ppm + 0.5%urea) which gave the highest yield (yield Kg/tree and ton/fed.) for the first and second seasons. Also, data revealed that trees sprayed at (first application) and (first and second time of application) scored the higher averages as compared with those sprayed at (second application). From the above results, it is

interest to notice that, GA<sub>3</sub> and KNO<sub>3</sub> at different rates spraying increased the yield of Washington navel orange trees. These results are in the same line with those obtained by (Smith, 1993; Babu and Lavania, 1985) in navel orange and (Brosh and Monselise, 1977) in tangerine hybrids; they found that gibberellic acid (GA<sub>3</sub>) increased fruit set. Also, (Ibrahim, *et al.* 1994) reported that, trees spraying with GA<sub>3</sub> increased the yield of Washington navel orange. Likewise, (El-Fangary, 1998 and Mostafa, *et al.*, 2005) found that, KNO<sub>3</sub> enhanced fruit set and increased yield of orange and mandarin trees.

#### **Fruit splitting percentage**

It is evident from the results shown in Fig. (1&2) that gibberellic acid and potassium nitrate treatments decreased fruit splitting percentage as compared with control treatment especially trees treated by KNO<sub>3</sub> 6% gave the lowest significant values of fruit splitting due to the peel thickness of fruits which increased with increasing the concentration of potassium nitrate Table (8). This finding is in parallel with the finding of (Josan *et al.*, 1995) in Lemon who reported that the thicker peel helps to resist fruit puncture. While the other treatments gave the intermediate values during two seasons (2008 &2009). Also, it can be noticed that, the splitting tendency is greatly depended on the stage of development of the fruit whereas, full bloom stage (first application) scored the maximum significant average for fruit splitting while, trees treated at (second application) or (first and second time of application) had the minimum significant averages of fruit splitting, although GA<sub>3</sub> treatments scored decreasing of fruit peel thickness, these results may be due to the elastic walls of fruit cells as a result of spraying with GA<sub>3</sub> in these stages. This explanation can be agreed with (Richards *et al.*, 2001) they found that, the gibberellins increase the cell wall flexibility by stimulating the synthesis of new cellulose polymers.

#### **Nitrite and nitrate content in fruit juice**

As shown in Tables (5& 6) results revealed that, nitrite and nitrate contents in fruit juice were affected by different treatments in the two seasons. It is clear that, trees sprayed with (KNO<sub>3</sub>) gave the highest values for nitrite and nitrate in their fruit juice especially with high concentrations, while, trees sprayed with gibberellic acid (GA<sub>3</sub>) either 50 ppm alone or plus 0.5 % urea gave the lowest values in fruit juice than the other treatments. This means that, the beneficial effect of spraying GA<sub>3</sub> either 50 ppm or plus 0.5% urea was mainly on reducing nitrite and nitrate in fruit juice. Also, it is noticed from Table (5&6) that the time of applications did not show any

distinctive effect on nitrite and nitrate contents in fruit juice in the first and second seasons. In this respect, (Ibraheem, 1994 and Abd El-Migeed, *et al.*,

2007) mentioned that, mineral nitrogen fertilization easily forms nitrate in fruit juice as compare with using organic fertilizers.

**Table (1) Fruit set percentage (after 30 & 60days) of Washington navel orange trees as affected by gibberellic acid and potassium nitrate during 2008 and2009 seasons**

| Experiment (1)        | Fruit set % (after 30days) |                     |              |              | Fruit set % (after 60days) |                     |              |              |
|-----------------------|----------------------------|---------------------|--------------|--------------|----------------------------|---------------------|--------------|--------------|
|                       | 2008, Season               |                     | 2009, Season |              | 2008, Season               |                     | 2009, Season |              |
|                       | Fruit cluster (Fruit/Tree) | Fruit diameter (mm) | (Fruit/Tree) | (Fruit/Tree) | Fruit cluster (Fruit/Tree) | Fruit diameter (mm) | (Fruit/Tree) | (Fruit/Tree) |
| Control               | 1.000                      | 1.000               | 1.000        | 1.000        | 1.000                      | 1.000               | 1.000        | 1.000        |
| G.A., 50ppm           | 14.56a                     | 44.47ab             | 11.25a       | 32.88a       | 75.1cd                     | 49.15cd             | 75.1cd       | 44.51d       |
| G.A., 100ppm          | 49.86cd                    | 59.26cd             | 71.86cd      | 72.97cd      | 62.57cd                    | 74.96cd             | 68.52cd      | 74.96cd      |
| G.A., 150ppm          | 75.56de                    | 61.92de             | 75.46de      | 72.13de      | 81.26e                     | 75.44e              | 81.13e       | 77.92de      |
| G.A., 200ppm          | 77.86de                    | 67.26de             | 72.76de      | 71.23de      | 84.26e                     | 80.64e              | 81.73e       | 77.92de      |
| G.A., 250ppm          | 77.86de                    | 67.26de             | 72.76de      | 71.23de      | 84.26e                     | 80.64e              | 81.73e       | 77.92de      |
| KNO <sub>3</sub> , 2% | 44.25a                     | 47.76a              | 48.47a       | 49.92a       | 69.25b                     | 71.75ab             | 72.76ab      | 74.96cd      |
| KNO <sub>3</sub> , 4% | 44.25a                     | 47.76a              | 48.47a       | 49.92a       | 69.25b                     | 71.75ab             | 72.76ab      | 74.96cd      |
| KNO <sub>3</sub> , 6% | 44.25a                     | 47.76a              | 48.47a       | 49.92a       | 69.25b                     | 71.75ab             | 72.76ab      | 74.96cd      |
| Mean (Error)          | 44.25a                     | 47.76a              | 48.47a       | 49.92a       | 69.25b                     | 71.75ab             | 72.76ab      | 74.96cd      |

Mean separation within columns by Duncan’s multiple range test, 5% level. Values that don’t share the same letter are significantly different.

**Table (2) Fruit set percentage (after 165 & 205days) of Washington navel orange trees as affected by gibberellic acid and potassium nitrate during 2008 and2009 seasons.**

| Experiment (1)        | Fruit set % (after 165days) |                     |              |              | Fruit set % (after 205days) |                     |              |              |
|-----------------------|-----------------------------|---------------------|--------------|--------------|-----------------------------|---------------------|--------------|--------------|
|                       | 2008, Season                |                     | 2009, Season |              | 2008, Season                |                     | 2009, Season |              |
|                       | Fruit cluster (Fruit/Tree)  | Fruit diameter (mm) | (Fruit/Tree) | (Fruit/Tree) | Fruit cluster (Fruit/Tree)  | Fruit diameter (mm) | (Fruit/Tree) | (Fruit/Tree) |
| Control               | 5.97g                       | 5.97g               | 5.97g        | 5.97g        | 6.56g                       | 6.56g               | 6.56g        | 6.56g        |
| G.A., 50ppm           | 7.61g                       | 7.61g               | 7.61g        | 7.61g        | 8.17g                       | 8.17g               | 8.17g        | 8.17g        |
| G.A., 100ppm          | 6.76f                       | 6.76f               | 7.46g        | 6.90f        | 7.51f                       | 6.94g               | 7.86g        | 7.26f        |
| G.A., 150ppm          | 9.46g                       | 9.46g               | 9.46g        | 9.46g        | 8.65g                       | 7.86g               | 8.65g        | 9.12g        |
| G.A., 200ppm          | 6.76f                       | 6.76f               | 7.46g        | 6.90f        | 7.51f                       | 6.94g               | 7.86g        | 7.26f        |
| G.A., 250ppm          | 6.76f                       | 6.76f               | 7.46g        | 6.90f        | 7.51f                       | 6.94g               | 7.86g        | 7.26f        |
| KNO <sub>3</sub> , 2% | 6.97g                       | 6.97g               | 6.97g        | 6.97g        | 6.51g                       | 6.51g               | 6.51g        | 6.51g        |
| KNO <sub>3</sub> , 4% | 6.76f                       | 6.76f               | 7.46g        | 6.90f        | 7.51f                       | 6.94g               | 7.86g        | 7.26f        |
| KNO <sub>3</sub> , 6% | 6.97g                       | 6.97g               | 6.97g        | 6.97g        | 6.51g                       | 6.51g               | 6.51g        | 6.51g        |
| Mean (Error)          | 6.76f                       | 6.76f               | 7.46g        | 6.90f        | 7.51f                       | 6.94g               | 7.86g        | 7.26f        |

Mean separation within columns by Duncan’s multiple range test, 5% level. Values that don’t share the same letter are significantly different.

**Table ( 3 ) Fruit weight and number of Washington navel orange trees as affected by gibberellic acid and potassium nitrate during 2008 and2009 seasons.**


| Experiment (1)        | Fruit weight (g)           |                     |              |              | Fruit number (g)           |                     |              |              |
|-----------------------|----------------------------|---------------------|--------------|--------------|----------------------------|---------------------|--------------|--------------|
|                       | 2008, Season               |                     | 2009, Season |              | 2008, Season               |                     | 2009, Season |              |
|                       | Fruit cluster (Fruit/Tree) | Fruit diameter (mm) | (Fruit/Tree) | (Fruit/Tree) | Fruit cluster (Fruit/Tree) | Fruit diameter (mm) | (Fruit/Tree) | (Fruit/Tree) |
| Control               | 212g                       | 212g                | 212g         | 212g         | 240                        | 240                 | 240          | 240          |
| G.A., 50ppm           | 286                        | 221g                | 231g         | 240          | 19g                        | 246g                | 246          | 246          |
| G.A., 100ppm          | 221g                       | 216                 | 216          | 216          | 240                        | 240                 | 240          | 240          |
| G.A., 150ppm          | 286                        | 216                 | 216          | 216          | 240                        | 240                 | 240          | 240          |
| G.A., 200ppm          | 286                        | 216                 | 216          | 216          | 240                        | 240                 | 240          | 240          |
| G.A., 250ppm          | 286                        | 216                 | 216          | 216          | 240                        | 240                 | 240          | 240          |
| KNO <sub>3</sub> , 2% | 286                        | 216                 | 216          | 216          | 240                        | 240                 | 240          | 240          |
| KNO <sub>3</sub> , 4% | 286                        | 216                 | 216          | 216          | 240                        | 240                 | 240          | 240          |
| KNO <sub>3</sub> , 6% | 286                        | 216                 | 216          | 216          | 240                        | 240                 | 240          | 240          |
| Mean (Error)          | 212g                       | 212g                | 212g         | 212g         | 240                        | 240                 | 240          | 240          |

Mean separation within columns by Duncan’s multiple range test, 5% level. Values that don’t share the same letter are significantly different.


**Table (4) Yield (Yield Kg. /tree and ton / fed.) of Washington navel orange trees as affected by gibberellic acid and potassium nitrate during 2008 and 2009 seasons.**

| Treatment(T)                            | 2008, Season                                     |   |   |         | 2009, Season                                     |   |   |         | 2008, Season                                     |   |   |         | 2009, Season                                     |   |   |         |
|---|--|---|---|---------|--|---|---|---------|--|---|---|---------|--|---|---|---------|
|   | Full Bloom (Time) <sup>1</sup> (T <sup>1</sup> ) | Fruit diameter (Time <sup>1</sup> -1.5Fruit diameter from (Time <sup>1</sup> )) (T <sup>1</sup> ) | (Time <sup>1</sup> ) <sup>2</sup> (T <sup>2</sup> ) | Mean(T) | Full Bloom (Time) <sup>1</sup> (T <sup>1</sup> ) | Fruit diameter (Time <sup>1</sup> -1.5Fruit diameter from (Time <sup>1</sup> )) (T <sup>1</sup> ) | (Time <sup>1</sup> ) <sup>2</sup> (T <sup>2</sup> ) | Mean(T) | Full Bloom (Time) <sup>1</sup> (T <sup>1</sup> ) | Fruit diameter (Time <sup>1</sup> -1.5Fruit diameter from (Time <sup>1</sup> )) (T <sup>1</sup> ) | (Time <sup>1</sup> ) <sup>2</sup> (T <sup>2</sup> ) | Mean(T) | Full Bloom (Time) <sup>1</sup> (T <sup>1</sup> ) | Fruit diameter (Time <sup>1</sup> -1.5Fruit diameter from (Time <sup>1</sup> )) (T <sup>1</sup> ) | (Time <sup>1</sup> ) <sup>2</sup> (T <sup>2</sup> ) | Mean(T) |
| Control                                 | 75.62  | 75.62   | 75.62   | 75.96C  | 77.54A   | 77.54A  | 77.54A  | 77.96C  | 11.52  | 11.52   | 11.52   | 11.82C  | 12.84A   | 12.84A  | 12.84A  | 12.84C  |
| GA <sub>3</sub> 50ppm                   | 127.56a  | 127.56a   | 127.56a   | 122.2A  | 133.66a  | 133.66a   | 133.66a   | 122.2B  | 20.16ab  | 19.76ab   | 20.16ab   | 19.76a  | 21.56ab  | 19.86a  | 20.96a  | 19.57B  |
| GA <sub>3</sub> 100ppm                  | 139.66a  | 131.96ab  | 126.96a   | 126.2A  | 133.76a  | 135.26a   | 134.76a   | 117.8C  | 20.96a   | 17.66ab   | 21.82a  | 20.2A   | 19.66a   | 19.86a  | 20.26a  | 19.86C  |
| GA <sub>3</sub> 50ppm+KNO <sub>3</sub>  | 127.56a  | 121.26ab  | 122.66a   | 127.8A  | 139.96a  | 137.76a   | 134.76a   | 127.5A  | 20.16ab  | 19.46ab   | 21.16ab   | 20.2A   | 22.36a   | 17.26ab   | 21.46a  | 20.37A  |
| GA <sub>3</sub> 100ppm+KNO <sub>3</sub> | 124.76a  | 122.86ab  | 125.26a   | 127.4A  | 126.76a  | 137.76a   | 128.76a   | 121.8C  | 19.66ab  | 19.76ab   | 21.46a  | 20.2A   | 20.56a   | 17.16ab   | 20.66a  | 19.98C  |
| KNO <sub>3</sub> 2%                     | 79.96b   | 74.62b  | 86.12bc   | 80.42C  | 76.62A   | 76.62A  | 83.81   | 82.82B  | 12.76b   | 11.96b  | 14.42bc   | 13.86C  | 12.91A   | 12.31A  | 14.81   | 13.22B  |
| KNO <sub>3</sub> 4%                     | 115.76ab   | 98.82ab   | 102.76b   | 106.7B  | 112.26b  | 99.22b  | 111.16ab  | 109.22B | 18.12ab  | 15.16ab   | 16.42bc   | 16.22B  | 18.11ab  | 15.86b  | 18.62ab   | 17.48B  |
| KNO <sub>3</sub> 6%                     | 105.86ab   | 97.76ab   | 102.76b   | 101.8B  | 105.66b  | 91.62b  | 101.66b   | 100.62B | 18.12ab  | 15.76ab   | 16.76bc   | 16.22B  | 16.96a   | 14.96b  | 16.76a  | 16.14B  |
| Mean (T <sup>1</sup> )                  | 118.4A   | 101.7B  | 102.2A  |         | 112.2A   | 96.22B  | 112.7A  |         | 17.66A   | 16.27B  | 17.97A  |         | 17.97A   | 16.48B  | 18.27A  |         |

Mean separation within columns by Duncan's multiple range test, 5% level. Values that don't share the same letter are significant different.



**Fig. (2) Fruit splitting percentage of Washington navel orange trees as affected by gibberellic acid and potassium nitrate during 2008 season.**



**Fig. (2) Fruit splitting percentage of Washington navel orange trees as affected by gibberellic acid and potassium nitrate during 2009 season.**

**potassium nitrate during 2009 season.**

**Table (5) Nitrite in fruit juice of Washington navel orange trees as affected by gibberellic acid and potassium nitrate during 2008 and 2009 seasons.**

| Treatments(T)                     | 2008, Season       |   |                   |         | 2009, Season       |   |                   |         |
|-----------------------------------|--------------------|---|-------------------|---------|--------------------|---|-------------------|---------|
|                                   | Full Bloom (Time1) | Fruit diameter from 1.52-2.04cm (Time2) | (Time1) + (Time2) | Mean(T) | Full Bloom (Time1) | Fruit diameter from 1.52-2.04cm (Time2) | (Time1) + (Time2) | Mean(T) |
|                                   | Control            | 3.13abf                                 | 3.13abf           | 3.13abf | 3.13B              | 3.10g                                   | 3.10g             | 3.10g   |
| GA <sub>3</sub> 50ppm             | 3.13abf            | 3.13abf                                 | 3.13abf           | 3.13B   | 3.10g              | 3.10g                                   | 3.10g             | 3.10C   |
| GA <sub>3</sub> 100ppm            | 3.08f              | 3.13abf                                 | 3.08f             | 3.08B   | 3.10g              | 3.10g                                   | 3.10g             | 3.10C   |
| GA <sub>3</sub> 50ppm + 0.5%area  | 3.13abf            | 3.13abf                                 | 3.13abf           | 3.13B   | 3.10ag             | 3.10ad                                  | 3.10ag            | 3.10B   |
| GA <sub>3</sub> 100ppm + 0.5%area | 3.13abf            | 3.13abf                                 | 3.13abf           | 3.13B   | 3.10ag             | 3.10ag                                  | 3.10ag            | 3.10B   |
| KNO <sub>3</sub> 2%               | 3.08ab             | 3.13ab                                  | 3.08ab            | 3.08B   | 3.08ab             | 3.08ab                                  | 3.08ab            | 3.08A   |
| KNO <sub>3</sub> 4%               | 3.13abf            | 3.13ab                                  | 3.13ab            | 3.13B   | 3.08ab             | 3.08ab                                  | 3.08ab            | 3.08A   |
| KNO <sub>3</sub> 6%               | 3.08ab             | 3.08ab                                  | 3.08ab            | 3.08B   | 3.08ab             | 3.08ab                                  | 3.08ab            | 3.08A   |
| Mean (Time)                       | 3.09 A             | 3.09 A                                  | 3.09 A            | 3.09 A  | 3.09 A             | 3.09 A                                  | 3.09 A            | 3.09 A  |

Mean separation within columns by Duncan's multiple range test, 5% level. Values that don't share the same letter are significantly different.

**Table (6) Nitrate in fruit juice of Washington navel orange trees as affected by gibberellic acid and potassium nitrate during 2008 and 2009 seasons.**

| Treatments(T)                     | 2008, Season       |   |                   |          | 2009, Season       |   |                   |         |
|-----------------------------------|--------------------|---|-------------------|----------|--------------------|---|-------------------|---------|
|                                   | Full Bloom (Time1) | Fruit diameter from 1.52-2.04cm (Time2) | (Time1) + (Time2) | Mean(T)  | Full Bloom (Time1) | Fruit diameter from 1.52-2.04cm (Time2) | (Time1) + (Time2) | Mean(T) |
|                                   | Control            | 46.40efg                                | 46.40efg          | 46.40efg | 46.40C             | 43.38fg                                 | 43.38fg           | 43.38fg |
| GA <sub>3</sub> 50ppm             | 40.52jkl           | 42.65ij                                 | 44.66ghi          | 42.61D   | 38.41i             | 39.53hi                                 | 44.66f            | 40.83E  |
| GA <sub>3</sub> 100ppm            | 40.51jkl           | 40.49jkl                                | 40.49jkl          | 40.50E   | 43.35fg            | 42.56fg                                 | 44.57f            | 43.49D  |
| GA <sub>3</sub> 50ppm + 0.5%area  | 38.24L             | 39.61kl                                 | 40.71jk           | 39.53E   | 38.61i             | 39.72hi                                 | 41.61gh           | 39.58E  |
| GA <sub>3</sub> 100ppm + 0.5%area | 40.60jkl           | 43.67hi                                 | 46.54efg          | 43.60D   | 48.41e             | 48.52e                                  | 54.52bc           | 50.48C  |
| KNO <sub>3</sub> 2%               | 47.41def           | 48.56cde                                | 45.30fgh          | 47.49C   | 53.68bcd           | 48.79c                                  | 51.75d            | 51.34C  |
| KNO <sub>3</sub> 4%               | 47.2bcf            | 48.66cde                                | 49.72bcd          | 48.55D   | 52.3abcd           | 52.3abcd                                | 55.72b            | 53.54D  |
| KNO <sub>3</sub> 6%               | 50.94abc           | 51.14ab                                 | 52.54a            | 51.54A   | 54.47bc            | 55.59b                                  | 59.62a            | 56.56A  |
| Mean (Time)                       | 43.99 B            | 45.15 A                                 | 45.80 A           | 46.89 B  | 46.31 B            | 46.31 B                                 | 50.62 A           | 48.62 A |

Mean separation within columns by Duncan's multiple range test, 5% level. Values that don't share the same letter are significantly different

**Physical properties of fruits**

**Fruit diameter and length (mm)**

Data in Table (7) indicated that the statistical analysis showed no significant differences between all treatments on Washington navel orange fruits in the first season, while trees sprayed with GA<sub>3</sub> 100 ppm enhanced significant increased fruit diameter and length (mm) in the second season. Anyhow, the differences between all treatments were low to be significant.

**Fruit size (mL)**

It is clear from Table (8) that, all treatments increased fruit size as comparing with control especially with high concentration of potassium nitrate in the first and second season (2008 & 2009) may be due to increasing of fruit peel thickness. The present results are in a general harmony with (Okada, et al., 1994) on Satsuma mandarin they mentioned that fruit size increased as K fertilization increased, also (Abd El-Rahman, 2005) found that spraying with promalin at 45 ppm and GA<sub>3</sub> at 50 ppm increased fruit size on navel orange fruits.

**Peel thickness (mm)**

Data in Table (8) showed that, peel thickness was affected by different treatments in the first and second seasons, and it is clear that, GA<sub>3</sub> treatments

reduced significant values for fruit peel thickness, on the other hand, it is noticed that, peel thickness increased with increasing the concentration of potassium nitrate. The same observation was reported by (Zaied, et al., 2006) on Washington navel orange.

Juice weight percentage of fruits is shown in Table (9); the changes of juice weight percentage were slightly fluctuated during two seasons. No obvious trend could be detected between all treatments. These results agree with the finding of (Abd El-Rahman, 2005) on navel orange trees, who mentioned that spraying gibberellic acid had no significant effect on juice weight percentage. In the same direction (Davies et al., 1999) stated that, effects of GA<sub>3</sub> on juice fruits were variable, this inconsistency may be related to time of application.

**Chemical properties**

**T.S.S percentage:**

Table (9) clearly showed that all treatments did not show a constant trend during two seasons for T.S.S percentage in fruit juice of Washington navel orange. Data also indicated that, spraying trees with KNO<sub>3</sub> at 4% gave the lowest significant values for T.S.S % comparing with other treatments. Anyhow, the differences between all treatments were low to be

significant.

A similar trend was obtained with navel orange (Ibrahim, *et al.* 1994 and Abel El-Rahman, 2005). On the other hand, (Mostafa and Saleh 2006) found that T.S.S content in the fruit juice increased with potassium sprays on Balady mandarin trees.

**Acidity**

Table (10) reflects that trees sprayed with GA<sub>3</sub> 50 ppm had the highest significant acidity for Washington navel orange fruits in the first season, while, trees treated by potassium nitrate treatments gave the maximum acidity for their fruits in the second seasons especially with higher concentration. Meanwhile, the other treatments scored the intermediate significant values in this regard for two seasons. These results are in the same line with those obtained by (Ibrahim, *et al.*, 1994) who found that total acidity were no significant effected by GA<sub>3</sub> treatment. Also, (Erner, *et al.* 1993) reported that trees spraying with KNO<sub>3</sub> increased juice acidity on Shamouti and Valencia oranges.

**T.S.S acid ratio**

It is evident from Table (10) that T.S.S acid ratio did not show any obvious trend in the first and second season. Concerning the time of foliar spraying, the results indicated that trees sprayed at second time (fruit diameter from 1.5-2.0 cm) recorded the highest significant values for T.S.S. acid ratio with KNO<sub>3</sub> 2% followed by GA<sub>3</sub> 100 ppm + 0.5% urea on Washington navel orange fruits than the other times of applications in the first season

(2008). While, GA<sub>3</sub> 100 ppm + 0.5% urea gave the highest significant values for T.S.S. acid ratio in (second application and first application) respectively in the second season (2009). Also, it could be noticed that, the differences between all treatments were high to be significant.

**Conclusion**

From the above mentioned results, it could be concluded that, fruit set, yield as weight ton/fed., and fruit quality especially in respect with reducing nitrite and nitrate in fruit juice of Washington navel orange were greatly improved with spraying trees GA<sub>3</sub> 50 ppm with or without 0.5 % urea. Concerning the application times it could be noticed that, both full bloom (first application ) or full bloom + fruit diameter from 1.5-2.0 cm (first and second time of application) gave the best results so, we could be recommended to use (first application ) to reduce the costs for growers and giving the best yield and fruit quality. Treatments that increase brix and reduce acid early in the season provide two benefits. Growers can gain the economic advantage of harvesting early and can avoid the economic loss due worsening crease as harvest is delayed as mentioned by (Carol J. Lovatt, 2000).

**Corresponding author**

Abd El-Rahman, GF.  
Citrus Department, Hort. Res. Institut., Agric. Res. Center, Giza, Egypt  
[gamalelshamanty@yahoo.com](mailto:gamalelshamanty@yahoo.com)

**Table ( 7 ) Fruit diameter and length of Washington navel orange trees as affected by gibberellic acid and potassium nitrate during 2008 and 2009 seasons.**

| Treatments(T)                      | Fruit diameter (cm)             |  |          |         |                                 |  |          |         | Fruit length (cm)               |  |          |         |                                 |  |          |         |
|------------------------------------|---------------------------------|--|----------|---------|---------------------------------|--|----------|---------|---------------------------------|--|----------|---------|---------------------------------|--|----------|---------|
|                                    | 2008 Season                     |  |          |         | 2009 Season                     |  |          |         | 2008 Season                     |  |          |         | 2009 Season                     |  |          |         |
|                                    | Full Bloom (1.5-2.0cm) (Treat1) | Fruit diameter from 1.5-2.0cm (Treat2) | (Treat3) | Mean(T) | Full Bloom (1.5-2.0cm) (Treat1) | Fruit diameter from 1.5-2.0cm (Treat2) | (Treat3) | Mean(T) | Full Bloom (1.5-2.0cm) (Treat1) | Fruit diameter from 1.5-2.0cm (Treat2) | (Treat3) | Mean(T) | Full Bloom (1.5-2.0cm) (Treat1) | Fruit diameter from 1.5-2.0cm (Treat2) | (Treat3) | Mean(T) |
| Control                            | 75.6abc                         | 75.6abc                                | 75.6abc  | 75.6A   | 79.6abcd                        | 79.6abcd                               | 79.6abcd | 79.6AB  | 79.6bc                          | 79.6bc                                 | 79.6bc   | 79.6bc  | 81.6abc                         | 81.6abc                                | 81.6abc  | 81.6BC  |
| GA <sub>3</sub> 50ppm              | 77.5abc                         | 74.6abc                                | 75.41c   | 74.8BA  | 79.6abcd                        | 80.9bcd                                | 81.6bc   | 79.6AB  | 81.6bc                          | 79.7bc                                 | 80.9bc   | 80.5bc  | 81.2abc                         | 80.9bc                                 | 81.2abc  | 81.1BC  |
| GA <sub>3</sub> 100ppm             | 75.2abc                         | 74.6abc                                | 74.2abc  | 74.6BA  | 82.9bcd                         | 82.9bcd                                | 82.9bcd  | 82.7A   | 79.7bc                          | 81.2bc                                 | 77.5bc   | 79.6bc  | 81.1abc                         | 81.5bc                                 | 82.9bc   | 81.6A   |
| GA <sub>3</sub> 50ppm + 0.5% urea  | 74.6abc                         | 74.6abc                                | 73.6abc  | 74.6BA  | 81.6bcd                         | 81.5bc                                 | 82.5bcd  | 81.6AB  | 79.6bc                          | 80.7bc                                 | 79.7bc   | 79.7bc  | 82.7abc                         | 82.1abc                                | 82.5abc  | 82.0BC  |
| GA <sub>3</sub> 100ppm + 0.5% urea | 74.6abc                         | 72.3abc                                | 74.2abc  | 73.6BA  | 80.3bcd                         | 81.6bcd                                | 82.5bcd  | 81.6AB  | 79.2bc                          | 78.4bc                                 | 79.2bc   | 79.2bc  | 81.6abc                         | 80.7bc                                 | 81.2abc  | 81.1AB  |
| KNO <sub>3</sub> 2%                | 72.6abc                         | 77.6abc                                | 76.6bc   | 74.6BA  | 75.5bcd                         | 79.5bcd                                | 80.5bcd  | 78.6BC  | 77.4bc                          | 80.5bc                                 | 81.0bc   | 80.5bc  | 79.7bc                          | 82.6abc                                | 81.2abc  | 81.0BC  |
| KNO <sub>3</sub> 4%                | 78.9abc                         | 77.6abc                                | 75.6abc  | 77.2BA  | 80.5bcd                         | 79.6bcd                                | 77.2bcd  | 79.2BC  | 80.9bc                          | 80.9bc                                 | 82.0bc   | 80.6bc  | 81.5abc                         | 82.6abc                                | 80.0bc   | 81.0BC  |
| KNO <sub>3</sub> 6%                | 77.6abc                         | 77.1abc                                | 77.1abc  | 77.6BA  | 79.5bcd                         | 79.5bcd                                | 79.5bcd  | 79.9BC  | 79.7bc                          | 79.6bc                                 | 82.0bc   | 80.5bc  | 82.5abc                         | 81.5abc                                | 81.5abc  | 81.0BC  |
| Mean (T-test)                      | 75.75A                          | 74.8A                                  | 75.15A   | 75.25A  | 80.02A                          | 80.85A                                 | 81.65A   | 80.82A  | 79.68A                          | 80.87A                                 | 80.68A   | 80.68A  | 82.21A                          | 82.21A                                 | 82.95A   |         |

Mean separation within columns by Duncan's multiple range test, 5% level. Values that don't share the same letter are significantly different.

**Table (8) Fruit size and Peel thickness of Washington navel orange trees as affected by gibberellic acid and potassium nitrate during 2008 and 2009 seasons.**

| Treatments(T)                           | 2008, Season     |                               |                             |         | 2009, Season     |                               |                             |         | 2008, Season     |                               |                             |         | 2009, Season     |                               |                             |         |
|---|------------------|-------------------------------|-----------------------------|---------|------------------|-------------------------------|-----------------------------|---------|------------------|-------------------------------|-----------------------------|---------|------------------|-------------------------------|-----------------------------|---------|
|   | Fruit Weight (g) | Fruit diameter (mm) ± (S.E.M) | Fruit length (mm) ± (S.E.M) | Mean(T) | Fruit Weight (g) | Fruit diameter (mm) ± (S.E.M) | Fruit length (mm) ± (S.E.M) | Mean(T) | Fruit Weight (g) | Fruit diameter (mm) ± (S.E.M) | Fruit length (mm) ± (S.E.M) | Mean(T) | Fruit Weight (g) | Fruit diameter (mm) ± (S.E.M) | Fruit length (mm) ± (S.E.M) | Mean(T) |
|   | Control          | 208                           | 268                         | 208     | 208BC            | 206                           | 276                         | 206     | 206              | 4.80                          | 4.80                        | 4.80    | 4.80D            | 4.70                          | 4.70                        | 4.70    |
| GA <sub>3</sub> 50ppm                   | 270              | 276                           | 276                         | 284.00  | 366ab,d          | 336ab                         | 372ab                       | 358BC   | 3.80             | 3.610                         | 3.70                        | 3.680   | 3.90             | 3.70                          | 3.80                        | 3.87C   |
| GA <sub>3</sub> 100ppm                  | 260              | 276                           | 243                         | 264.79  | 326ab,f          | 310ab                         | 326ab,f                     | 314.70  | 3.80             | 4.53                          | 4.20                        | 4.20C   | 3.70             | 3.65                          | 4.00                        | 3.68    |
| GA <sub>3</sub> 200ppm+KNO <sub>3</sub> | 270              | 276                           | 276                         | 283.76  | 306              | 326ab,f                       | 326ab,f                     | 310     | 3.90             | 4.53                          | 4.00                        | 4.20C   | 3.80             | 3.70                          | 3.20                        | 3.88    |
| GA <sub>3</sub> 100ppm+KNO <sub>3</sub> | 280              | 276                           | 280                         | 284.70  | 326ab,f          | 377abg                        | 336ab                       | 332.879 | 4.90             | 4.10                          | 4.00                        | 4.20    | 3.80             | 3.80                          | 3.70                        | 3.78    |
| KNO <sub>3</sub> 2%                     | 260              | 276                           | 267                         | 273.80  | 326ab            | 336ab                         | 366ab,d                     | 343.349 | 4.70             | 4.87ab                        | 4.20                        | 4.70C   | 4.80             | 4.20                          | 3.80                        | 4.60    |
| KNO <sub>3</sub> 4%                     | 260              | 276                           | 276                         | 276.70  | 326ab            | 366                           | 356ab                       | 346.740 | 4.70             | 5.10                          | 4.80d                       | 4.70D   | 4.70             | 4.70                          | 4.60                        | 4.68    |
| KNO <sub>3</sub> 6%                     | 260              | 276                           | 270                         | 268.64  | 366ab            | 336ab,d                       | 366                         | 368.34  | 4.70             | 4.60                          | 4.80                        | 4.70A   | 4.80             | 4.50                          | 4.80                        | 4.64    |
| Mean (T Error)                          | 268.5A           | 268.0B                        | 268.8C                      |         | 323.9A           | 328.3A                        | 328.6A                      |         | 4.80C            | 4.53A                         | 4.40B                       |         | 3.80C            | 3.80B                         | 4.01A                       |         |

Mean separation within columns by Duncan's multiple range test, 5% level. Values that don't share the same letter are significantly different

**Table (9) Juice weight and T.S.S percentage in Washington navel orange fruits as affected by gibberellic acid and potassium nitrate during 2008 and 2009 seasons.**

| Treatments(T)                           | 2008, Season     |                     |                     |         | 2009, Season     |                     |                     |         | 2008, Season     |                     |                     |          | 2009, Season     |                     |                     |         |
|---|------------------|---------------------|---------------------|---------|------------------|---------------------|---------------------|---------|------------------|---------------------|---------------------|----------|------------------|---------------------|---------------------|---------|
|   | Juice weight (g) | T.S.S (%) ± (S.E.M) | T.S.S (%) ± (S.E.M) | Mean(T) | Juice weight (g) | T.S.S (%) ± (S.E.M) | T.S.S (%) ± (S.E.M) | Mean(T) | Juice weight (g) | T.S.S (%) ± (S.E.M) | T.S.S (%) ± (S.E.M) | Mean(T)  | Juice weight (g) | T.S.S (%) ± (S.E.M) | T.S.S (%) ± (S.E.M) | Mean(T) |
|   | Control          | 56.80g              | 56.80g              | 56.80g  | 56.80C           | 55.70g              | 55.70g              | 55.70g  | 55.70C           | 10.50g              | 10.50g              | 10.50g   | 10.50ABC         | 11.00g              | 11.00g              | 11.00g  |
| GA <sub>3</sub> 50ppm                   | 59.70            | 61.50               | 62.70               | 61.3A   | 51.70            | 56.50g              | 54.80g              | 54.80C  | 10.50g           | 10.50g              | 10.50g              | 10.50ABC | 10.50g           | 10.50g              | 10.50g              | 10.50B  |
| GA <sub>3</sub> 100ppm                  | 61.50            | 60.00               | 58.50               | 60.00   | 55.50g           | 54.80g              | 53.40g              | 54.50C  | 10.50g           | 11.00g              | 10.50g              | 10.50ABC | 11.00g           | 11.00g              | 10.50g              | 10.50AB |
| GA <sub>3</sub> 200ppm+KNO <sub>3</sub> | 60.00            | 61.50               | 61.50g              | 60.80   | 56.50g           | 52.60               | 53.40g              | 54.50C  | 9.50g            | 9.50g               | 10.25g              | 9.80BC   | 10.50g           | 9.50g               | 10.25g              | 10.00BC |
| GA <sub>3</sub> 100ppm+KNO <sub>3</sub> | 56.70g           | 55.50               | 56.60               | 56.20   | 52.60            | 55.80g              | 56.80g              | 54.30   | 11.00g           | 10.50g              | 10.50g              | 10.50A   | 11.00g           | 10.50g              | 10.50g              | 10.50AB |
| KNO <sub>3</sub> 2%                     | 56.80g           | 55.50g              | 54.80g              | 55.30C  | 55.50g           | 57.50g              | 60.00g              | 57.60   | 11.00g           | 10.50g              | 10.50g              | 10.50AB  | 10.50g           | 11.00g              | 10.50g              | 10.70AB |
| KNO <sub>3</sub> 4%                     | 58.00g           | 55.00               | 56.60               | 56.20   | 56.60g           | 62.50               | 62.50               | 60.5A   | 10.50g           | 9.50g               | 9.50g               | 9.50C    | 10.00g           | 9.50g               | 9.50g               | 9.70C   |
| KNO <sub>3</sub> 6%                     | 59.70            | 58.00               | 58.00               | 58.50   | 58.00            | 58.00               | 58.70               | 58.20   | 11.00g           | 9.80g               | 10.50g              | 10.50ABC | 9.80g            | 10.50g              | 10.50g              | 10.10BC |
| Mean (T Error)                          | 58.87A           | 56.20C              | 57.20B              |         | 55.40B           | 56.21AB             | 56.71A              |         | 10.61A           | 10.61A              | 10.61A              |          | 10.61A           | 10.61A              | 10.61A              |         |

Mean separation within columns by Duncan's multiple range test, 5% level. Values that don't share the same letter are significantly different.

**Table (10) Acidity percentage and T.S.S / Acid ratio in Washington navel orange fruits as affected by gibberellic acid and potassium nitrate during 2008 and 2009 seasons.**

| Treatments(T)                           | 2008, Season |                     |                     |         | 2009, Season |                     |                     |         | 2008, Season |                     |                     |         | 2009, Season |                     |                     |         |
|---|--------------|---------------------|---------------------|---------|--------------|---------------------|---------------------|---------|--------------|---------------------|---------------------|---------|--------------|---------------------|---------------------|---------|
|   | Acidity (%)  | T.S.S (%) ± (S.E.M) | T.S.S (%) ± (S.E.M) | Mean(T) | Acidity (%)  | T.S.S (%) ± (S.E.M) | T.S.S (%) ± (S.E.M) | Mean(T) | Acidity (%)  | T.S.S (%) ± (S.E.M) | T.S.S (%) ± (S.E.M) | Mean(T) | Acidity (%)  | T.S.S (%) ± (S.E.M) | T.S.S (%) ± (S.E.M) | Mean(T) |
|   | Control      | 1.07g               | 1.07g               | 1.07g   | 1.07C        | 0.90g               | 0.90g               | 0.90g   | 0.90C        | 9.01                | 9.01                | 9.01    | 9.01C        | 11.54               | 11.54               | 11.54   |
| GA <sub>3</sub> 50ppm                   | 1.25         | 1.36                | 1.22a               | 1.23A   | 0.93a        | 0.91a               | 0.76                | 0.82D   | 8.80         | 8.80                | 8.80                | 8.80C   | 12.50        | 12.50               | 12.50               | 12.50B  |
| GA <sub>3</sub> 100ppm                  | 1.07g        | 1.02a               | 1.16a               | 1.08    | 0.93a        | 0.90a               | 0.90g               | 0.90C   | 9.54f        | 9.52                | 9.50g               | 9.50B   | 12.50        | 12.50               | 10.45               | 11.60C  |
| GA <sub>3</sub> 200ppm+KNO <sub>3</sub> | 1.06f        | 1.05g               | 1.16a               | 1.090C  | 0.90f        | 0.91a               | 0.80                | 0.89C   | 9.15g        | 9.15g               | 9.05g               | 9.05B   | 10.65        | 11.50               | 11.54               | 11.00D  |
| GA <sub>3</sub> 100ppm+KNO <sub>3</sub> | 1.06f        | 0.91                | 1.07g               | 1.00    | 0.92a        | 0.70                | 0.83g               | 0.79E   | 10.10        | 11.20               | 9.54f               | 10.20B  | 11.65        | 14.30               | 12.45               | 13.00A  |
| KNO <sub>3</sub> 2%                     | 1.23ab       | 0.80                | 0.90                | 0.90F   | 1.00a        | 1.07ab              | 1.02f               | 1.03A   | 9.02         | 11.62a              | 10.45               | 10.72A  | 9.70         | 10.27b              | 10.60               | 10.20E  |
| KNO <sub>3</sub> 4%                     | 0.90         | 1.06af              | 0.90                | 0.90F   | 1.10         | 0.90f               | 1.00                | 1.00A   | 10.00        | 9.90                | 9.90                | 9.90C   | 9.90         | 9.90                | 9.90                | 9.90F   |
| KNO <sub>3</sub> 6%                     | 1.17a        | 0.90                | 1.02a               | 1.030E  | 1.10ab       | 1.00ab              | 1.10ab              | 1.07A   | 9.90         | 9.15g               | 9.02                | 9.40D   | 9.45         | 9.00                | 9.50                | 9.50F   |
| Mean (T Error)                          | 1.08a        | 1.00C               | 1.07B               |         | 0.97A        | 0.93B               | 0.80B               |         | 9.50B        | 9.81A               | 9.60B               |         | 11.00B       | 11.05A              | 10.60B              |         |

Mean separation within columns by Duncan's multiple range test, 5% level. Values that don't share the same letter are significantly different.

**References:**

**A.O.A.C.,(1995).** Official methods of analysis of A.O.A.C. Pub.By official A.O.A.C. chapter 4, p.18-20, 37, p. 10, 44 p.8-9.  
**Abd El-Rahman, G.F. (2005).** Physiological and

anatomical studies on fruit drop in navel orange trees. Ph.D. Thesis, Fac. Agric., Al Azhar Univ.  
**Abde El-Migeed, M.M.M.;Saleh, M.M.S. and Mostafa, E.A.M. (2007).** The beneficial effect of minimizing mineral nitrogen fertilization on Washington navel orange



- trees by using organic and bio-fertilizers, World Journal of Agric. Sci. 3 (1): 80-85.
- Agusti, M.F and V.O. Almela.( 1991).** Aplicación de fitoreguladores em citricultura. Barcelona, Aedos., pp. 261.
- Babu, -Ghur and Lavania, ML. (1985).** Effect of plant growth regulators on fruit set and fruit drop of plant lemon-1 (*Citrus limon Burn*). Indian Journal of Horticulture, 42 (3/4) 237-240.
- Brosh,P. and Monselise, S.P. (1977).** Increasing yields of "Topaz" mandarin by gibberellin and girdling in the presence of "Minneola" Pollinizers. Scientia Hort. 7:369-372.
- Brosh, P.; Oren, Y. and Shani,M. (1975).**Preharvest autumn drop of grapefruit in central area. Alon ha Notea, 29: 506-509 (Hebrew).
- Carol J. Lovatt.(2000).** Use of Urea and Growth Regulators to Increase Fruit Set and/or Size and Quality of Citrus.Citrus\_Research Board 2000 Annual Report. Department of Botany and Plant Sciences, University of California at Riverside
- Clarke, G.M.and Kempson, R.E. (1977).**Introduction to the design and analysis of experiments. Arnold, A. Member of the Holder-Headline Group. 1<sup>st</sup> Edt. London, UK.
- Davies, F.S.;Campbell,C.A. and Fidelibus, M.W. (2001).** Gibberellic acid tank mix and adjuvant effects on peel quality and juice yield of "Hamlin" oranges. Hort Technology 11:171-174.
- Davies, F.S.;Campbell, C.A.; Zalman, G. and Fidelibus, M. (1999).** Gibberellic acid application timing effects on juice yield and peel quality of "Hamlin" oranges, Proc. Fla. State Hort. Soc. 112: 22-24.
- Davies, P.J., (1987).** Plant hormones and their role in plant growth and development. Martinus Nijhoff Publishers, Dordrecht, p. 681.
- Duncan, D.B. (1955).** Multiple ranges and multiple F-tests. Biometrics, 11: 1-42.
- El-Deeb,M.D.E. (1989).** Studies on the nutritional status of citrus in some areas of Kaliobyia,Ph.D. Thesis. Fac.of Agric. , Moshtohor, Zagazig Univ.
- El-Fangary,M.A. (1998).** Physiological studies on growth and fruiting of citrus trees. Ph.D. Thesis, Fac. Agric. Moshtohor , Zagazig Univ.
- El-Otmani, M.(1992).** Principal growth regulator uses in citrus production.Proc.2nd.Seminar on Citrus Physiology. Bebedouro. Ed. Cargill. pp. 55-69.
- Erner,Y.; Kaplany. ; Artzi, B. and Hamou, M. (1993).** Increasing citrus fruit size using auxins and potassium. Acta Horticulturae, No. 329, 112-119.
- Fornes,F., P.J.J. Van-Rensburg, M. Sánchez- Perales and J.L. Guardiola.(1992).**Fruit setting treatments effects on two Clementine mandarin cultures.Proc.in.Soc.Citric. 1:489-492.
- Garcia, L.;Duarte, A.M.M.; Porrás, I.; Garcia, A. and Guardiola, J.L. (1994).** Fruit splitting in "Nova" hybrid mandarin in relation to the anatomy of the fruit and fruit set treatments. Scientia Horticulturae Volume 57, Issue 3, April 215-231.
- Ibrahim,T. (1994).**Water pollution, Part I. Science and life series, Egyptian organization for Books, Cairo, Egypt, pp. 58-61.
- Ibrahim,T.A.; Salem, S.E. and Guindy, L.F. (1994).**The influence of gibberellic acid and promalin on the yield and fruit quality of Washington Navel orange. Bulletin, Fac. Agric. Univ. Cairo, 45: 711-722.
- Josan, J.S., Sandhu, A.S.and Jasbir,K.(1995).**Pericarp anatomy in relation to fruit cracking in Lemon (Citrus Limon) .Indian Journal of agricultural Science, 65 (6), 410- 430.
- Krezdorn, A.H.(1965).**Fruit setting problems in citrus Proc. Carib. Reg. Amer. Soc. Hort. Sci. 9:85-92.
- Lavon, R.;Shapchiski, S., Mohel, E. and Zur.N.(1992).** Nutritional and hormonal sprays decreased fruit splitting and fruit creasing of "Nova" Hasade, 72: 1252-1257 (Hebrew).
- Mostafa, E.A.M.; Hassan, H.S.A. and El-Sabag,A.S. (2005).** Influence of spraying GA<sub>3</sub> and KNO<sub>3</sub> on yield, (2005).Influence of spraying GA<sub>3</sub> and KNO<sub>3</sub> on yield, fruit quality and leaf mineral contents of Balady mandarin trees. Minufiya J. Agric. Res. 30 (1):283-295.
- Richards,D.E., K.E. King, T. Ait-ali and N. Harberd. (2001).**How gibberellin regulates plant growth and development: A molecular genetic analysis of gibberellin signaling. Annu. Rev. Plant. Physiol. Plant. Mol. Biol. 52: 67-88.
- Ruiz, R., A. García-Luiz, C. Monerri and J.L. Guardiola.( 2001).**Carbohydrate availability in relation to fruit-set abscission in Citrus.**Annals of Botany. 87:805-812.**
- Sen,N.P. and Donaldson, B.D (1978).**Improved colorimetric method for determining nitrate and nitrite.Food J. Assoc. Anal. Chem., 6: 1389-1395.
- Smith, C.J. (1993).** Improving of fruit set of navels in the citrusdal area. In ligtings bulletin. Institute – Vir-Tropiese-en, subtropiese-Gewasse. No. 240, 27-32.
- Talón, M., F.R. Radeo,A. Gómez-Cadena and E. Primo-Millo.(1997).**Regulación hormonal de la caída de frutos inducida por defoliaciones en mandarinos Satsuma. Agric. Vergel 186:334-350.
- Tominaga,S. (1998).** GA sprays delay and reduce physiological fruit drop in Ponkan mandarin (*Citrus reticulate Blanco*), Acta Horticulturae, 463: V. III International symposium on plant bio-regulation in fruit production.
- Villafane,VE. ; Munoz, F. JE and Torres, HR. (1989).** Flowering growth and ripening of the orange Washington Valley Citrus, Acts. Agronomica, Universidad-Nacional de Colombia 39: (3-4) 142-149.
- Wardowski, W.F.; Nagy, S. and Grierson,W. (1985).**Fresh citrus fruits. Avi. Publ. Co., Inc. Westport, USA, pp. 79-83.
- Zaied, .S.; Khafagy,S.A.A. and Saleh, M.A. ( 2006).**Effect of nitrogen and potassium fertilization on vegetative growth, fruit set and quality of Washington navel orange trees. Journal of Applied Sciences Research, 2 (11): 851-857