

# The most frequent c.239A>G SNP of *NKX2.5* is not involved in Congenital Heart Disease

S. M. Dinesh, Kusuma L, and N. B. Ramachandra

Genomics Laboratory, Department of Studies in Zoology, University of Mysore, Manasagangothri, Mysore 570 006, Karnataka, India. E-mail: [nallurbr@gmail.com](mailto:nallurbr@gmail.com)

**Abstract:** Congenital heart disease (CHD) is the most developmental errors in humans, affecting 8 out of 1000 newborns. Cardiac septal defects (CSD) constitute the majority of CHD, however, the etiology of CSD remains unclear. About 85 single nucleotide polymorphisms (SNPs) of *NKX2.5* have been associated with various forms of CHD. A total of 150 CHD patients and 70 controls were screened for *NKX2.5* SNPs by PCR amplification and direct sequencing, as well as Mass array techniques. The sequence analysis was done by using Accelrys gene software. In this study, six SNPs, c.239A>G (48.66%), c.608A>G (0.6%), c.646C>T (25.33%), c.852G>A (3%), c.896C>A (0.6%) and 1212G>T (40%) were identified, of which 239A>G was the most frequent in both cases and controls. The SNP c.239A>G (rs2277923), leads into synonymous change of glutamine, in turn, the effect is neutral, thereby its frequency is almost equal in both controls and cases. Therefore, this SNP is not involved in the manifestation of CHD. [New York Science Journal 2010;3(8):43-47]. (ISSN: 1554-0200).

**Keywords:** *NKX2.5*; Congenital Heart Disease, Mass array, Direct sequencing, Accelrys gene software.

## 1. Introduction

Congenital heart disease (CHD) is the most common developmental errors in humans, affecting 8 out of 1,000 newborns (Posch et al., 2008). Cardiac septal defects (CSDs) constitute the majority of CHD, however, the etiology of CSD remains unclear. The available data suggest that fetal heart development is regulated by a group of highly conserved transcription factors, including *NKX2.5*, *TBX5*, *GATA4*, *TBX1*, *ZIC3*, *TFAP2B*, *FOG2*, and others (Ramegowda and Ramachandra, 2005; Clark et al., 2006; Chen et al., 2010). *NKX2.5*, a vertebrate homologue of *Drosophila* tinman, is a homeobox transcription factor highly expressed in early progenitors and in adult cardiomyocytes (Posch et al., 2008). Targeted disruption of *NKX2.5* in rodent models results in early embryonic lethality due to failed cardiac looping and defects in chamber formation (Lyons et al., 1995; Posch et al., 2008).

*NKX2.5* was the first gene identified as a relevant disease gene for non-syndromic human CHD (Posch et al., 2008). Schott et al., (1998) described heterozygote mutations in *NKX2.5* that resulted in atrioventricular (AV) conduction block; many genotype-positive individuals also had a secundum atrial septal defect (ASD). Subsequent family studies identified extensive variable expressivity, in addition to AV block and ASD, genotype positive individuals also had ventricular septal defect (VSD), tetralogy of fallot (TOF), double outlet right ventricle (DORV), and tricuspid valve abnormalities, including Ebstein

anomaly (Benson, 2010). These results suggest an essential role for *NKX2.5* in atrial, ventricular, and conotruncal septation, AV valve formation, and maintenance of AV conduction. These findings have been confirmed by numerous other studies (Benson, 2009).

About 85 Single nucleotide polymorphisms (SNPs) of *NKX2.5* have been associated with various forms of CHD (Schott et al., 1998; Benson et al., 1999; Hosoda et al., 1999; Kasahara et al., 2000; Goldmuntz et al., 2001; Ikeda et al., 2002; Watanabe et al., 2002; Elliot et al., 2003; McElhinney et al., 2003; Kasahara et al., 2004; Hirayama-Yamada et al., 2005; Sarkozy et al., 2005; Gutierrez-Roelens et al., 2006; Konig et al., 2006; Pabst et al., 2008). *NKX2.5* sequence variants have also been found in sporadic CHD, although these variants' contribution to the disease phenotype remains uncertain (Zhang et al., 2009). Here we made an attempt to investigate the role of c.239A>G SNP of *NKX2.5* in the manifestation of CHD.

## 2. Materials and Methods

A total of 150 CHD patients at K.R Hospital and J.S.S Hospital, Mysore (South India) were diagnosed by their past histories, physical examination and echocardiograms by experts. This study includes, 56 patients with VSD, 21 with ASD, 10 with ASD-VSD, 4 with TOF and 8 patients with Patent Ductus Arteriosus (PDA) and remaining with specific CHD phenotypes and complex forms. Informed consent was obtained from their parents or guardians. 70

individuals with the same ethnic background as well as no clinical history of CHD were used as controls. The research protocols and procedures were approved by the research ethics committee of both Hospitals and the ethics committee of University of Mysore.

EDTA tubes were used for venous blood collection. Genomic DNA was extracted from peripheral leukocytes by using Wizard genomic DNA purification kit (Promega). Both the exons of human *NKX2.5* (Gene Bank Accession Number: NM004387) were amplified through polymerase chain reaction (PCR) using exon specific primers (Benson et al., 1999). The PCR was performed in a 20 $\mu$ l reaction mixture containing 10x assay buffer, 60ng of genomic DNA, 0.2mM of each dNTP, 0.5 $\mu$ M of each primer and 1U taq DNA polymerase. The PCR condition was as follows: 95°C of initial denaturation for 3 minutes followed by 95°C for 30 seconds, annealing temperature of 68–72°C for 30 seconds, and extension at 72°C for 45 seconds for 35 cycles. A final extension was conducted at 72°C for 5 minutes. Further, the amplified products were purified using the Bangalore genei PCR purification kit (Cat # 117310) and sequenced for both the forward and reverse strands. The sequence analysis was done by using Accelrys gene software. The SNPs identified through these

investigations in patients and controls were reconfirmed through resequencing. Mass array analysis was carried out by using Sequenom-iPLEXR Gold SNP genotyping platform with Spectro CHIPR and MALDI-TOF Mass spectrometer (Sladek et al., 2007; Wilke et al., 2009).

### 3. Results

Figure 1 presents the organization of *NKX2.5*, which is 3.125 Kb with two exons that code for 324 amino acid residues having transcript length of 1.580 bp. It has homeodomain (HD), TN domain and NK2 specific domain (NK2-SD). Of the 85 CHD specific SNPs reported till date, only six SNPs, c.239A>G (48.66%), c.608A>G (0.6%), c.646C>T (25.33%), c.852G>A (3%), c.896C>A (0.6%) and 1212G>T (40%) were identified in the present study. Among these, 239 A>G is the most frequent (48.66%) SNP observed in the present study.

The SNP c.239A>G was observed in 73 cases out of 150 patients with CHD. Of which, 14 patients with cardiac septal defects (10 ASD, 27 VSD, 5 ASD with VSD) and 8 with PDA. In addition, this SNP was also observed in 36 out of 70 controls.




Fig.1 *NKX2.5* gene showing the most frequent SNP c.239A>G.

### 4. Discussion

Mutations in *NKX2.5* cause a spectrum of congenital heart defects, including ASD, VSD, and cardiac conduction abnormalities (Srivastava et al., 2000; Posch et al., 2008; Benson, 2010). Of the 85 SNPs, most of them found only in CHD cases and not in the control samples (Benson et al., 1999; Goldmuntz et al., 2001; McElhinney et al., 2003; Reamon-Buettner et al., 2004; Hirayama-Yamada et al., 2005; Bjornstad et al., 2009). Perusal of the

literature revealed that the c.239A>G SNP of *NKX2.5* is more frequent in both CHD and control samples (Table 1). However, in some cases, it was seen only in controls (Benson et al., 1999) and in some cases, it was observed in some CHD cases (Gutierrez-Roelens et al., 2002). The present study revealed that this SNP was also seen both in CHD cases and controls, which is a synonymous SNP coding for glutamine. Taking together of all these studies into consideration, the frequency of occurrence of this SNP in both CHD

cases (42.48%) and controls (41.36%) is almost same. The inconsistent reports of this SNP in some studies could be the error of sample size. One of the possibilities of occurrence of this SNP in more frequency in both CHD patients and controls could be due to its neutral effect, wherein although changes

seen at nucleotide level, at the amino acid level the change is synonymous and no effect in the coding property of glutamine. Therefore, this SNP become more common in the population and not under the selection pressure.

Table1. The available data on c.239A>G SNP of *NKX2.5*, screened in the congenital heart disease patients and controls.

Frequency of c.239A>G SNP		
CHD Cases	Controls	References
-	18/52 (36%)	Benson et al., 1999
0/49 (0%)	53/100 (53%)	Gutierrez-Roelens et al., 2002
55/68 (80%)	17/45 (38%)	Reamon-Buettner et al., 2004
5/12 (41.6%)	-	Khetyar et al., 2008
97/205 (47.3%)	-	Posch et al., 2008
17/28 (60.71%)	17/28 (60.71%)	Draus Jr et al., 2009
107/230 (46.5%)	98/200 (49%)	Zhang et al., 2009
12/99 (12.1%)	3/90 (3.3%)	Ding et al., 2009
11/168(6.5%)	-	Liu et al., 2009
140/208 (67.3%)	-	Hamanoue et al., 2009
73/150 (48.6%)	36/70 (51.4%)	The Present study, 2010
<b>517/1217 (42.48%)</b>	<b>242/585 (41.36%)</b>	

Thus, this SNP is seen in almost equal frequency in both CHD cases and controls as well as only in controls, one can state that this SNP is not involved in causing CHD.

### Conclusion

Since the change of SNP c.239A>G of *NKX2.5* is synonymous, the effect is neutral, thereby its frequency is almost equal in both controls and cases. Therefore, this SNP is not involved in the manifestation of CHD.

### Acknowledgments

We thank Council for Scientific and Industrial Research (CSIR), New Delhi, [No.27 (0156)/06/EMR-II dated 19.10.2006] for the financial support. We thank all the patients, families participated in this investigation, Doctors and PG students for their kind support, Investigators of Unit on Evolution and Genetics for the laboratory facilities and also Prof. H. A. Ranganath for his encouragement

**Corresponding Author:**

Prof. N. B. Ramachandra  
 Department of Studies in Zoology,  
 University of Mysore,  
 Manasagangotri, Mysore-570006, India.  
 Telephone: +91-8212419781 (O)  
 Fax: +91-8212516056  
 E-mail: [nallurbr@gmail.com](mailto:nallurbr@gmail.com)

**References**

1. Posch M G, Perrot A, Schmitt K, Mittelhaus S, Esenwein E M, Stiller B, Geier C, Dietz R, Gessner R, Ozcelik C, Berger F. Mutations in GATA4, NKX2.5, CRELD1, and BMP4 Are Infrequently Found in Patients with Congenital Cardiac Septal Defects. *A J Med Gen* 2008; 146:251–253.
2. Ramegowda S, Ramachandra NB. An understanding the genetic basis of congenital heart disease. *Ind J of Hu Gen* 2005; 11: 14-23.
3. Clark KL, Yutzey KE, Benson DW. Transcription factors and congenital heart defects. *Annu Rev Physiol* 2006; 68:97-121.
4. Chen MW, Pang YS, Guo Y. GATA4 Mutations in Chinese patients with congenital cardiac septal defects. *Pediatr Cardiol* 2010; 31:85-89.
5. Lyons I, Parsons LM, Hartley L, Li R, Andrews JE, Robb L, Harvey RP. Myogenic and morphogenetic defects in the heart tubes of murine embryos lacking the homeo box gene *Nkx2-5*. *Gene Dev* 1995; 9:1654-1666.
6. Schott J, Benson D W, Basson C T, Pease W, Silberbach G M, Moak J P, Maron B J, Seidman C E, Seidman J G. Congenital heart disease caused by mutations in the transcription factor *NKX2.5*. *Science* 1998; 281: 108-111.
7. Benson DW. Genetic origins of pediatric heart disease. *Pediatr Cardiol* 2010; 31:422-429.
8. Benson D W, Silberbach G M, Kavanaugh - McHugh A, Cottrill C, Zhang Y, Riggs S, Smalls O, Johnson M C, Watson M S, Seidman J G, Seidman C E, Plowden J, Kugler J D. Mutations in the cardiac transcription factor *NKX2.5* affect diverse cardiac developmental pathways. *J Clin Invest* 1999; 104: 1567-1573.
9. Hosoda T, Komuro I, Shiojima I, Hiroi Y, Harada M, Murakawa Y. Familial atrial septal defect and atrioventricular conduction disturbance associated with a point mutation in the cardiac homeobox gene *CSX/NKX2-5* in a Japanese patient. *Jpn. Circ. J* 1999; 63: 425–426.
10. Kasahara H, Lee B, Schott J, Benson DW, Seidman JG, Seidman CE, Izumo S. Loss of function and inhibitory effects of human *CSX/NKX2.5* homeoprotein mutations associated with congenital heart disease. *J Clin Invest* 2000; 106: 299-308.
11. Goldmuntz E, Geiger E, Benson DW. *NKX2.5* mutations in patients with tetralogy of Fallot. *Circ* 2001; 104: 2565-2568.
12. Ikeda Y, Hiroi Y, Hosoda T, Utsunomiya T, Matsuo S, Ito T, Inoue J, Sumiyoshi T, Takano H, Nagai R, Komuro I. Novel point mutation in the cardiac transcription factor *CSX/NKX2.5* associated with congenital heart disease. *Circ J* 2002; 66: 561 –563.
13. Watanabe Y, Benson D W, Yano S, Akagi T, Yoshino M, Murray J C. Two novel frameshift mutations in *NKX2.5* result in novel features including visceral inversus and sinus venosus type ASD. *J Med Genet* 2002; 39:807–811
14. Elliot D A, Kirk E P, Yeoh T, Chandar S, McKenzie F, Taylor P, Grossfeld P, Fatkin D, Jones O, Hayes P, Feneley M, Harvey RP. et al. Cardiac homeobox gene *NKX2-5* mutations and congenital heart disease. Associations with atrial septal defect and hypoplastic left heart syndrome. *J Am Coll Cardio* 2003; 41: 2072–2076.
15. McElhinney D B, Geiger E, Blinder J, Benson D W, Goldmuntz E. *NKX2.5* mutations in patients with congenital heart disease. *J Am Coll Cardiol* 2003; 42: 1650-1655.
16. Kasahara H, Benson DW. Biochemical analysis of eight *NKX2.5* homeodomain missense mutations causing atrioventricular block and cardiac anomalies. *Cardiovasc Res* 2004; 64:40-51.
17. Hirayama-Yamada K, Kamisago M, Akimoto K, Aotsuka H, Nakamura Y, Tomita H, Furutani M, Imamura S, Takao A, Nakazawa M, Matsuoka R. Phenotypes with *GATA4* or *NKX2.5* Mutations in Familial Atrial Septal Defect. *Am J Med Genet* 2005; 135:47–52.
18. Sarkozy A, Conti E, Neri C, D'Agostino R, Digilio MC, Esposito G, Toscano A, Marino B, Pizzuti A, Dallapiccola B. Spectrum of atrial septal defects associated with mutations of *NKX2.5* and *GATA4* transcription factors. *J Med Genet* 2005; 42:1-5.
19. Gutierrez-Roelens, De Roy L, Ovaert C, Sluysmans T, Deverieendt K, Brunner H G, Vikkula M. A novel *CSX/NKX2.5* mutation causes autosomal-dominant AV block: are atrial fibrillation and syncopes part of the phenotype? *Eur J Hum Genet* 2006; 14: 1313-1316.
20. Konig K, Will JC, Berger F, Müller D, Benson DW. Familial congenital heart disease, progressive atrioventricular block and the cardiac homeobox transcription factor gene

- NKX2.5: identification of a novel mutation. *Clin Res Cardiol* 2006; 95:499–503.
21. Pabst S, Wollnik B, Rohmann E, Hintz Y, Glänzer K, Vetter H, Nickenig G, Grohé C. A novel stop mutation truncating critical regions of the cardiac transcription factor NKX2-5 in a large family with autosomal dominant inherited congenital heart disease. *Clin Res Cardiol* 2008; 97:39–42.
  22. Zhang W, Li X, Shen A, Jiao W, Guan X, Li Z. Screening NKX2.5 Mutation in a sample of 230 Han Chinese Children with Congenital Heart Diseases. *Gen Test and mol Biom* 2009; 13:159-162.
  23. Sladek R, Rocheleau G, Rung J, Dina C, Shen L, Serre D, Boutin, Vincent D, Belisle A, Hadjadj S, Balkau B, Heude B, Charpentier G, Hudson TJ, Montpetit A, Pshzhetsky AV, Prentki M, Posner BI, Balding DJ, Meyre D, Polychronakos C, Froguel P. A genome-wide association study identifies novel risk loci for type 2 diabetes. *Nature* 2007; 445: 881-885.
  24. Wilke RA, Simpson RU, Mukesh BN, Bhupati SV, Dart RA, Ghebranious NR, McCarty CA. Genetic variation in CYP27B1 is associated with congestive heart failure in patient with hypertension. *Pharmacology* 2009; 10:1789-1797.
  25. Srivastava D, Olson EN. A genetic blueprint for cardiac development. *Nature* 2000; 407:221-226.
  26. Reamon-Buettner S M, Hecker H, Spanel-Borowski K, Craatz S, Kuenzel E, Borlak J. Novel NKX2–5 Mutations in Diseased Heart Tissues of Patients with Cardiac Malformations. *Am J Pathol* 2004; 164: 2117-2125.
  27. Bjørnstad P G, Leren T P. Familial atrial septal defect in the oval fossa with progressive prolongation of the atrioventricular conduction caused by mutations in the NKX2.5 gene. *Cardiol Young* 2009; 19:540-544.
  28. Gutierrez-Roelens I, Sluysmans T, Gewillig M, Deverieendt K, Vikkula M. Progressive AV-Block and anomalous venous return among cardiac anomalies associated with two novel missense mutations in the CSX/NKX2-5 gene. *Hum Mut* 2002; 512: 1-7.
  29. Khetyar M, Tinworth L, Syrris P, Abushaban L, Abdulazzaq Y, Silengo M, Carvalho J, Carter N. NKX2.5/NKX2.6 Mutations Are Not a Common Cause of Isolated Type 1 Truncus Arteriosus in a Small Cohort of Multiethnic Cases. *Genet Test* 2008; 12: 467-469.
  30. Draus Jr J M, Hauck M A, Goetsch M, Austin III E H, Tomita- Mitchell A, Mitchell M E. Investigation of somatic NKX2-5 mutations in congenital heart disease. *J Med Genet* 2009; 46: 115-122.
  31. Ding JD, Li KR, Zhang XL, Yao YY, Reng LQ, Tao SY, Fang X, Ma GS. Preliminary exploration of transcription factor Nkx2.5 mutations and congenital heart diseases. *Zhonghua Yi Xue Za Zhi* 2009; 16:1114-1116.
  32. Liu XY, Yang YQ, Yang Y, Lin XP, Chen YH. Mutations of NKX2-5 gene in patients with atrial septal defect. *Zhonghua Er Ke Za Zhi* 2009; 9:696-700.
  33. Hamanoue H, Rahayuningsih SE, Hirahara Y, Itoh J, Yokoyama U, Mizuguchi T, Saitsu H, Miyake N, Hirahara F, Matsumoto N. Genetic screening of 104 patients with congenitally malformed hearts revealed a fresh mutation of GATA4 in those with atrial septal defects. *Card in the Y* 2009; 19:482-485.

**25/05/2010**