

Biochemical and Microbiological Evaluation of Fermented Camel Milk

Abeer A. Khayal¹ and Ragia O.M.²

¹Chemistry Department, Faculty of Science and Arts, Al Jouf University, Kingdom of Saudi Arabia

²Food Technol. Res. Inst. Agriculture Research Centre, Egypt

E-mail: abeerkhaial@yahoo.com

Abstract: Camel milk has an important role in human nutrition in the hot regions and arid countries. The general composition of camel milk varies in various part of the world with range of 3.07-5.50% fat, 3.5-4.5% protein, 0.7-0.95% ash and 3.4-5.6 % lactose, 12.1-15% total solid. Camel milk is different from other ruminant milk, having low cholesterol, low sugar, high minerals (sodium, potassium, iron, copper, zinc and magnesium), high vitamin C. Camel milk contains more proteins and whey protein than cow milk. Camel milk is remedy for viruses causing diarrhea as Rota Virus considering it could be important from public health point of view to anticipate the vital hazard in camel raw milk and its produces. In this study, we used rat model of rotavirus infection which causing diarrhea. This investigation was designed to prepare fermented Camel milk with low fat (1.5%) and using it in formulas for feeding diarrhea rats (25%, 50% and 75%) to perform the role of the macro elements (sodium, potassium, phosphorous and calcium) as well as detection of *Staphylococcus aureus* of the three prepared formulas for diarrhea rats. Results indicated that the fermented camel milk had higher content in sodium and potassium than the row camel milk and there was stopping diarrhea on formula (3), this is due to the formula (3) can be considered antimicrobial and a strong vital inhibitor to human Rota Virus which represents the main diarrhea-causing agent in infants. It can be concluded that fermented camel milk can be considered as a good food of high nutritive and therapeutic applications. Meanwhile, the high content of antimicrobial agents in camel milk may explain its potential as an activity especially against diarrhea.

[Abeer A. Khayal and Ragia O.M. **Biochemical and Microbiological Evaluation of Fermented Camel Milk.** *N Y Sci J* 2013;6(9):74-79]. (ISSN: 1554-0200). <http://www.sciencepub.net/newyork>. 11

Key words: Camel milk- diarrhea-antimicrobial- minerals

Introduction

Camel's milk in particular is a good source of various vitamins and minerals and is characterized for its low cholesterol and high concentration of insulin (Agrawal et al., 2005) & (Badriah., 2012). Camel, which is also known as ship of desert, is used for transportation and a source of milk, meat and wool (Meiloud et al., 2011). It has medicinal properties and antibacterial and antiviral activity (El-Agamy et al., 1992). which may be due to higher concentration of lactoferrin in camel's milk (Yagil, 1994). Camel milk is used therapeutically against hepatitis, dropsy, problems of spleen, asthma, (Mal et al., 2000). Camel milk is also having low cholesterol, low sugar, high minerals (sodium, potassium, iron, copper, zinc and magnesium), high vitamin C, low protein and large concentration of insulin (Agrawal et al., 2004) & (Arrowal.,2005). There are no allergens, and it can be consumed by lactase deficient persons and those with weak immune systems. The milk is considered as having medicinal properties. In Sahara, fresh butter is not eaten, but is often used as base for medicines. The products developed also include cosmetics or pharmaceuticals. It claimed that the value of camel milk is to be found in the high concentrations of volatile acids especially, linoleic acid and polyunsaturated acids, which are essential for human

nutrition (Agrawal et al., 2004). A series of metabolic and autoimmune diseases are successfully being treated with camel milk. In India, camel milk is used therapeutically against dropsy, jaundice, problems of spleen, asthma, anemia, piles and diabetes role of raw camel milk in chronic pulmonary tuberculosis patients has been observed (Mal et al., 2000 & Agrawal et al., 2004).

Milk plays a significant role in human's nutrition for conducted to test the effect of camel's milk in comparison the wonderful reason that they are excellent source of cow's milk on blood glucose level, triglycerides, various nutrients. Milk diet has been suggested in the cholesterol and ALT and AST levels in healthy albino management of various diseases (Badriah., 2012).

In several parts of world, camel milk is used as a remedy for some diseases as tuberculosis, juvenile diabetes (Beg et al., 1985), liver cirrhosis, rickets, constipation, asthma (Yagil, 1987). Camel milk was not only contains more nutrients compared to cow milk, but also it has therapeutic and antimicrobial agents (Barbour et al, 1984 & El-Agamy et al., 1992). The antiviral activity in camel milk was also found (El-Agamy et al., 1992). Camel milk is remedy for viruses causing diarrhea as Rota Virus considering it could be important from public health point of view to anticipate

the vital hazard in camel raw milk and its produces. Therefore, Camel milk can be considered a strong vital inhibitor to human Rota Virus which represents the main diarrhea-causing agent in infants (El-Mougi, 1999).

In the present study, we prepared fermented camel milk with low fat (1.5%) in formulas for feeding diarrhea rats and perform the role of the macro elements (sodium, potassium, phosphorous and calcium) as well as detection of *Staphylococcus aureus* of the three prepared formulas for diarrhea rats.

Material & Methods

I- Materials:

(A). Camel milk used in the current study obtained from local market at GIZA, Egypt freeze dried direct vet set (FD-DVS) MYE 96 yoghurt culture was obtained from Rodia Food France. The culture was intended for direct inoculation to the process milk at 0.1%.

(B). Animals: Twenty four Sraye-Dawley White Albino rats, of an average weight 90 ± 0.2 gm were used in this study. The rats were obtained from experimental animal house in Food Technology – Research Institute, Agriculture Research Centre, Giza - Egypt.

Animal inoculation: Rats were inoculated by oral gavage with 0.5 ml phosphate-buffered saline (PBS), 6.75×10^8 PFU of RRV, or 1.45×10^8 PFU of HAL1166 rotavirus. Control animals were always handled prior to virus-inoculated animals. To determine the serum 50% antigenemia dose (AD50) and the 50% diarrhea dose (DD50) of RRV, rats were inoculated with 0.5 ml of serial 10-fold dilutions of RRV. For a subset of experiments, the rats were examined daily for evidence of rotavirus-induced diarrhea by gentle abdominal palpation. Diarrhea was noted and scored from 0 to 4 based on stool color, amount, and consistency. A score of 2 or greater was considered diarrhea (Ciarlet, 2002).

II-Methods:

Biochemical analysis

Milk sample were analyzed for Moisture, pH, titratable acidity, specific gravity, total solids, and fat according to Ling (1963).

Determination of minerals:

Calcium, Sodium, Phosphorus, Potassium, Iron, Zinc concentration were determined by atomic absorption (Thermo-Tarrell, Ash, Smith-Hieftje (1000) in their digested solutions according to A.O.A.C (2000).

Protein determination:Protein concentration was measured spectrophotometrically according to AOAC (1980).

Prepared fermented Camel milk

Fermented Camel milk was prepared with low fat (1.5%). Camel milk was heated after decreased fat to 1.5% at 90°C for 10 min, immediately cooled to 42°C and inoculated with 0.1% DVS yoghurt culture. The inoculated milk was dispensed into plastic cups, fitted with press on lids. Cups were incubated at 42°C until titratable acidity of 0.8% was reached. Control diet (Casein-based diet) was composed of 11.6% casein (equal 10% protein), 5% corn oil, 4% mineral mixture, 1%vitamin mixture, 5% cellulose and corn starch up to 100%.

Microbiological analysis

Fermented camel milk samples (11 ml) were homogenized for one minute in 99 ml (1/10) of a sterile solution of 0.1% (w/v) peptone water (Oxoid CM9) using a Stomacher Lab blender (Model400, Seward Laboratory, London). From these samples serial decimal dilutions were prepared in sterile 0.1% peptone water. The microorganism's counts were carried out by the pour-plate method with duplicate plating on selective agar media (Parrow, 1978). The coliforms were estimated in duplicate pour plates of Violet Red Bile Agar (VRBA, Oxoid CM107) medium and the plates were overlaid after solidification with 3 to 4 ml of additional Violet Red Bile Agar. The plates were incubated in an inverted position at $30^\circ\text{C} \pm 2$ for 18 – 24 h (Mehlman, 1984). The yeasts and molds were counted on acidified Potato Dextrose Agar, (Oxoid CM139) which was acidified by the addition of the proper amount of sterile 10% tartaric acid (Fluka-AG-Buchs.SG), then the plates were incubated at $25^\circ\text{C} \pm 1$ for 3 - 7 days (Koburger and Marth, 1984). The lactic acid bacteria were enumerated in pour plates of de Man, Rogosa and Sharp (MRS) medium (Oxoid CM359). The plates were incubated at 37°C for 48 h under microaerobic conditions using Gas Pak (H₂+CO₂) (BBL, Microbiology Systems, Div. Becton Dickinson and Co., Cockeysville. Med.) anaerobic systems (Gilliland et al., 1984) . The detection of *Staphylococcus aureus*, was applied according to the methods described in the FDA (1998) .

Experimental diets:

The composition of experimental diets used in this work for feeding diarrhea rats is shown in table (1):

Table (1): The experimental rat groups and their diets

Items	Basel diet	% additional		
		25%	50%	75%
Group 1 (control)	100	-	-	-
Group 2	75	25	-	-
Group 3	50	-	50	-
Group 4	25	-	-	75

Biological experiments

Twenty four rats were fed the control diet for 7 consecutive days, rats were infected by rotavirus to induce diarrhea. After that, the rats were divided into four groups; each of it consists of six rats. The first group was the control one that fed on Casein-based diet. The rats of other groups (groups 2, 3, and 4) were fed on fermented camel milk. The total feeding period was 28 days. The rats were weighted after the 7 and 14 days at the beginning of the experiment and once a week. The gain weight was calculated by differences between the final body weight (gm) and the initial body weight (gm). The protein efficiency ratio (PER) was calculated according to A.O.A.C (2000) as follow:

$$\text{PER} = \frac{\text{Gain in body weight (gm)}}{\text{Protein consumed (gm)}}$$

The corrected PER was calculated according to Jansen et al., 1978 as follow:

$$\text{Correct PER} = \text{PER} \times \frac{2.5}{\text{PER for stander casein}}$$

Blood Sampling:

In all previously mentioned animal groups, blood samples were collected after 12 hours fasting at the start of the experiment and after 2 and 4 weeks, from orbital venous by means of micro capillary glass tubes. The blood of the six rats of each group were placed in a dry clean centrifuge tubes and left to clot in a water bath (37°C) for half an hour. The blood was centrifuged for 10 minutes at 4000 r.p.m to separate the

serum. Serum was carefully aspirated and transferred into clean curvet quartz tubes and kept frozen at -20°C till analysis.

Statistical analysis

Statistical analysis was performed using the SPSS software package for Windows [SPSS (UK) Ltd., Surrey, United Kingdom]. ANOVA was used to determine the difference between the means of the groups. Further analysis was carried out using a t-test for comparing two variables. P value considered significant when it was < 0.05

RESULTS AND DISCUSSION

Camel milk used in this study has Total Solids (T.S), Fat, Total Nitrogen (T.N) and Titratable Acidity (T.A), with contents of 11.92%, 3.1%, 0.49% and 0.18% respectively. This data is similar with Badran.,2004, While fermented camel milk was analysis T.S, Fat, P.N, N.P.N and acidity were 12.3%, 1.5%, 0.432%, 0.054 and 0.88% respectively. In this study we used separation fat partly from camel milk to (1.5%) before manufacture of fermented camel milk. we prepared quantity different formulas from it (25%, 50% and 75%) as shown as in table (1). Three prepared formulas from fermented camel milk used for therapeutics diarrhea.

The macro elements (sodium, potassium, phosphorous and calcium) as well as micro elements (iron and zinc) of the three prepared formulas for diarrhea rats were determined and the results were presented in table (2):

Table (2): Minerals content of formulas for diarrhea rats (mg/100gm)

Formulas	macro elements				micro elements	
	Na	K	P	Ca	Fe	Zn
Formula 1	70	180	113	135	0.02	0.50
Formula 2	75	185	115	140	0.10	0.70
Formula 3	80	195.5	120	145	0.15	1.20

Minerals content noticed in fermented camel milk were distinguished by the higher content of the sodium and potassium with the values 70.0 and 180.0 mg/100g than the row camel milk which recorded 52.0 and 145.0 mg/100g for sodium and potassium, respectively. These obtained results were agree with these obtained by (Watt and Merrill, 1963), (Abu-Lehia, 1989)],

(Aderson, 1991), (El-Agamy et al., 1998) and (Shamsia, 2009).

The data indicated that formula (3) characterized by the highest levels of Na, K, P and Ca, they were 80.0 and 195.5, 120, and 145 mg/100g, respectively. While formula 1 was the least in its content of these elements, with values of 70.0, 180.0, 113, and 135 mg/100g

respectively. Also, the values of microelements (Iron and Zinc) in formula 3 were 0.15 and 1.2 mg/100gm respectively compared to formula 1, 2. In general, it was clearly noticed that most formulas have nearly the same content of the studied minerals.

Sodium deficiency was seen when severe diarrhea or vomiting occur. Partially, all the body sodium was found in the extra-cellular fluid which bathes the tissues. It does not cross the cell membrane to any great extent. Sodium functions are important in the regulation of acid-base equilibrium maintenance of osmotic pressure and of water balance. Potassium deficiency occur when there was prolonged failure treat and in pathologic conditions such as severe diarrhea (Robinson and Lowler, 1989).

The results of protein efficiency (PER), protein, sodium and potassium contents in serum of diarrhea rats are showed in table (3) and fig (1):




Table (3): Means of protein efficiency (PER), value of protein, sodium and potassium contents in serum of diarrhea rats fed on control and experimental diets

Groups	Serum parameters after periods in days											
	PER			Protein (mg/l)			Sodium (m mol/l)			Potassium(m mol/l)		
	7	14	21	0	7	21	0	7	21	0	7	21
Control	2.8	2.9	3.0	6.5	6.9	7.2	150.0	151.9	151.5	5.6	5.7	5.6
Group 1	2.4	2.5	2.5	6.8	7.0	7.2	150.2	151.3	151.2	5.5	5.5	5.6
Group 2	2.6	2.7	2.7	6.9	7.1	7.2	150.7	151.5	151.4	5.5	5.6	5.6
Group 3	2.8	2.9	3.0	7.0	7.2	7.4	151.0	151.8	151.6	5.7	5.7	5.6
L.S.D	0.2	0.3	0.1	0.5	0.8	0.9	0.6	0.7	0.8	0.6	0.8	0.9

L.S.D: (Less Significance Difference at = 0.05)

From table (3), we noticed that all prepared diets were of good PER with ranged between (2.4 to 3.0). The highest PER was detected for rats fed on the diet prepared from formula 3; it was the same with rats fed on the control casein diet. Also, this table shows the means of total protein, sodium and potassium contents in the serum of diarrhea rats. These parameters were significantly equal for all rats at all the detected periods. Serum protein levels of rats fed control and tested diets were higher than the normal range this is in agreement with (El-Agamy et al., 1998) who found higher fat , protein (especially casein) and ash contents but lower whey protein and lactose contents in camel milk, Also with (Shamsia., 2009) who revealed that camel milk proteins contained satisfactory balance of essential amino acids. So, camel milk can be considered as good source of protein and can meet part of the daily needs of humans from these nutrients.

Fermented milk products are known for their taste, nutritive value and therapeutic properties. According to the International Dairy Federation (1969), fermented milks are “products prepared from milk” whole or fully skimmed, concentrated or milk substituted from partially or full skimmed milk, either homogenized or


un-homogenized, pasteurized or sterilized and fermented by mean of specific microorganisms. Milk from camel has been used to make traditional fermented milk products throughout the world. The people who had domesticated these milk animal usual accepted fermented milk by necessity (Kroger et al., 1989), and (Abdel Moneim et al., 2009).

As shown in table (4), the infectious of rats produced by a Rotavirus to induce diarrhea caused a great loss in the initial body weight of all rats. When the diarrhea rats were fed on formula 3, diarrhea was stopped at the second day compared to formula 1, 2 and control casein diet. Formula (2) was followed by formula 3 in stopping diarrhea after three days which contained 50% fermented camel milk, 50% control diet, while formula (1) was stopping diarrhea after four days, its similar with the control casein diet which contained 25% fermented camel milk. These observations could be considered that formula (3) was the adequate one for stopping diarrhea compared to other prepared formula and the control, followed by formula (2) and formula (1) as shown in fig (2).

Table (4): Means of body weigh (B.W), gain weight (G.W) and diarrhea stopping day of diarrhea rats fed on control and experimental diets

Groups	Initial body weigh (g)	B.W and G.W (g) after periods in days								Diarrhea stopping day
		4 days		7 days		14 days		21 days		
		B.W.	W. loss	B.W.	G.W.	B.W.	G.W.	B.W.	G.W.	
Group1 (control)	90.2	80.2	10.0	92.5	12.3	110.5	30.3	127.0	46.8	4
Group 2	85.0	70.0	15.0	88.0	18.0	100.0	30.0	130.7	60.7	4
Group 3	88.2	75.0	13.2	89.4	14.4	105.0	30.0	135.0	60.0	3
Group 4	90.1	80.0	10.1	91.4	11.4	111.0	30.0	136.0	56.0	2
L.S.D	0.9	0.8	0.4	1.0	0.5	0.4	0.9	0.6	1.0	-

After infectious of rats which produced by a Rotavirus to induce diarrhea *



Results were agreement with previously reported (Abu-Lehia, 1987, Aderson, 1991 and El-Agamy et al., 1998).

The results of this work showed that final products of fermented camel milk prepared in the lab by using cultures had no *Staphylococcus aureus* in formula(3).

The stopping diarrhea at the second day on feeding formulas (3) may be attributed to its higher content of fermented camel milk (75%), besides this formula contained the highest level of protein, Na and K,. The stopping diarrhea may be due to the interaction effect between all these factors in formula (3) causing diarrhea.

The concentrations of macro elements (Na, K, P, and Ca) were higher in formula 3 than in formula 1 and 2. Therefore, it considered that the formula 3 as a good source of these minerals. Also, formula 3 contained higher values of microelements (Fe and Zn).

Besides, the fermented camel milk can stopping diarrhea because it contained specific antibodies to Rota virus which cause diarrhea and in addition, the fermented camel had antiviral activity against human Rotavirus, so, it used as a remedy for virus (Shamsia, 2009)].

It can be concluded that fermented camel milk can be considered as a good food of high nutritive and therapeutic applications. Meanwhile, the high content of antimicrobial agents in camel milk may explain its potential as an activity especially against diarrhea.

Corresponding Author:

Dr. Abeer A.Khayal
Department of Chemistry
Al Jouf University
Kingdom of Saudi Arabia
E-mail: abeerkhaial@yahoo.com

References:

1. Abdel Moneim E. Sulieman, Hamid, A.Dirar, Elamin A. Elkhalifa and Ali O.Ali.(2009): "Effect of fermentation on some chemical components of the Sudanese fermented butter milk" J. Sc. Tech Vol.10(1): 1-10
2. Abu-Lehia, (1987): "Composition of camel milk". *Milchwissenschaft*, 42:368-371
3. Abou-Lehia J., Al-Mohizea, L.S. and Behiry M. (1989): "Studies on the production of ice-cream from camel milk products". *Australian G. Dairy. Tech. Vol (44): 32-34*
4. Aderson R.R. (1991): Comparison of minerals in milk of four species. *Comparative biochemistry & Physiology. A Comparative physiology*, 100 (4): 1045-1048
5. Agrawal, R.P., R. Beniwal, D.K. Kochar, F.C. Tuteja, S.K. Ghorui, M.S. Sahani and S. Sharma, 2005. Came IMilk as an Adjunct to Insulin Therapy Improves Long-Term Glycaemia Control and Reduction in Doses of Insulin in Patients with Type-1 Diabetes A 1 Year Randomized Controlled Trial. *Diabetes Research and Clinical Practice*, 68: 176-177.
6. Agrawal R.P., Kochar D.K., Sahani M.S., Tuteja F.C. and Ghru S.K. (2004): "Hypoglycaemic activity of camel milk in streptozotocin induced diabetic rats" *Int. J. Diab. Dev. Countries Vol. 24:47-49*.
7. AOAC, (1980): *International Official Methods of Analysis*, 13th Ed. Association of official Analytical Chemists, Washington DC.
8. AOAC, (1998): *Official Methods of Analysis*, 15th Ed. Association of official agricultural chemists, Washington, DC.
9. AOAC, (2000): *Association of official Analytical chemists. Official Methods of Analysis*, 17th Ed A0AC, Washington, DC.
10. Arrowal, R.P., Beniwal, R. Kochar, D.K., Tuteja, F.C., Ghorui, S.K., Sahani, M.S., Sharma, S. (2005): *Camel milk as an adjunct to insulin*

- therapy improves long-term glycaemia control and reduction in doses of insulin in patients with type-1 diabetes A 1 year randomized controlled trial. – *Diabetes Res. Clin. Pract.* 68:17
11. Badran M.S., Sherein A.L., Abou Dawood and El-Assar M.A. (2004): Evaluation of fermented camel milk prepared using yoghurt starter. Proc.9 th Egyptian Conf. for Dairy Sci & Tech.
 12. Badriah A.(2012): Effect of Camel Milk on Blood Glucose, Cholesterol, Triglyceride and Liver Enzymes Activities in Female Albino Rats *World Applied Sciences Journal* 17 (11): 1394-1397.
 13. Barbour, E.K., Nabbut, N.H., Ferisch, W.M., Ai-Nakhli, H.M. (1984): Inhibition of pathogenic bacteria by camel's milk: relation to whey lysozyme and stage of lactation. *J. Food Protec.* 47: 838-840.25.
 14. Beg, O.U, H. Von Bahr-Lind Strom, Z.H. Zaidi, H. Tornvall. (1986): A camel milk protein rich in half cystine, primary structure assessment of variations, internal repeat patterns and relationship with neurophysin and other active polypeptides. *Eur. J.Biochem*, Aug., 15(1): 195-201.
 15. Beg O.U., von-Bahr-Lind strom H, Zaidi Z.H., Jorvall H. (1985):“Characterisation of camel milk protein rich praline identifies a new beta casein fragment. *Regul Pept* vol. 15: 55-62.
 16. Ciarlet, M., M. E. Conner, M. J. Finegold, and M. K. Estes. (2002). Group A rotavirus infection and age-dependent diarrheal disease in rats: a new animal model to study the pathophysiology of rotavirus infection. *J. Virol.* 76:41-57.[
 17. El-Agamy, E.I., Abou-shloue Z.I, Abdel-kader Y.I (1998): Gel electrophoresis of proteins, physicochemical characterization and vitamin C content of milk of different species. *Alexandria J. Agric. Res.* 43(2): 57-70
 18. El-Agamy, E.I., Ruppner, R., Ismail, A., Champagne, C.P., Assaf, R. (1992): Antimicrobial and antiviral activity of camel milk protective proteins. – *J. Dairy Res.* 59: 169-175.
 19. El-Mougi M.T. (1999): Basic pediatrics. Faculty of Medicine, Al- Azher Univ., Cairo, Egypt: 194-213.
 20. FDA (1998). *Bacteriological Analytical Manual*. 8th ed. Revision A, Chapter. 4, 5, 10, 12 and 14. US. Food and Drug Administration Center for Food Safety and Applied Nutrition. Gaithersburg: AOAC.International.
 21. Gilliland SE, Sadine WE, Vendamuthu ER (1984). Acid-producing microorganisms. In. Speck ML (ed.), *Compendium of Methods for the Microbiological Examination of Foods*. American Public Health Association. Washington, D.C. pp.184-194
 22. Hanna J. (2001): Over the hump. In: Jack Hanna's Animal Adventures. TV series (USA).
 23. International Dairy Federation (IDF) (1969): *Composition and standards for fermented milks*. FIL-IDF, (47-52).
 24. Koburger JA, Marth EH (1984). *Yeasts and Molds*. In: Speck ML (Ed), *Compendium of Methods for the Microbiological Examination of Foods*. American Public Health Association, Washington. pp. 197- 201.
 25. Kroger, M., Kurman, J. A. & Rasic, L. (1989): Fermented milks- past, present and future. *Food Technology*, 43(1):92-99
 26. Ling E.R (1963): *A text book of Dairy Chemistry*. Vol.2, 3 rd ED. Chapman & Hall, London
 27. Mal G., Suchitra Sena D., Jain V.K., Singhvi N.M., Sahani M.S., (2000).: Therapeutic utility of camel milk as an adjuvant nutritional supplement against multiple drug resistant patient. Proc. Int. Camelid Conf., Almaty, Kazakhstan, p.99
 28. Mehlman IJ (1984). Coliforms, fecal coliforms, Escherichia coli and enteropathogenic E.coli. In: Speck ML(ed), *Compendium of Methods for the Microbiological Examination of Foods*. American Public Health Association Washington, D.C. P. 265-285.
 29. Meiloud, G.M., I.N. Ould Bouraya, A. Samb and A. Houmeida, 2011. *Composition of Mauritanian Camel Milk: Results of First Study*. *International Journal of Agriculture Biology*, 13: 145-147.
 30. Parrow T (1978). *Manual of Recommended Laboratory Methods for Sampling and Microbiological Testing of Single Service Food Packages and their Components*. Syracuse, NY: Syracuse University Res. Corp.
 31. Robinson C.H. and Lowler (1989): “Normal and therapeutic nutrition 15th ED. Published by Macmillan publishing Company, INC. New York pp.127-128
 32. Shamsia S.M. (2009): “Nutritional and therapeutic properties of camel and human milk” *International J. of Genetics and molecular Biology* vol.1 (2), pp.052-058
 33. Watt B.K and Merrill A.L. (1963): *Composition of foods No. 8 consumer and food economics*. Research Division. Agric. Research. Services. United State Department of Agric. Washington, D.C.
 34. Yagil R. (1982): *Camels and camel milk*. Italy: FAO (Food and agricultural Organization of the UN).
 35. Yagil R (1987): Camel milk –A review. *Int. J. Anim. Sci.* Vol (2): 81-89.