

Phytochemicals and hypertension

¹Akram kooshki, ²Mahmood Rivandi

¹. Asistan professor, Traditional and Complementary Medicine Research Center, Sabzevar University of Medical Sciences, Sabzevar, Iran

² Master science in Organic Chemistry, Bureau of Education, Sabzevar, Iran

[E-mail:akooshki.nutr@yahoo.com](mailto:akooshki.nutr@yahoo.com)

Abstract: Hypertension is above the systolic blood pressure above 140 mmHg or diastolic blood pressure greater than 90mmHg or both. It is that can be controlled by diet containing an adequate supply of phytochemicals. The article is based on non-systematic literature searches in PubMed, Ovid and Sciencedirect and ISC. Hypertension is one of the major chronic diseases that can affect people around the world. Today's DASH diet (dietary approach to stop hypertension) is used to reduce and control blood pressure. The diet containing plenty of fruits, vegetables, nuts and whole grains that also contain nutrients such as vitamins and minerals and non- nutrient elements are called phytochemical. Phytochemicals are materials at very low concentrations, effect on the expression of many genes, including transcription factors AP-1, NF-Kappa B, SREBPs, PPARs and their cellular metabolism and gene regulation and they have antioxidant properties and anti-hypertensive effects. Moreover, increased intake of phytochemicals is associated to reduction of death from disease cardiovascular diseases, stroke and damage secondary to hypertension. Possible mechanisms of action of phytochemicals in reducing blood pressure are including of inhibition of angiotensin-converting enzyme and inactivation bradikinin and regulation of bioavailability of nitrous oxide. In conclusions based on the results, increased intake of phytochemicals in daily diet is associated with the opposite of death from heart disease - coronary, stroke and damage secondary to hypertension. [Kooshki A, Rivandi M. **Phytochemicals and hypertension**. *Biomedicine and Nursing* 2016;2(1): 67-70]. ISSN 2379-8211 (print); ISSN 2379-8203 (online). <http://www.nbmedicine.org>. 11. doi:[10.7537/marsbnj02011611](https://doi.org/10.7537/marsbnj02011611)

Key words: hypertension, phytochemicals, plant foods, poly phenols, flavonoids, carotenoids

Introduction:

Hypertension or high blood pressure defined as systolic blood pressure of greater than 140 mmHg or diastolic blood pressure greater than 90mmHg or both, affected 11% people in the Iran(1). Hypertension is the third risk factor for death from cardiovascular disease, with LDL-C and smoking has been called the silent killer(2,3).

Also, hypertension is the major risk factor for stroke and renal dysfunction. Lowering the blood pressure, through dietary may decrease damage of hypertension.

Epidemiological studies suggest that consumption of fruit, vegetables (4,5,6,7,8) and mineral water, water (9,10) and tea may protect against high blood pressure (11,12,13,14).Vegetables and fruits and tea may decrease blood pressure due to phytochemicals(15,16).

Functional non- nutrient food factors (phtochemicals) are in vegetable and fruits that considered to be effective for health promotion and disease prevention (15,17)

Methods:

The article is based on non-systematic literature searches in PubMed, Ovid and Science direct and ISC.

Phytochemical:

Flavonoids, polyphenols and carotenoids are included in phytochemical database (15).

This review summarizes current evidence on the effect of phytochemical on blood pressure.

High intake of fresh fruit and vegetable was associated with a low stroke mortality, an effect that was attributed to preventing hypertension(18).

Flavonoids and hypertension:

Flavonoid compounds included flavonols (quercetin, kaempferol, myricetin), flavones (apigenin, lutein) and isoflavones (coumestrol, daidzein, genistein, equol) (18).

Therefore, vegetarians had approximately 40% higher intake of flavonoids than non – vegetarians. Flavonoids are scavengers of free radicals such as superoxide anions and lipid proxy radicals and by preventing oxidation of LDL-c, because oxidized LDL is believed to be atherogenic. Therefore, flavonoids may decrease the formation of atherosclerotic plaques and reduce arterial stiffness, Leaving arteries more responsive to the endogenous stimuli of vasodilation (18,5).

The major dietary of flavonoid intake is black tea studies have shown that consumption of fruit, vegetable and tea (black and green) containing high amount of flavonoid may lead to lower blood pressure

and may provide protection against coronary heart disease and stroke. It is possible that their antioxidant effect are responsible.

Poly phenols and hypertension:

The Mediterranean diet has been associated with greater longevity and quality of life in epidemiological studies. Because of the abundance of fruits and vegetable are provided high amounts of polyphenols. Several polyphenol- rich sources such as grape-derived products, cocoa, tea, soy, pomegranate and tomatoes have been shown to decrease blood pressure in hypertensive patients (6,19,20,21,22,23,24).

Polyphenol compounds included chlorogenic acid, caffeic acid, coumaric acid, gallic acid, quercetin, isorhammetin, kaempferol, hesperetin, narigenin, phloretin, enterolacton and enterodiol (25).

Recent studies suggest that chlorogenic acids which are the main components of the polyphenol class in coffee, decrease blood pressure (26). The hypotensive effects of chlorogenic acids was associated with nitric oxide, the suppression of mRNA expression of NADPH oxidase, and the improvement in endothelium-dependent vasodilation in the aorta (27,28).

Grassi and etal suggest that consumption of dark chocolate for 15 day reduce systolic blood pressure in healthy subjects as well as young and elderly hypertensive patients (29).

Polyphenols decrease blood pressure by increasing in endothelial nitric oxide bioavailability via their antioxidant action and their capacity to activate vascular endothelial nitric oxide synthase.

Carotenoids and hypertension:

In the last years, It is known the excess of reactive oxygen species(ROS), is associated to inflammation, growth and vasoconstriction contributing to vascular injury in many cardiovascular disease such as hypertension, hyperlipidemia and diabetes (30,31). ROS play an important pathophysiological role in hypertension.

Antioxidant such as carotenoids inhibit NAD(P)H oxides- driven generation of ROS reduces, improvement of endothelial function, enhance nitric oxide (NO) generation and reduce blood pressure in hypertensive patients (30). Astaxanthin, an oxycarotenoid has antihypertensive effect by normalization of sensitivity of adreno-receptor sympathetic pathway, decrease in oxidative stress and improvement in NO bioavailability (30).

Lycopene, a carotenoid without pro-vitamin A, is red pigment in tomatoes and watermelon. High lycopene consumption has been associated with a decrease blood pressure (32,33). Meta analysis of the effect of lycopene on systolic blood pressure of all

trials is suggested a significant blood pressure reducing effect (-5.60 + 5.26 mmHg) but no significant effect on diastolic blood pressure (34).

Blood pressure lowering properties of lycopene have been attributed to the stimulation of nitric oxide production in the endothelium(34).

In conclusions based on the results, increased intake of phytochemical in daily diet is associated with the opposite of death from heart disease - coronary, stroke and damage secondary to hypertension. Therefore, it is emphasized to consume foods rich in phytochemical such as vegetables, fruits, nuts, tea and coffee, etc, in the diet order to control and prevention of hypertension.

Acknowledgement:

The present study was supported by Sabzevar university of Medical Sciences (No:11577_ xxx).

Corresponding Author

Akram kooshki, Traditional and Complementary Medicine Research Center, Sabzevar University of Medical Sciences, Sabzevar, Iran
E-mail: akooshki.nutr@yahoo.com

References:

1. Ranjbar F, Akbarzade F, Khaneshi M. The study of type A personality in essential hypertensive patients. *Medical Journal of Kordestan University of Medical Sciences* 2006; 11: 26-31.
2. Kooshki A, Yaghoobifar M A, Behnam Vasshani H R. Relation between drinking water and blood pressure. *Sabzevar J Medical Science* 2004;10(3):23-28.
3. Kooshki A, Mohajeri N, Movahedi A. Prevalence of CVD Risk Factors related to diet in Patients Referring to Modarres Hospital in Tehran in 1999. *Sabzevar J Medical Science* 2004;10(2):17-22.
4. Liu S, Manson JE, Lee IM, Cole SR, Hennekens CH, Willett WC, Buring JE. Fruit and vegetable intake and risk of cardiovascular disease: the Women's Health Study. *Am J Clin Nutr* 2000 Oct;72(4):922-8.
5. de Paula TP, Steemburgo T, de Almeida JC, Dall'alba V, Gross JL, de Azevedo MJ. The role of Dietary Approaches to Stop Hypertension (DASH) diet food groups in blood pressure in type 2 diabetes. *Br J Nutr* 2011; Dec 6:1-8.
6. Alonso A, de la Fuente C, Martín-Arnau AM, de Irala J, Martínez JA, Martínez-González MA. Fruit and vegetable consumption is inversely associated with blood pressure in a Mediterranean population with a high vegetable-fat intake: the Seguimiento Universidad de Navarra (SUN) Study. *Br J Nutr* 2004

- Aug;92(2):311-9.
7. Wang L, Manson JE, Gaziano JM, Buring JE, Sesso HD. Fruit and Vegetable Intake and the Risk of Hypertension in Middle-Aged and Older Women. *Am J Hyperten* 2011 Oct; 13.
 8. doi: 10.1038/ajh.2011.186. [Epub ahead of print]
 9. Kooshki A, Golafrooz Shahri M. Assessment of macro and micro nutrients on blood pressure in diabetics patients. *Ofogh Danesh* 2004;11(1):34-37.
 10. Kooshki A, Aminpour A, Azar M, Kalantari N, Ardakani H. The effect of drinking damavand mineral water on sodium, potassium, magnesium and calcium cations in men with hypertension. *Sabzevar J Medical Science* 2003;9(4):31-38.
 11. Kooshki A, Aminpour A, Azar M. Influence of type and duration of water intake on blood pressure in hypertensive men. *Ofogh Danesh* 2007;12(2):50-54.
 12. Yang YC, Lu FH, Wu JS, Wu CH, Chang CJ. The protective effect of habitual tea consumption on hypertension. *Arch Intern Med* 2004 Jul 26;164(14):1534-40.
 13. Giday A, Tadesse B. Prevalence and determinants of hypertension in rural and urban areas of southern Ethiopia. *Ethiop Med J* 2011 Apr;49(2):139-47.
 14. Sae-tan S, Grove KA, Lambert JD. Weight control and prevention of metabolic syndrome by green tea. *Pharmacol Res* 2011 Aug;64(2):146-54.
 15. Kurita I, Maeda-Yamamoto M, Tachibana H, Kamei M. Antihypertensive effect of Benifuuki tea containing O-methylated EGCG. *J Agric Food Chem* 2010 Feb 10;58(3):1903-8.
 16. Dauchet L, Amouyel P, Hercberg S, Dallongeville J. Fruit and vegetable consumption and risk of coronary heart disease: a meta-analysis of cohort studies. *J Nutr* 2006 Oct;136(10):2588-93.
 17. Seymour E.M, Singer A. M, Bennink M.R, Parikh R.W, Kirakosyan A, Kaufman P.B and Bolling S F. Chronic Intake of a Phytochemical-Enriched Diet Reduces Cardiac Fibrosis and Diastolic Dysfunction Caused by Prolonged Salt-Sensitive Hypertension. *J Gerontol A Biol Sci Med Sci* 2008 October; 63(10): 1034–1042.
 18. Melby MK, Murashima M, Watanabe S. Phytochemical intake and relationship to past health history in Japanese women. *Biofactors*. 2004;22(1-4):265-9.
 19. Moline J, Bukharovich I. F, Wolf M. S, Phillips R. Dietary flavonoids and hypertension: is there a link? *Medical Hypotheses* 2000; 55(4): 306–309.
 20. Cassidy A, O'Reilly ÉJ, Kay C, Sampson L, Franz M, Forman JP, Curhan G, Rimm EB. Habitual intake of flavonoid subclasses and incident hypertension in adults. *Am J Clin Nutr* 2011 Feb;93(2):338-47.
 21. Jouad H, Lacaille-Dubois M.A, Lyoussi B, Eddouks M. Effects of the flavonoids extracted from *Spergularia purpurea* Pers. on arterial blood pressure and renal function in normal and hypertensive rats. *Journal of Ethnopharmacology* 2001;76(2):159-163.
 22. Schini-Kerth VB, Auger C, Etienne-Selloum N, Chataigneau T. Polyphenol-induced endothelium-dependent relaxations role of NO and EDHF. *Adv Pharmacol* 2010;60:133-75.
 23. Stoclet JC, Chataigneau T, Ndiaye M, Oak MH, El Bedoui J, Chataigneau M, Schini-Kerth VB. Vascular protection by dietary polyphenols. *Eur J Pharmacol* 2004 Oct 1;500(1-3):299-313.
 24. Nicholson SK, Tucker GA, Brameld JM. Effects of dietary polyphenols on gene expression in human vascular endothelial cells. *Proc Nutr Soc* 2008 Feb;67(1):42-7.
 25. Medina-Remón A, Zamora-Ros R, Rotchés-Ribalta M, Andres-Lacueva C, Martínez-González MA, Covas MI and et al. Total polyphenol excretion and blood pressure in subjects at high cardiovascular risk. *Nutr Metab Cardio vas Dis* 2011 May;21(5):323-31.
 26. Ghosh D, Scheepens A. Vascular action of polyphenols. *Mol Nutr Food Res* 2009 Mar;53(3):322-31.
 27. Mennen LI, Sapinho D, Ito H, Bertrais S, Galan P, Hercberg S, Scalbert A. Urinary flavonoids and phenolic acids as biomarkers of intake for polyphenol-rich foods. *Br J Nutr* 2006 Jul;96(1):191-8.
 28. Suzuki A, Fujii A, Yamamoto N, Yamamoto M, Ohminami H, Kameyama A and et al. Improvement of hypertension and vascular dysfunction by hydroxyhydroquinone-free coffee in a genetic model of hypertension. *FEBS Lett* 2006 Apr 17;580(9):2317-22.
 29. Zhao Y, Wang J, Balleve O, Luo H, Zhang W. Antihypertensive effects and mechanisms of chlorogenic acids. *Hyperten Res* 2011 Nov 10. doi: 10.1038/hr.2011.195. [Epub ahead of print].
 30. Grassi D, Desideri G, Necozione S, Lippi C, Casale R, Properzi G, et al. Blood pressure is reduced and insulin sensitivity increased in glucose-intolerant, hypertensive subjects after 15 days of consuming high-polyphenol dark chocolate. *The Journal of Nutrition and Disease* November 2011;1671-6.
 31. Monroy-Ruiz J, María-Ángeles S, Rosalía C, María-José M. Astaxanthin-enriched-diet reduces blood pressure and improves cardiovascular

- parameters in spontaneously hypertensive rats. *Pharmacological Research* 2011; 63: 44–50.
32. Kooshki A, Golafruz Shahri M. Relationship between carotenoid intake and serum sugar and lipids in diabetics type 2 patients. *Ardebil J Med Scien* 2007;6(3):287-9.
 33. Tapiero H, Townsend DM, Tew KD. The role of carotenoids in the prevention of human pathologies. *Biomed Pharmacother* 2004 Mar;58(2):100-10.
 34. Paran E, Novack V, Engelhard YN, Hazan-Halevy I. The effects of natural antioxidants from tomato extract in treated but uncontrolled hypertensive patients. *Cardio vas Drugs Ther* 2009 Apr;23(2):145-51.
 35. Karin Ried, Peter Fakler. Protective effect of lycopene on serum cholesterol and blood pressure: Meta-analyses of intervention trials. *Maturitas* 2011;68: 299–310.

3/26/2016