

Biological Invasions, Its Type And Impact On Global Scale: A Review

¹Kuldip S. Dogra, ²Ravinder K. Kohli, ¹Sarvesh K. Sood, ²Parveen K. Dobhal and ¹Seema Sharma

¹Department of Biosciences, Himachal Pradesh University, Summer Hill, Shimla-171005, India

²Department of Botany, Panjab University Chandigarh- 160014, India

E.mail: dograks6@yahoo.co.in; Mobile: 919816740648; 919459170683

ABSTRACT: Biological invasion is now recognized world over and causing loss of indigenous wealth on earth. Biological invasions are altering ecosystem functions by reducing indigenous species diversity, altering soil dynamics and ecology. Economic losses due to invasion process found to be huge and difficult to measure. If this process of biological invasion is remain continuous for years to come than we can only transfer monocultures of species to our future generations. To preserve our indigenous species diversity it is important to understand the biological invasion its type and impact on global scale. In view of this present paper presents an insight into the biological invasion, its type and impact.

[Kuldip S. Dogra, Ravinder K. Kohli, Sarvesh K. Sood, Parveen K. Dobhal and Seema Sharma. **Biological Invasions, Its Type And Impact On Global Scale: A Review**. Report and Opinion 2011;3(1):9-20]. (ISSN: 1553-9873). <http://www.sciencepub.net>.

Keywords: Alien species, Biological Invasion, Biodiversity, Ecosystems, Invasive species,

INTRODUCTION

Biological invasions are important drivers of the escalating homogenization of the world's biota, and cause a substantial threat to biodiversity and ecological integrity of natural ecosystems (Booth et al 2003; Hulme 2003). Invasion by the species cause extensive effects on the habitats they invade, like impact on native species diversity, soil nutrient composition, altering forest fire cycles and loss of productivity of invading ecosystems. It also becomes a threat to endangered or threatened plant species around the world (Pimentel et al 2005). It is supposed that 10% of plant species, on an average, from any region are good colonizers. Thus, it can be estimated that from 260,000 vascular plant species known around the world, only 10% are potential invaders. Further, there are about 10,000 recognized invasives and 40% of these have been interchanged among different regions of the world (Rapoport 1991). It is also estimated that 20% or more of the plant species are exotics in many continental areas and 50% or more on many islands (Rejmanek and Randall 1994). The disturbances in the natural ecosystems provide the great opportunities to the alien invaders to establish themselves. The frequency of the exotic herbal plants increased in the areas of human interference such as forest fragmentation (Higgins et al 1996).

Introduced species that can rapidly achieve high densities may have greater establishment success

(Kolar and Lodge 2001) and dominate invaded communities to the exclusion of native species (Ortega and Pearson 2005). The species capable of rapid colonization are, in general, more likely to have negative impacts on biodiversity. Although the same traits that allow a species to successfully invade a broad range of communities could also magnify their impact (Callaway and Ridenour 2004). It is estimated that as many as 50% of invasive species in general can be classified as ecologically harmful, based on their actual impacts (Richardson et al 2000).

1 BIOLOGICAL INVASION AND ITS TYPE

Biological Invasion dramatically affects the distribution, abundance and reproduction of many native species (Sala et al 1999). Because of these ecological effects, alien species can also influence the evolution of natives exposed to novel interactions with invaders (Parker et al 1999). Evolutionary changes in natives in response to selection from aliens are usually overlooked, yet common responses include altered anti-predator defenses, changes in the spectrum of resources and habitats used, and other adaptations that allow native populations to persist in invaded areas (Mooney and Cleland 2001). So, introduction of such invasive species leads to change in the structure and composition of native communities (Rice and Emery 2003). It is basically of two types:

1. Human introduced invasions
2. Natural invasions

1.1 Human introduced invasions

The human-made introductions in the new habitats are quick and responsible for rapid change in the structure and composition of the communities (Ridenour and Callaway 2001). The invasion process by humans increased in the past five centuries, especially during the twentieth century, due to rapid increase in trade and travel across the globe. Planes, ships, and other forms of modern transport have allowed both deliberate and inadvertent movement of species, often resulting in unexpected and sometimes disastrous consequences (Moore 2004). Some times the species are introduced in such environments which can not be chosen by the species themselves for their growth and establishment. The introduction of new species in the balanced ecosystems and habitats can affect the natural process which leads towards destruction or loss of biodiversity (Louda et al 2003). Introduction of *Eucalyptus citriodora* Hook., *Populus deltoides* Marsh. and *Lantana camara* L. species in India is an example of human introduced invasions (Kohli et al 2004).

1.2 Natural invasions

The impact of natural invasion is almost similar to that of human made invasions but this kind of invasion mostly depends upon the dispersal ability of the invading plants and animals. The time scale for natural invasion can range from few years to several years. The sources for natural invasion are birds, animals, water and wind etc (Herbold and Moyle 1986). *Ageratum conyzoides* L. and *Parthenium hysterophorus* L. are examples of such type of invasions in India (Kohli et al 2004).

2 IMPACT OF BIOLOGICAL INVASIONS

Understanding the impacts of historical invasions can help us to predict the course of current invasions because the ecological effects of invasive species may change over time (Holway et al 2002). Studies of past introductions demonstrate that the effects of invasive species are complex and can permanently alter the structure of communities (Carlton 2003). Invasive alien species pose a threat to native plant communities globally, especially where these communities are disturbed (D'Antonio et al 2001). However, relatively few alien plant species seem to have the capacity to invade undisturbed native plant communities (Rejmánek 1989).

2.1 IMPACT ON THE NATURAL PLANT SPECIES

Disturbed and unattended habitats are more prone to the invasion as compare to the well-managed ecosystems and habitats. The habitats which have more diverse communities are highly competitive and resist invasion (Crawley 1987). For example, direct competition with the native flora can result in monocultures of an alien species, such as by *Psidium cattleianum* Sabine (strawberry guajava) in Mauritius and *Parthenium hysterophorus* L. (white top) in Australia and India (Evans 1997).

The invasive trees introduced in Florida (USA) cause major threats to the native vegetation. The tree species included Brazilian pepper (*Schinus terebinthifolius*), Australian paperbark (*Melaleuca quinquenrvia*) and Australian pine i.e. *Casuarina* sp. (Schmitz et al 1997). Invasion by different pine species becomes a problem to natural habitats in Australia, New Zealand, and South Africa (Richardson et al 1994). It was reported that in Christmas Island (Australia) 52.70% species have been found to be aliens (Claussen 2001) and most of them are confined to disturbed regions such as minefields, overburden dumps, and road sides. These include *Leucaena leucocephala*, *Muntingia calabura*, *Ricinis communis*, *Carica papaya* and *Psidium guajava* (Green et al 2004).

The alien terrestrial invaders are responsible for extensive and un-predictable irreversible changes to the natural habitats (Higgins et al 1999). Many exotic tree species worldwide introduced for commercial exploitation, economic reasons and for ornamental purpose have subsequently become noxious invaders. These tree species have impacted the natural above-ground herbal and other native vegetation. The allelopathic nature of these trees suppresses the growth of understory vegetation (Calder et al 1992). Some times, under the alien conditions or in new invaded ecosystems, such type of species become naturalized and expand over native ecosystems (Richardson 1998). For example the introduction of pines in the Southern Hemisphere has affected large areas of natural grass and shrub lands. It brought a lot of change in the dominant life forms, decreased the species composition and modified vegetation patterns and nutrient cycles (Richardson et al 1994).

The disturbed forest understories are more prone to invasion as compared to the undisturbed one. There are many species that establish and dominate low light forest understories in the Northwestern USA, such as *Alliaria petiolata*

(Meekins and McCarthy 2001), *Acer platanoides* (Webb and Kaunzinger 1993), *Lonicera bella*, *Rhamnus cathartica* (Harrington et al 1989) and *Berberies thunbergii* (Silander and Klepeis 1999).

The invasive species survive under the shade because rapid growth takes place in the gaps (Sanford et al 2003). Further, impact of some major invasive species on global scale is listed in the table, I.

Table I: Impact of some major invasive plants on global scale

S.N.	Botanical, Common Name and Habit of Alien Plants	Origin	Introduced	Threat	Reference
1.	<i>Acer platanoides</i> L. (Norway Maple) Tree	Norway	Suburban Ithaca, New York., USA	It displace native flora especially understorey vegetation, where it casts deep shade.	Martin 1999
2.	<i>Acroptilon repens</i> (L.) DC. (Russian Knapweed) Forb	Russia	North America.	Effecting the seedling emergence and growth of native grasses	Grant et al 2003
3.	<i>Agropyron cristatum</i> (L.) Gaertn Grass	North American	Asia	It is a strong competitor which can displace or prevent the establishment of native species.	Bakker and Wilson 2001
4.	<i>Ailanthus altissima</i> (Mill.) Swingle, (Tree of Heaven) Tree	China	USA	It is a serious threat to ecosystems in introduced areas, as the plant is very competitive, especially due to chemicals that may inhibit the growth of many native plants	Ding et al 2007
5.	<i>Alliaria petiolata</i> (Bieb.) Cav. & Grand. (Garlic Mustard) Herb	European	Eastern US and Canada	Its invasion poses a severe threat to native plant communities.	Nuzzo 1999
6.	<i>Alternanthera philoxeroides</i> Mart.) Griseb (Alligator Weed) Herb	South America	China	By forming dense mats of interwoven stems over water or land, this invasive weed may threaten the native flora and fauna, reduce crops yields, block ships, and promote flooding	Holm et al 1997
7.	<i>Andropogon gayanus</i> Kunth (Gamba Grass) Grass	Africa	Humid Central and Northern Brazil	Threat to native species	Fisher et al 1995
8.	<i>Andropogon gayanus</i> Kunth. (Gamba Grass) Grass	Africa	Northern Australia	Serious threat to northern Australia's savannas, with the potential to alter vegetation structure and initiate a grass fire cycle.	Rossiter et al 2003
9.	<i>Ardisia elliptica</i> Thunb. (Duck;s Eye) Shrub	Southeast Asia	Southern Florida, USA	Displacing native species since its invasion in Florida.	Koop 2004
10.	<i>Arundo donax</i> L. (Giant Reed) Grass	India	California	Displacing native species around the water channels.	Duke and Mooney 2004
11.	<i>Berberies thunbergii</i> DC	Japan	USA	Dense clusters of its	Cassidy et al 2004

	(Japanese Barberry) Shrub			own monocultures and altering the ecosystem process and functions in the undisturbed forests. Now, considered as a serious problem in forest management.	
12.	<i>Bothriochloa ischaemum</i> (L.) Keng (Yellow Blue Stem) Grass	Eurasian	Edwards Plateau of Central Texas, USA,	Fast spreading invasive grass which reduces native herbaceous plant diversity.	Gabbard 2007
13.	<i>Brachiaria brizantha</i> (A. Rich) Stapf (Signal Grass, Pasto Alambe) Grass	Humid tropical Africa	Humid Tropics	Threat to native plants.	Fisher et al 1995
14.	<i>Brachiaria mutica</i> (Forsk.) Stapf. (Pará Grass) Grass	Africa Tropical Africa	Humid tropics, sub-tropics	Serious threat to native vegetation.	Williams and Baruch 2000
15.	<i>Carduus acanthoides</i> L. (European Thistle) Herb	Europe	North America	It has a high probability of establishment and persistence and has a wide impact on native species.	Jongejans et al 2007
16.	<i>Carrichtera annua</i> (L. Aschers.) (Ward's Weed) Herb		South Australia.	Affecting the Native communities in chenopod shrublands of South Australia.	Harris and Facelli 2003
17.	<i>Celastrus orbiculatus</i> Thunb. (Oriental or Asian Bittersweet) Deciduous Liana	Southeast Asia	Southern Illinois, USA	It has created a ecological and economic threat to native ecosystems, particularly in disturbed temperate forest areas.	Pande et al 2007
18.	<i>Chromolaena odorata</i> (L.) R.M. King & H. Robin. (King in the Bush) Shrub	South Africa	America	It's foliage is reportedly flammable (contains essential oils), making it a threat to indigenous coastal forest patches, which are not resilient to fire.	Witkowski and Wilson 2001
19.	<i>Cinnamomum verum</i> Presl. (The True Cinnamon) Tree	Western Ghats of India and Sri Lanka,	Seychelles, Switzerland	It influences the course of forest succession by differentially affecting regeneration of native tree species.	Kueffer et al 2007
20.	<i>Conyza sumatrensis</i> (Retz.) E. Walker, (Fleabane) Herb		England UK	It spreads along transport routes from the capital (Greater London), such as railway lines, major roads and motorways. A threat to indigenous flora	Wurzell 1994
21.	<i>Cryptostegia grandiflora</i> (Roxb.) R. Br. (Rubber Vine) Vine	Madagascar	Australia	A threat to native biodiversity	Grice et al 2000
22.	<i>Cytisu scoparius</i> (L.) Link (English or Scotch Broom) Shrub	Europe	South Australia	It displaces native understorey vegetation and grasses, finally forming monospecific stands.	Fogarty and Facelli 1999

23.	<i>Elaeagnus angustifolia</i> L. (Russian Olive) Tree or shrub	Russia	North America	These invaders strongly influence the species composition, ecological processes, productivity, and biodiversity of native riparian forests	Lesica and Miles 2001
24.	<i>Eragrostis lehmanniana</i> Nees. Grass	South Africa	Arizona, Southwestern USA	Exist in a wide range of ecological conditions with little to no genetic variation	Schussman et al 2007
25.	<i>Heracleum mantegazzianum</i> Sommier & Levier (Giant Hogweed) Herb	Western part of the Greater Caucasus Russia, Georgia	Europe, North America and New Zealand	It is a problem species, because it forms monocultures with a high cover, replaces resident vegetation, and produces photosensitive sap that is toxic for humans.	Nehrbass et al 2007
26.	<i>Heracleum mantegazzianum</i> Sommier et Levier (Giant Hogweed) Herb	Western part of the Greater Caucasus Russia, Georgia	Czech Republic	Replacement of native vegetation and injuries to human skin caused by phototoxic substances are the main reasons for attempts to eradicate it.	Pys'ek et al 2007
27.	<i>Heracleum persicum</i> Desf. ex Fischer. Herb	Turkey, Iran, and Iraq.	Scandinavia, Europe	Posing threat to native plant diversity.	Jahodová et al 2007
28.	<i>Heracleum sosnowskyi</i> Manden Herb	Eastern and central Caucasus, Transcauc-asia and northeast Turkey.	Russia, Belarus, Ukraine, Baltic countries, Eastern Germany.	Widely invasive species in various habitats.	Nielsen et al 2005
29.	<i>Hyparrhenia rufa</i> (Nees) Stapf. (Jaraguá Grass) Grass	Tropical and South Africa and Madagascar	Humid and sub-humid tropics, Central and South America, Cuba	African grass-dominated pasture produces radical changes in hydrologic balance.	Williams and Baruch 2000
30.	<i>Imperata cylindrica</i> (L.) Beauv. (Cogongrass, Alang-Alang) Grass	Subtropical and tropical Asia.	Southeastern USA	It displaces native plant and animal species and alters fire regimes. The Phosphorus addition by this species reduces invasion of a longleaf pine savanna in Southeastern USA.	Brewer and Cralle 2003
31.	<i>Ligustrum robustum</i> (Roxb.) (Srilakan Privet) Shrub	South Indian Hills (Western Gh'ats) and Sri Lanka (Central Highlands)	La R'union, Mauritius	This species is a major threat to the native ecosystems and one of the worst pest plants of Mauritian wet forests.	Lavergne et al 1999
32.	<i>Ligustrum sinense</i> Lour. (Chinese Privet) Shrub	China	North Carolina, USA	Has impact on the native plant diversity and on forest regeneration.	Merriam and Feil 2002
33.	<i>Lolium multiflorum</i> (Pers.) (Italian Raygrass) Grass	Spain	Alkali sites in California, USA	It may pose a long-term threat to native alkali biodiversity.	Dawson et al 2007
34.	<i>Lonicera maackii</i> (Rupr.) (Herder Amur-Honeysuckle)	Northeastern Asia,	Eastern North America,	Associated with reduced tree seedling density in Ohio	Gorchov and Trisel 2003

	Shrub			forests.	
35.	<i>Lythrum salicaria</i> L. (Purple loosestrife) Herb	Eurassia	North America	The plant is considered a problem because it displaces native plants, clogs waterways, and reduces the quality of habitat for wildlife.	Shadel and Molofsky 2002
36.	<i>Melia azadratica</i> L. (Dreak) Tree	India	South America	Threat to the native plant species diversity.	Luti et al 1979
37.	<i>Melinis minutiflora</i> Beauv. (Molasses Grass, Gordura) Grass	Tropical West Africa, Angola to Cameroon	Tropics of Central and South America, West Indies, Puerto Rico	The conversion of native forest to pasture and the establishment of African grasses can significantly alter water balance of ecosystems and watersheds.	Williams and Baruch 2000
38.	<i>Microstegium vimineum</i> (Trin.) A. Camus, (Japanese Stilt Grass) Grass	Japan	Eastern and South Eastern United States	Threatening the growth of native species.	Redman 1995
39.	<i>Panicum maximum</i> Jacq., (Guinea Grass) Grass	Tropical Africa to sub-tropics of South Africa	South Eastern US, West Indies, Tropics of Central and South America	These effects are produced primarily by the extreme structural changes of the vegetation.	Williams and Baruch 2000
40.	<i>Pennisetum clandestinum</i> Hochst. (Buffel Grass) Grass	Hotter and drier parts of eastern and southern Africa	Semi-arid and arid tropics and sub-tropics, North Mexico, southwestern US	Posing a threat to the native species.	Williams and Baruch 2000
41.	<i>Phacelia tanacetifolia</i> Benth. (Lacy Phacelia) Herb	North America	Europe	Affect pollinator visitation and female reproductive success of a native plant, <i>Melampyrum pretense</i> L. in recently disturbed and undisturbed boreal forests.	Totland et al 2006
42.	<i>Polygonum perfoliatum</i> L (Devil's Tail Tear-Thumb or Mile-a-Minute Weed) Herb	India, China, Korea, Japan, Bangladesh, and the Philippines	Northeastern USA	Causing ecological problems in invaded areas, as the plant grows rapidly and covers shrubs and other vegetation, dominating in its new community.	Ding et al 2007
43.	<i>Prosopis juliflora</i> (Swartz) DC. (Vilayeti Babul) Tree	Jamaica	India	Aggressive and has not only successfully invaded several habitats but has also caused substratum degradation in these by causing loss of finer soil particles.	Sharma and Dakshini 1998
44.	<i>Pueraria Montana</i> (Lour.) Merr. var. lobata (Willd.) Maesen & Almeida (Kudzu) Vine	Asian	Philadelphia, USA	It poses a serious threat to biodiversity as it completely replaces existing vegetation and few plants can survive once smothered by kudzu.	Ding et al 2007
45.	<i>Rubus alceifolius</i> Poirlet (The Giant Bramble) Shrub	Southeastern Asia and Malaysia,	Island of La-Réunion	Invaded a wide variety of habitats- lowland rainforest, mountain and submountain rainforest, <i>Acacia</i>	Baret et al 2004

				<i>heterophylla</i> rainforest and disturbing native species.	
46.	<i>Senecio madagascariensis</i> Poir. (Fireweed) Shrub	Afro- Madagascan	Hawaii	Competes strongly with existing pasture flora for light, moisture, and soil nutrients (notably P and N), leading to the ultimate deterioration of pastures.	Roux et al 2006
47.	<i>Sorghum halepensis</i> L. (Johnson Grass) Grass	Mediterranean North Africa	Sub-tropics and warm temperate North and South America	Threat to indigenous plants.	Williams and Baruch 2000
48.	<i>Spartina densiflora</i> Braigu (Cordgrass) Grass	Chile	Gulf of Cadiz, Spain,	Poses a threat to the biodiversity of southern European marshes.	Castillo et al 2000
49.	<i>Tamarix ramosissima</i> Ledeb. (Salt Cedar) Tree	Russia	North America	These invaders strongly influence the species composition, ecological processes, productivity, and biodiversity of native riparian forests	Stohlgren et al 1998
50.	<i>Tibouchina herbacea</i> (DC) Cogn. (Glory Bush) Shrub	South America	Hawaii and Maui	Effecting the growth of native vegetation under the native forests.	Almasi 2000
51.	<i>Tradescantia fluminensis</i> Vell. (Small Leaf Spiderwort) Herb	South America	New Zealand	Effecting the litter decomposition and nutrient availability in a remnant of New Zealand lowland podocarp broadleaf forest.	Standish et al 2004
52.	<i>Ziziphus mauritiana</i> Lam (Indian Jujuba) Shrub or Tree	India	Australia	A threat to native biodiversity.	Grice et al 2000

2.2 IMPACT ON THE SOIL DYNAMICS

Invasion by alien plant species affects the dynamics and composition of soil on a wide scale and have great impact on ecosystem functions such as soil nutrient cycling. Since these impacts result from differences in traits between the exotic and resident species, novel physiological traits such as nitrogen cycling may cause large alterations in ecosystem function (Yelenik et al 2007).

The invasion of two exotic plants – *Berberis thunbergii* and *Microstegium vimineum* – in hardwood forests of New Jersey, Europe has shown a significant increase of pH in soils under the invasive plants as compared to soils from under native shrubs (*Vaccinium* sp.). Further, available nitrate and net potential nitrification were significantly higher in soils under the two exotic species (Kourtev et al 1999). The introduced *Prosopis juliflora* is fast

growing, highly aggressive and invasive, and causes substratum degradation in the semi-arid and arid areas of north and north-west India as compared to native species *P. cineraria*. This lack of integration amongst plant and soil characteristics and the ability to meet its nutrient requirements in all situations could be the basis of the phenomenal spread of *P. juliflora* across varying environmental conditions, in contrast to *P. cineraria* (Sharma and Dakshini 1998).

Comparisons between habitats with contrasting levels of soil resource availabilities suggest that an increase in resource availability tends to increase invasion of native grassland communities by non-native plants. For example, nutrient enrichment (Davis et al 2000; Kolb et al 2002) has been consistently shown to increase the abundance of non-native plant species and decrease the abundance of native ones (White et al 1997).

2.3 ECONOMIC LOSS DUE TO INVASION

The risks associated with the invasion of alien species are increasing, with increasingly rapid international exchange and convenient transportation (Chen and Xu 2001). Invasive alien species expedites the losses of species and genetic biodiversity (Li and Xie 2002; Wan et al 2002), destroys the structure and functions of ecosystems (Zhang and Ye 2002), and causes huge economic losses. Invasive alien species have caused losses worth USD 138 billion to the USA (Pimentel et al 2000). The total economic losses caused by invasive alien species to China were to the time of USD 14.45 billion, with direct and indirect economic losses accounting for 16.59% and 83.41% of total economic losses, respectively (Xu and Ding 2003). Oerke et al (1994) calculated 13% loss in the world's agricultural output due to weeds (based on eight major crops). In maize alone, an actual loss due to weeds from 1997 to 1999 was around 1.7 billion USD. There has been an extensive movement of plant species around the world by humans as a consequence of trading activities. This has resulted in exotic species forming a significant part of the agricultural weed flora, and in natural ecosystems invasive weeds are almost exclusively alien (Groves et al 2001).

CONCLUSIONS

Biological invasion of species alter native community composition, deplete species diversity, affect ecosystem process and thus cause huge economic and ecological imbalance. Studies of invasive species introductions in the past revealed that the impacts of their invasion are complex and can permanently alter the structure and function of communities, cause local extinctions and changes in ecosystem processes. The increased incidence of invasion around the world poses a major threat to indigenous biological diversity.

Increase in the rate of invasion and deliberate introduction of aliens into an area by man is the byproduct of the globalization of regional economics. Large parts of the world are currently dominated by human modified ecosystems that often comprise a greater biomass of introduced than native organisms (Vitousek et al 1997). Besides human actions, several other factors contribute to successful invasion by alien plants. The climatic and edaphic similarities between the original and new habitats are very important factors for the establishment of alien species (Holdgate 1986). Thus, humid tropics of the Asia and Africa with highly leached soils are similar to Latin American home of species such as *Lantana camara*, *Ageratum conyzoides*, *Eupatorium*

odoratum, *Parthenium hysterophorus* and *Mikania micrantha* enabling them to invade and colonise appropriate sites on these two continents (Ramakrishnan 1991).

The magnitude and net effects of biological invasion escalated rapidly over the twentieth century. During each decade, more species become invasive, more ecosystems were irreversibly altered, and an ever-increasing array of functions and processes was impacted by invasive alien species (Rejmanek 2000). There are thousands of alien species known to establish around the world and many more introduced species remain undetected or unrecognized (Ruiz et al 2000a). Their invasion cause a wide range of high-impact and high profile impacts, including decline in population of threatened and endangered species, habitat alteration and loss, increased frequency of fires, shifts in food webs and nutrient cycling, loss of agricultural crops and productive lands. So, Biological invasion are clearly a potent force of change, operating on a global scale and affecting many dimensions of society (Wilcove et al 1998; Ohlemuller et al 2006). In view of the wide range of impacts of biological invasion as mentioned above comprehensive studies on long term basis are required at a global scale.

REFERENCES

1. Almasi KN (2000) A non-native perennial invades a native forest. *Biol. Invasions* 2: 219-230.
2. Bakker J and Wilson S (2001) Competitive abilities of introduced and native grasses. *Plant Ecol* 157: 117-125.
3. Baret S, Maurice S, Le Bourgeois T. and Strasberg D (2004) Altitudinal variation in fertility and vegetative growth in the invasive plant *Rubus alceifolius* Poir. (Rosaceae), on Reunion Island. *Plant Ecol* 172: 265-273.
4. Booth BD, Murphy SP and Swanton CJ (2003) *Weed Ecology in Natural and Agricultural Systems*. CABI Publishing, Willingford, Oxfordshire, UK, Pp. 288.
5. Brewer JS and Cralle SP (2003) Phosphorus addition reduces invasion of a longleaf pine savanna (Southeastern USA) by a non-indigenous grass (*Imperata cylindrica*). *Plant Ecol* 167: 237-245.
6. Calder JA, Wilson JB, Mark AF and Ward G (1992) Fire, succession and reserve management in a New Zealand snow tussock grassland. *Biol Conser* 62: 35-45.
7. Callaway RM and Ridenour WM (2004) Novel weapons: Invasive success and the evolution of increased competitive ability. *Front Ecol Environ* 2: 436-443.

8. Carlton JT (2003) Community assemblage and historical biogeography in the North Atlantic Ocean: The potential role of human-mediated dispersal vectors. *Hydrobiology* 503: 1-8.
9. Cassidy TM, Fownes JH and Harrington RA (2004) Nitrogen limits an invasive perennial shrub in forest understorey. *Biol Invasions* 6: 113-121.
10. Castillo JM, Fernandez-Baco L, Castellanos EM, Luque CJ, Figueroa ME and Davy AJ (2000) Lower limits of *Spartina densiflora* and *S. maritima* in a Mediterranean salt marsh determined by different ecophysiological tolerances. *J Ecol* 88: 801-812.
11. Chen LY and Xu HG (2001) Australian management strategy for invasive alien species and references available to China. *Biod Sci* 9(4): 466-471.
12. Claussen J (2001) *Plants of Christmas Island*. Version 2. Parks Australia North, Christmas Island, Indian Ocean.
13. Crawley MJ (1987) What makes a community invulnerable? In: *Colonization, Succession and Stability* (Eds Gray AJ, Crawley MJ and Edwards PJ). *The 26th Symposium of the British Ecological Society*. Blackwell Scientific Publications, London, UK, Pp. 429-453.
14. D'Antonio CM, Levine JM and Thomson M (2001) Ecosystem resistance to invasion and the role of propagule supply: A California perspective. *J Mediterr Ecol* 27: 233-245.
15. Davis MA, Grime JP and Thompson K (2000) Fluctuating resources in plant communities: A general theory of invasibility. *J Ecol* 88: 528-534.
16. Dawson K, Veblen KE and Young TP (2007) Experimental evidence for an alkali ecotype of *Lolium multiflorum*, an exotic invasive annual grass in the central valley, CA, USA. *Biol Invasions* 9: 327-334.
17. Ding J, Reardon R, Wu Y, Zheng H and Fu W (2007) Biological control of invasive plants through collaboration between China and the United States of America: A perspective. *Biol Invasions* 8: 1439-1450.
18. Duke JS and Mooney HA (2004) Disruption of ecosystem processes in western North America by invasive species. *Rev Chil Hist Nat* 77: 411-437.
19. Evans HC (1997) *Parthenium hysterophorus*: A review of its weed status and the possibilities for biological control. *Biocon News Infor* 18: 89-98.
20. Fisher MJ, Rao IM, Ayarza MA, Lascano CE, Sainz JI, Thomas RJ and Vera RR (1995) Scientific correspondence, reply. *Nature* 376: 473.
21. Fogarty G and Facelli JM (1999) Growth and competition of *Cytisus scoparius*, an invasive shrub, and Australian native shrubs. *Plant Ecol* 144: 27-35.
22. Gabbard BL and Fowler NL (2007) Wide ecological amplitude of a diversity-reducing invasive grass. *Biol Invasions* 9: 149-160.
23. Gorchov DL and Trisel DE (2003) Competitive effects of the invasive shrub, *Lonicera maackii* (Rupr.) Herder (Caprifoliaceae) on the growth and survival of native tree seedlings. *Plant Ecol* 166:13-24.
24. Grant DW, Debra PCP, George KB and Harold DF (2003) Influence of an exotic species, *Acroptilon repens* (L.) DC. on seedling emergence and growth of native grasses. *Plant Ecol* 166: 157-166.
25. Green PT, Lake PS and O'Dowd JD (2004) Resistance of island rainforest to invasion by alien plants: Influence of microhabitat and herbivory on seedling performance. *Biol Invasions* 6: 1-9.
26. Grice AC, Radford IJ and Abbott BN (2000) Regional and landscape-scale patterns of shrub invasion in tropical savannas. *Biol Invasions* 2: 187-205.
27. Groves RH, Panetta FD and Virtue JG (2001) *Weed Risk Assessment*. CSIRO Publishing, Collingwood, Australia, Pp. 244.
28. Harrington RA, Broom BJ and Reich PB (1989) Ecophysiology of exotic and native shrub in southern Wisconsin: Relationship of leaf characteristics, resources availability and phenology to seasonal patterns of carbon gain. *Oecologia* 80: 356-367.
29. Harris MR and Facelli JM (2003) Competition and resource availability in an annual plant community dominated by an invasive species, *Carrichtera annua* (L. Aschers.), in South Australia. *Plant Ecol* 167: 19-29.
30. Herbold B and Moyle PB (1986) Introduced species and vacant niches. *Amer Natur* 128: 751-760.
31. Higgins SI, Richardson DM and Cowling RM (1996) Modeling invasive plant spread: The role of plant-environment interactions and model structure. *Ecology* 77: 2043-2054.
32. Higgins SI, Richardson DM, Cowling RM and Smith THT (1999) Predicating the landscape scale distribution of alien plants and their threat to biodiversity. *Conserv Biol*. 13: 303-313.
33. Holdgate MW (1986) *Summary and Conclusions: Characteristics and Consequences of Biological Invasions*. Philosophical Transactions of the Royal Society, London.
34. Holm L, Doll J, Holm E, Pancho JV and Herberger JP (1997) *World Weeds: Natural Histories and Distribution*. John Wiley and Sons Inc., New York.
35. Holway DA, Lach L, Tsutsui ND and Case TJ (2002) The causes and consequences of ant invasions. *Ann Rev Ecol Syst* 33: 181-233.
36. Hulme PE (2003) Biological invasions: Winning the science battles but losing the conservation war? *Oryx* 37: 178-193.
37. Jahodová S, Trybush I S, Pysek P, Wade M. and Karp A (2007) Invasive species of *Heracleum* in Europe:

- An insight into genetic relationships and invasion history. *Diversity Distrib* 13: 99-114.
38. Jongejans E, Skarpaas O, Tipping PW and Shea K (2007) Establishment and spread of founding populations of an invasive thistle: The role of competition and seed limitation. *Biol Invasions* 9: 317-325.
 39. Kohli RK, Dogra KS, Batish DR and Singh HP (2004) Impact of invasive plants on the structure and composition of natural vegetation of north western Indian Himalayas. *Weed Tech* 18: 1296-1300.
 40. Kolar CS and Lodge DM (2001) Progress in invasion biology: Predicting invaders. *Trends Ecol Evol* 16: 199-204.
 41. Kolb A, Alpert P, Enters D and Holzapfel C (2002) Patterns of invasion within a grassland community. *J Ecol* 90: 871-881.
 42. Koop AL (2004) Differential seed mortality among habitats limits the distribution of the invasive non-native shrub *Ardisia elliptica*. *Plant Ecol* 172: 237-249.
 43. Kourtev PS, Huang WZ and Ehrenfeld JG (1999) Differences in earthworm densities and nitrogen dynamics in soils under exotic and native plant species. *Biol Invasions* 1: 237-245.
 44. Kueffer C, Schumacher E, Fleischmann K, Edwards PJ and Dietz H (2007) Strong below-ground competition shapes tree regeneration in invasive *Cinnamomum verum* forests. *J Ecol* 95: 273-282.
 45. Lavergne C, Rameau J-C and Figier J (1999) The invasive woody weed *Ligustrum robustum* subsp. *walkeri* threatens native forests on La Réunion. *Biol Invasions* 1: 377-392.
 46. Lesica P and Miles S (2001) Natural history and invasion of Russian olive along eastern Montana rivers. *Western North Amer Natur* 61: 1-10.
 47. Li ZY and Xie Y (2002) *Invasive Alien Species in China* (in Chinese). Forestry Publishing Company of China, Beijing.
 48. Louda SM, Pemberton RW, Johnson MT and Follett PA (2003) Non-target effects: The Achilles heel of biological control? *Ann Rev Entom* 48: 365-396.
 49. Luti R, Galera MA, Muller N, Berzal M, Nores M, Herrera M and Barrera JC (1979) Vegetación. In: *Geografía Física de la Provincia de Córdoba* (Eds Vazquez JB, Miatello R and Roque M). Boldt, Buenos Aires, Pp. 297-368.
 50. Martin P (1999) Norway maple (*Acer platanoides*) invasion of a natural forest stand: Understorey consequence and regeneration pattern. *Biol Invasions* 1: 215-222.
 51. Meekins JF and McCarthy BC (2001) Effects of environmental variation on the invasive success of non-indigenous forest forbs. *Ecol Appl* 11(5): 1336-1348.
 52. Merriam RW and Feil E (2002) The potential impact of an introduced shrub on native plant diversity and forest regeneration. *Biol Invasions* 4: 369-373.
 53. Mooney HA and Cleland EE (2001) The evolutionary impact of invasive species. *Proceedings of the National Society of Sciences, USA*. 98, Pp. 5446-5451.
 54. Moore PD (2004) Favoured aliens for the future. *Nature* 427: 594.
 55. Nehrbass N, Winkler E, Müllerová J, Pergl J, Pysěk P and Perglova I (2007) A simulation model of plant invasion: Long-distance dispersal determines the pattern of spread. *Biol Invasions* 9: 383-395.
 56. Nielsen C, Ravn HP, Cock M and Nentwig W (Eds) (2005) *The giant hogweed best practice manual. Guidelines for the management and control of an invasive alien weed in Europe*. Forest and Landscape Denmark, Hoersholm, Denmark.
 57. Nuzzo V (1999) Invasion pattern of the herb garlic mustard (*Alliaria petiolata*) in high quality forests. *Biol. Invasions* 1: 169-179.
 58. Oerke EC, Dehne DW, Schonbeck F and Weber A (1994) *Crop Production and Crop Protection: Estimated Losses in Major Food and Cash Crops*. Elsevier, Amsterdam, Pp. 808.
 59. Ohlemuller R, Walker S and Wilson JB (2006) Local vs regional factors as determinants of the invisibility of indigenous forest fragments by alien plant species. *Oikos* 112: 493-501.
 60. Ortega YK and Pearson DE (2005) Weak vs. strong invaders of natural plant communities: Assessing invasibility and impact. *Ecol Appl* 15: 651-661.
 61. Pande A, Williams CL, Lant CL and Gibson DJ (2007) Using map algebra to determine the mesoscale distribution of invasive plants: The case of *Celastrus orbiculatus* in Southern Illinois, USA. *Biol Invasions* 9: 419-431.
 62. Parker IM, Simberloff D, Lonsdale WM, Goodell K, Wonham M, Kareiva PM, Williamson M.H, Von-Holle B, Moyle PB, Byers JE and Goldwasser L (1999) Impact: Toward a framework for understanding the ecological effects of invaders. *Biol Invasions* 1: 3-19.
 63. Pimentel D, Lach L, Zuniga R and Morrison D (2000) Environmental and economic costs of non indigenous species in the United States. *Biosciences* 50: 53-65.
 64. Pimentel D, Zuniga R and Morrison D (2005) Update on the environmental and economic costs associated with alien-invasive species in the United States. *Morrison* 52: 273-288.
 65. Pysek P, Krinke L, Jarosik V, Perglova I, Pergl J and Moravcova L (2007) Timing and extent of tissue removal affect reproduction characteristics of an

- invasive species *Heracleum mantegazzianum*. *Biol Invasions* 9: 335-351.
66. Ramakrishnan PS (1991) *Ecology of Biological Invasions in the Tropics*. International Scientific Publications, New Delhi.
 67. Rapoport EH (1991) Tropical versus temperate weeds: A glance into the present and future. In: *Ecology of Biological Invasion in the Tropics* (Ed Ramakrishnan PS). International Scientific Publications, New Delhi.
 68. Redman DE (1995) Distribution and habitat types for Nepal *Microstegium* [*Microstegium vimineum* (Trin.) Camus] in Maryland and the district of Columbia. *Castanea* 60: 270-275.
 69. Rejmanek M (1989) Invasibility of plant communities. *Biological Invasions: A Global Perspective* (Eds Drake JA, Mooney HA, di-Castri F, Groves RH, Kruger FJ, Rejmanek M and Williamson M). Wiley and Sons, Chichester, England, Pp 369-388.
 70. Rejmanek M (2000) Invasive plants: Approaches and predictions. *Aust. Ecol.* 25: 497-506.
 71. Rejmanek M and Randall JM (1994) *Invasive alien plants in California*: 1993 Summary and comparison with other areas in North America. *Madrono* 41: 161-177.
 72. Rice KJ and Emery NC (2003) Managing microevolution: Restoration in the face of global change. *Front Ecol Environ* 9: 469-478.
 73. Richardson DM (1998) Forestry trees as invasive aliens. *Conser Biol* 12: 18-26.
 74. Richardson DM, Pysek P, Rejmánek M, Barbour MG, Panetta FD and West CJ (2000) Naturalization and invasion of alien plants: Concepts and definitions. *Diversity Distrib* 6: 93-107.
 75. Richardson DM, Williams PA and Hobbs RJ (1994) Pine invasions in the Southern Hemisphere: Determinants of spread and invadability. *J Biogeography* 21: 511-727.
 76. Ridenour WM and Callaway RM (2001) The relative importance of allelopathy in interference: The effects of an invasive weed on a native bunchgrass. *Oecologia* 126: 444-450.
 77. Rossiter NA, Setterfield SA, Douglas MM and Hutley LB (2003) Testing the grass-fire cycle: Alien grass invasion in the tropical savannas of northern Australia. *Diversity Distrib* 9: 169-176.
 78. Ruiz GM, Fofonoff PW, Carlton JT, Wonham MJ and Hines AH (2000) Invasion of coastal marine communities in North America: Apparent patterns, processes, and biases. *Ann Rev Ecol.Syst* 31: 481-531.
 79. Sale OE, Chapin FS, Gardner RH, Lauenroth WK, Mooney HA and Ramakrishnan PS (1999) Global change, biodiversity and ecological complexity. In: *The Terrestrial Biosphere and Global Change: Implications for Natural and Managed Ecosystems* (Eds Walker B, Steffen W, Canadell J and Ingram J). Cambridge University Press, Cambridge, UK, Pp. 304-328.
 80. Sanford NL, Harrington RA and Fownes JH (2003) Survival and growth of native and alien woody seedlings in open and understorey environments. *For Ecol Manage* 183: 377-385.
 81. Schmitz DC, Simberloff D, Hofstetter RH, Haller W and Sutton D (1997) The ecological impact of non indigenous plants. In: *Strangers in Paradise. Impact and Management of Non Indigenous Species in Florida* (Eds Simberloff D, Schmitz DC and Brown TC). Island Press, Washington, DC, Pp. 39-61.
 82. Schussman H, Geiger E, Mau-Crimmins T and Ward J (2007) Spread and current potential distribution of an alien grass, *Eragrostis lehmanniana* Nees, in the southwestern USA: Comparing historical data and ecological niche models. *Diversity Distrib* 12: 582-592.
 83. Shadel WP and Molofsky J (2002) Habitat and population effects on the germination and early survival of the invasive weed, *Lythrum salicaria* L. (purple loosestrife). *Biol Invasions* 4: 413-423.
 84. Sharma R and Dakshini KMM (1998) Integration of plant and soil characteristics and the ecological success of two *Prosopis* species. *Plant Ecol* 139: 63-69.
 85. Silander JAJ and Klepesis DM (1999) The invasion ecology of Japness Barberry (*Berberis thunbergia*) in the New England landscape. *Biol Invasions* 1: 189-201.
 86. Standish RJ, Williams PA, Robertson AW, Scott NA and Hedderley DI (2004) Invasion by a perennial herb increases decomposition rate and alters nutrient availability in warm temperate lowland forest remnants. *Biol Invasions* 6: 71-81.
 87. Stohlgren TJ, Bull KA and Otsuki Y (1998) Comparison of rangeland vegetation sampling techniques in the central grasslands. *J Range Manage* 51: 164-172.
 88. Totland Q, Nielsen A, Bjerknes A-L and Ohlson M (2006) Effect of an exotic plant and habitat distribution on pollinator visitation and reproduction in a boreal forest herb. *Amer J Bot* 93 (6): 868-873.
 89. Vitousek PM, Mooney HA, Lubchenco J and Melillo JM (1997) Human domination of earth's ecosystems. *Science* 277: 494-499.
 90. Wan FH, Guo JY and Wang DH (2002) Alien invasive species in China: Their damages and management strategies. *Biod Sci* 10(1): 119-125.
 91. Webb SA and Kaunzinger CK (1993) Biological invasion of the Drew University (New Jersey) forest preserve by Norway maple (*Acer platanoides* L.). *Bull Torr Bot Club* 120: 343-349.

92. White TA, Campbell BD and Kemp PD (1997) Invasion of temperate grassland by a subtropical annual grass across an experimental matrix of water stress and disturbance. *J Veg Sci* 8: 847-854.
93. Wilcove DS, Rothstein D, Dobow J, Phillips A and Losos E (1998) Quantifying threats to imperiled species in the US. *Biosciences* 48: 607-615.
94. Williams DG and Baruch Z (2000) African grass invasion in the Americas: Ecosystem consequences and the role of ecophysiology. *Biol Invasions* 2: 123-140.
95. Witkowski ETF and Wilson M (2001) Changes in density, biomass, seed production and soil seed banks of the non-native invasive plant, *Chromolaena odorata*, along a 15 year chronosequence. *Plant Ecol* 152: 13-27.
96. Wurzell B (1994) A history of *Conyza* in London. *Botanical Society of British Isles New.* 65: 34-39.
97. Xu HG and Ding H (2003) Countermeasures for the prevention of invasive alien species. In: *Conserving Biodiversity and Strengthening Nature Reserve Management* (Eds Wang DH and Fang C). China Environment Sciences Press, Beijing, Pp. 128-139.
98. Yelenik SG, Stock WD and Richardson DM (2007) Functional group identity does not predict invader impacts: Differential effects of nitrogen-fixing exotic plants on ecosystem function. *Biol Invasions* 9: 117-125.
99. Zhang LY and Ye WH (2002) Community invasibility and its influencing factors. *Acta Phytocologica Sinica* 26(1): 109-114.

7/1/2010