

Effect of diet supplemented with pumpkin (*Cucurbita moschata*) and black seed (*Nigella sativa*) oils on performance of rabbits: 2- Productive and reproductive performance of does and their offspring

Gaafar, H.M.A.; Ayat A. Ragab and K.F.A. El-Reidy

Animal Production Research Institute, Agriculture Research Center, Dokki, Giza. hamedgaafar@gmail.com

Abstract: Thirty six New Zealand white (NZW) doe rabbits with initial body weight of 1445.2±39.4 g and aged 12 weeks were used in a complete randomized design experiment with four dietary treatments. Does were fed commercial diet without additive (control, G1) or supplemented with 5 g pumpkin seed (PS) oil/kg diet (G2), 5 g black seed (BS) oil/kg diet (G3) and 2.5 g PS oil plus 2.5 g BS oil /kg diet (G4). Rabbit does in G4 showed the highest (P<0.05) body weight (BW) and average daily gain at the different age intervals, followed by those in G2 and G3, while does in G1 had the lowest weights. Does in G4 showed significantly (P<0.05) the highest values of BW at mating and parturition, the 1st service conception rate and the lowest number of service per conception, followed by those in G2 and G3, while does in G1 had the poorest values. Does in G2, G3 and G4 increased average daily milk yield (as overall for the suckling period) by 9.57, 20.00 and 27.63% compared with G1, respectively. Milk of does in G4 showed significantly (P<0.05) the highest percentage of fat, protein, lactose, total solids and solids not fat, followed by milk of does in G2 and G3, while milk of G1 had the lowest percentages. In blood serum of does, G4 showed significantly (P<0.05) the highest concentrations of total protein and globulin, while those in G2 showed significantly (P<0.05) the highest albumin and glucose concentrations. Does in G1 showed significantly (P<0.05) the highest total lipids, triglycerides, total cholesterol, HDL and LDL concentrations. Activity of AST and ALT was not affected by supplementation. The concentrations of prolactin and thyroxine (T4) hormones were higher in G4, while triiodothyronine hormone (T3) was higher in G3. The concentrations of IgG and IgM in serum of both does and their offspring were the highest in G4 as compared to other groups, but the differences were not significant. Does in G3 showed the highest litter size at birth, but those in G4 showed the highest (P<0.05) litter size at 7, 14, 21 and 28 days of suckling period. Bunnies of G1 showed the highest (P<0.05) mortality rate, followed by those in G2 and G3, while G4 had the lowest rate. Does in G2 showed the heaviest (P<0.05) litter weight at birth, while G4 had the highest litter weight at 28 days of suckling period. Bunnies of G3 and G4 recorded the heaviest (P<0.05) litter weight gain for the whole suckling period. Bunnies in G2 showed significantly (P<0.05) the highest BW and weight gain during the different suckling periods, while the lowest weight gain was recorded for bunnies in G1. Pumpkin and black seeds oils supplementation improved economic efficiency of does and their offspring, which net revenue of G2, G3 and G4 increased by 9.77, 25.23 and 34.75% compared with G1, respectively. Does rabbits fed diet supplemented with a combination of pumpkin and black seeds oils (2.5 g PS oil plus 2.5 g BS oil/ kg diet) and their offspring showed the best results.

[Gaafar, H.M.A.; Ayat A. Ragab and K.F.A. El-Reidy. **Effect of diet supplemented with pumpkin (*Cucurbita moschata*) and black seed (*Nigella sativa*) oils on performance of rabbits: 2- Productive and reproductive performance of does and their offspring.** *Rep Opin* 2014;6(1):60-68]. (ISSN: 1553-9873). <http://www.sciencepub.net/report>. 11

Keywords: NZW rabbits, pumpkin, black seed, reproduction, offspring growth.

1. Introduction

Rabbits can play a significant role in solving the problem of meat shortage in many parts of the world, due to their high potential for reproduction, rapid growth rate, short generation interval, ample nutritional spectrum, limited vital space and ease of rearing. However, pre- and post-weaning mortality until marketing limits the crop of rabbits in kilograms and a lower income (Rashwan ana Marai, 2000).

Eight fatty acids (99.5%) and thirty-two compounds (86.7%) have been identified in the fixed and volatile oils of *Nigella sativa* seeds, respectively. The main fatty acids of the fixed oil were linoleic acid (55.6%), oleic acid (23.4%) and palmitic acid (12.5%). The major compounds of the volatile oil were trans-

anethole (38.3%), p-cymene (14.8%), limonene (4.3%) and carvone (4.0%) (Nickavar *et al.*, 2003). The use of diet supplemented with *Nigella sativa* seed oil improved growth performance, feed conversion efficiency and immune response of chickens (Omar *et al.*, 2002). Also, oral *Nigella sativa* L. treatment might decrease the diabetes-included disturbances of heart rate and some haematological parameters of alloxan-induced diabetic rabbits (Meral *et al.*, 2004).

In pumpkin seed (PS) oil, the saturated fatty acid content was 27.73% and comprises of 16.41% palmitic acid and 11.14% stearic acid. The unsaturated fatty acid value was 73.03% and consisting mainly of 18.14% oleic acid and 52.69% linoleic acid (Alfawaz, 2004). Up to 60.8%, of the pumpkin seed (PS) oil is

from the fatty acids, oleic (up to 46.9%), linolenic (up to 40.5%) and palmitic and stearic (up to 17.4%), the ratio of monounsaturated to polyunsaturated acids from 0.60 to 0.75 g (Nakiae *et al.*, 2006). The dietary supplementation with 5 g kg⁻¹ DM from PS oil in corn-soybean meal-wheat based diet can be profitable because it reduced broiler chickens mortality and it did not have any adverse effect on birds performance.

The objective of this study was to investigate the effect of pumpkin and black seeds oils on growth performance, milk yield and composition, blood parameters, immune response and reproductive performance of does rabbits as well as growth performance and immune response of their offspring during the suckling period.

2. Materials and methods

The current work was carried out at Sakha Animal Production Research Station, Animal Production Research Institute, Agricultural Research Center, Ministry of Agriculture during the period from January to May 2013.

Thirty six New Zealand white (NZW) doe rabbits with initial body weight of 1445.2±39.4 g and aged 12 weeks were used in a complete randomized design experiment with four dietary treatments. Does were fed commercial diet without additive (control, G1) or supplemented with 5 g pumpkin seed (PS) oil/kg diet (G2), 5 g black seed (BS) oil/kg diet (G3) and 2.5 g PS oil plus 2.5 g BS oil /kg diet (G4). Rabbits were fed commercial rabbit diet to cover their requirements according to NRC (1977). The formulation and chemical composition of commercial rabbit diet is shown in Table (1). Chemical analysis of diet was determined according to AOAC (1995).

Table 1. Formulation and composition of commercial rabbit diet

Ingredient	%	Composition	%
Berseem hay	30	DM	88.06
Wheat bran	16	OM	90.57
Soybean meal	20	CP	16.54
Yellow corn	20	CF	12.33
Barley grain	10	EE	2.25
molasses	2	NFE	59.45
limestone	1	Ash	9.43
Common salt	0.5		
Premix*	0.5		
Total	100		

* Each one kg of premix (minerals and vitamins mixture) contains vit. A, 20000 IU; vit. D3, 15000 IU; vit. E, 8.33 g; vit. K, 0.33 g; vit. B1, 0.33; vit. B2, 1.0 g; vit. B6, 0.33 g; vit. B5, 8.33 g; vit. B12, 1.7 mg; pantothenic acid, 3.33 g; biotinic, 33 mg; folic acid, 0.83 g; choline chloride, 200 mg.

Rabbits were housed separately in individual wired cages (30 x 50 x 30 cm) and arranged in double tier batteries allocated in two rows till insemination. After insemination, rabbits were placed again in individual cages (50 x 60 x 35 cm) of galvanized wire net and arranged in double tier batteries allocated in two rows. Nest boxes (30 x 25 x 30 cm) were attached to the front sides of the cages five days prior to kindling and removed at 28 day of lactation (weaning age). The bucks were housed in individual cages as that of females, but without nest boxes. All cages were equipped with feeders (made of galvanized steel sheets) and nipples (automatic drinkers). The system provided animals with fresh water all over the experimental period. Rabbits in all treatment groups were kept under similar managerial system and environmental conditions.

Doe rabbits were individually weighed at the beginning of the experiment and at weekly intervals during the experimental period. Does were naturally inseminated using four NZW bucks rabbit fed the control diet. Pregnancy diagnosis was done by palpation at 10-12 days post-mating and does failed to conceive post 1st mating were reinseminated and number of services per conception was calculated. During the suckling period, milk yield was determined by the differences in LBW before and after suckling and milk composition was determined using milkoscan (Model 133B).

Post-kindling, litter size and weight were recorded for each group at birth and at weekly intervals up to weaning at 28 days. Feed intake and weight gain during pregnancy and lactation and kindling rate of does as well as litter size and weights, stillbirths and mortality rate of bunnies were also recorded.

Blood samples were taken from does at 20 weeks of age and at 28 days of suckling period from does and their offspring. Blood samples were taken from ear vein into clean sterile tubes. Blood samples were let to coagulate and centrifuged at 3500 rpm for 15 minutes and serum was separated and stored at -20 °C till assay. Concentrations of total protein, albumin, globulin (by difference), glucose, total cholesterol, HDL, LDL (by difference), triglycerides, total lipids and activity of aspartate (AST) and alanine (ALT) aminotransferases as well as prolactin, T3 and T4 (one sample from each group) were determined in blood serum using spectrophotometer (Spectronic 21 DUSA) and commercial kits (Combination Pasteur Lap.). Also, concentration of immunoglobulins (IgG and IgM) was determined in serum of does and offspring using rabbit immunoglobulin M Elsa Kits.

Data were statistically analyzed using general linear models (GLM) procedures adapted by SPSS (2008) for user's guide with one-way ANOVA.

Duncan test within SPSS program was done to determine the degree of significance level among means.

3. Results and discussions

3.1. Growth performance of does:

Average of live body weight (LBW), daily gain (ADG) and total feed intake of does are presented in Table (2). Does in G4 showed significantly ($P<0.05$) the highest LBW and ADG at each age interval as compared to other groups versus the lowest values in G1. However, the differences in total feed intake were not significant.

3.2. Reproductive performance of does:

Reproductive traits of does rabbit are shown in Table (3). Results showed that does in G4 showed significantly ($P<0.05$) the highest LBW at mating and kindling, 1st service conception rate (CR) and the lowest number of services per conception, followed by those in G2 and G3, while G1 had the poorest values. However, gestation period length was nearly similar

for all treatment groups, ranging from 31.3 to 32.0 days.

3.3. Milk yield and composition:

Results in Table (4) showed that does in G4 recorded significantly ($P<0.05$) the highest average daily milk yield (ADMY) at different suckling intervals, followed by those in G2 and G3, while G1 had the lowest yield. Overall mean of daily milk yield during the suckling period significantly ($P<0.05$) increased by 9.57, 20.00 and 27.63% for G2, G3 and G4 compared with G1, respectively.

Milk composition of does including the percentages of fat, protein, lactose, total solids (TS), solids not fat (SNF) and ash are presented in Table (5). Results showed that diet of does in G4 significantly ($P<0.05$) improved milk composition in terms of the highest percentage of fat, protein, lactose, TS and SNF in milk. Does in G2 and G3 ranked the 2nd and 3rd groups, while G1 had the lowest percentages. However, ash percentage was not affected by supplementation. It is of interest to note that milk composition showed an opposite trend to milk yield.

Table 2: Live body weight and average daily gain of does rabbit at different age interval.

Age (week)	Experimental group				MSE
	G1	G2	G3	G4	
Body weight (g):					
12	1283.60 ^b	1455.00 ^{ab}	1450.00 ^{ab}	1592.10 ^a	39.44
16	1785.40 ^c	1980.80 ^{bc}	2021.80 ^{ab}	2191.80 ^a	42.64
20	2237.00 ^c	2454.00 ^d	2536.40 ^{ab}	2731.50 ^a	48.28
24	2658.50 ^c	2895.70 ^b	3016.70 ^{ab}	3235.20 ^a	55.21
Average daily gain (g):					
12-16	17.92 ^c	18.78 ^{bc}	20.42 ^{ab}	21.42 ^a	0.46
16-20	16.13 ^c	16.90 ^{bc}	18.38 ^{ab}	19.27 ^a	0.41
20-24	15.05 ^c	15.77 ^{bc}	17.15 ^{ab}	17.99 ^a	0.38
12-24	16.37 ^c	17.15 ^{bc}	18.65 ^{ab}	19.56 ^a	0.42
Total feed intake (kg)	13.59	13.60	13.67	12.99	0.19

a, b, c: Values in the same row with different superscripts differ significantly ($P<0.05$).

Table 3: Reproductive traits of does rabbit in experimental groups.

Item	Experimental group				MSE
	G1	G2	G3	G4	
No. of does	9	9	9	9	
LBW at mating (g)	2711.70 ^c	2953.60 ^b	3077.10 ^{ab}	3299.90 ^a	56.31
LBW at kindling (g)	2791.40 ^c	3040.40 ^b	3167.60 ^{ab}	3397.00 ^a	57.97
Conception rate %					
1 st service	66.67 ^b	77.78 ^{ab}	77.78 ^{ab}	88.89 ^a	6.93
2 nd service	22.22 ^a	22.22 ^a	22.22 ^a	11.11 ^b	2.69
3 rd service	11.11 ^a	0.00 ^b	0.00 ^b	0.00 ^b	1.50
No. service/conception	1.44 ^a	1.22 ^{ab}	1.22 ^{ab}	1.11 ^b	0.08
Gestation period (day)	32.00	31.43	31.71	31.29	0.17

a, b, c: Values in the same row with different superscripts differ significantly ($P<0.05$).

Table 4: Average daily milk yield (g/day) of does rabbit at different suckling intervals.

Suckling period (day)	Experimental groups				MSE
	G1	G2	G3	G4	
1	73.57 ^b	88.57 ^{ab}	96.57 ^a	100.71 ^a	6.57
7	88.57 ^c	105.00 ^b	117.86 ^a	122.14 ^a	7.57
14	149.29 ^b	151.43 ^b	168.57 ^a	170.71 ^a	9.59
21	159.29 ^c	178.57 ^b	182.86 ^b	212.86 ^a	12.03
28	118.57 ^b	122.14 ^b	139.29 ^a	145.71 ^a	7.11
Overall mean	117.86 ^c	129.14 ^b	141.43 ^{ab}	150.43 ^a	8.35

a, b, c, d: Values in the same row with different superscripts differ significantly (P<0.05).

Table 5: Milk composition of doe rabbits at different suckling periods.

Suckling period (day)	Experimental group				MSE
	G1	G2	G3	G4	
Fat%:					
1	16.63 ^b	17.33 ^{ab}	17.83 ^{ab}	18.33 ^a	0.26
7	14.97 ^b	15.80 ^{ab}	16.05 ^{ab}	16.50 ^a	0.24
14	14.47 ^b	15.08 ^{ab}	15.52 ^{ab}	15.95 ^a	0.23
21	14.14 ^b	14.73 ^{ab}	15.16 ^{ab}	15.58 ^a	0.22
28	14.64 ^b	15.25 ^{ab}	15.69 ^{ab}	16.13 ^a	0.23
Protein%:					
1	20.67 ^b	21.83 ^{ab}	22.17 ^{ab}	23.77 ^a	0.46
7	16.87 ^c	17.60 ^{bc}	18.93 ^{ab}	19.77 ^a	0.48
14	16.50 ^b	17.13 ^{ab}	17.77 ^{ab}	18.80 ^a	0.35
21	15.90 ^b	17.00 ^{ab}	17.70 ^{ab}	18.67 ^a	0.39
28	17.47 ^b	18.53 ^{ab}	18.73 ^{ab}	19.67 ^a	0.32
Lactose%:					
1	2.06 ^b	2.18 ^{ab}	2.20 ^{ab}	2.38 ^a	0.05
7	1.76 ^b	1.78 ^{ab}	1.81 ^{ab}	1.88 ^a	0.02
14	1.52 ^c	1.62 ^{bc}	1.72 ^{ab}	1.88 ^a	0.05
21	1.41 ^c	1.58 ^{bc}	1.64 ^{ab}	1.84 ^a	0.05
28	1.72 ^b	1.78 ^b	1.82 ^{ab}	1.98 ^a	0.04
Total solids%:					
1	41.41 ^b	43.42 ^{ab}	44.30 ^{ab}	46.60 ^a	0.62
7	35.35 ^c	36.75 ^{bc}	38.58 ^{ab}	39.95 ^a	0.65
14	34.17 ^c	35.53 ^{bc}	36.73 ^{ab}	38.73 ^a	0.55
21	33.09 ^c	34.97 ^{bc}	36.19 ^{ab}	37.79 ^a	0.50
28	35.54 ^c	37.29 ^b	38.00 ^{ab}	39.56 ^a	0.54
Solids not fat%:					
1	24.77 ^c	26.08 ^b	26.47 ^b	28.27 ^a	0.49
7	20.38 ^b	21.15 ^{ab}	22.53 ^{ab}	23.45 ^a	0.49
14	19.70 ^b	20.45 ^{ab}	21.21 ^{ab}	22.41 ^a	0.44
21	18.95 ^b	20.24 ^{ab}	21.03 ^{ab}	22.21 ^a	0.36
28	20.90 ^b	22.04 ^{ab}	22.31 ^{ab}	23.43 ^a	0.34
Ash%:					
1	2.05	2.07	2.10	2.12	0.03
7	1.75	1.76	1.79	1.80	0.03
14	1.68	1.70	1.72	1.74	0.03
21	1.64	1.66	1.69	1.70	0.03
28	1.71	1.73	1.75	1.77	0.03

a, b, c: Values in the same row with different superscripts differ significantly (P<0.05).

Table 6: Blood serum parameters of does in different groups.

Item	Experimental group				MSE
	G1	G2	G3	G4	
At 20 weeks of age:					
Total protein (g/dl)	6.92 ^b	7.72 ^a	6.76 ^b	7.99 ^a	0.33
Albumin (g/dl)	3.17 ^{ab}	3.51 ^a	3.35 ^{ab}	3.06 ^b	0.12
Globulin (g/dl)	3.75 ^c	4.21 ^b	4.07 ^{bc}	4.93 ^a	0.25
Glucose (g/dl)	47.75 ^c	63.48 ^a	57.74 ^b	59.50 ^{ab}	3.13
Total lipids (mg/dl)	282.87 ^a	261.77 ^{ab}	269.30 ^{ab}	248.59 ^b	19.22
Triglycerides (mg/dl)	40.80 ^a	38.67 ^{ab}	36.80 ^{ab}	35.87 ^b	2.58
Total cholesterol (mg/dl)	53.26 ^a	45.94 ^{ab}	43.06 ^b	43.04 ^b	4.07
HDL (mg/dl)	41.93 ^a	36.15 ^{ab}	34.12 ^{ab}	33.11 ^b	3.26
LDL (mg/dl)	11.33 ^a	9.78 ^{ab}	8.94 ^b	9.33 ^{ab}	1.31
Activity of AST (IU/ml)	38.42	38.37	37.09	38.41	0.47
Activity of ALT (IU/ml)	25.69	25.07	24.62	25.17	0.44
At 28 day of suckling period:					
Prolactin (ng/ml)	5.37	5.48	6.15	6.46	
T3 (nmol/l)	1.12	1.96	2.23	1.29	
T4 (nmol/l)	15.35	17.12	31.69	60.77	

a, b, c: Values in the same row with different superscripts differ significantly (P<0.05).

Table 7: Immunoglobulin concentration in blood serum of does and their offspring for the different groups at 28 days of suckling period.

Item	Experimental group			
	G1	G2	G3	G4
Does:				
IgG (ng/ml)	823	847	888	924
IgM (ng/ml)	328	422	455	589
Offspring:				
IgG (ng/ml)	977	1013	1023	1140
IgM (ng/ml)	387	433	478	605

Table 8: Litter size / doe and mortality rate (%) in different groups during suckling period.

Age (day)	Experimental group				MSE
	G1	G2	G3	G4	
Litter size:					
1	7.14 ^{ab}	6.71 ^b	7.57 ^a	7.29 ^{ab}	0.26
7	6.29 ^b	6.29 ^b	6.86 ^{ab}	7.29 ^a	0.25
14	6.29 ^b	6.00 ^b	6.86 ^{ab}	7.29 ^a	0.27
21	5.86 ^b	5.86 ^b	6.71 ^a	7.00 ^a	0.27
28	5.57 ^b	5.43 ^b	6.29 ^a	6.43 ^a	0.31
mortality rate (%):					
1-7	11.90 ^a	5.16 ^c	7.76 ^b	0.00 ^d	2.43
8-14	0.00 ^b	3.57 ^a	0.00 ^b	0.00 ^b	0.89
15-21	5.95 ^a	1.59 ^c	1.43 ^c	3.57 ^b	1.40
22-28	4.76 ^b	7.14 ^a	4.90 ^b	8.16 ^a	2.15
1-28	22.62 ^a	17.46 ^b	14.08 ^{bc}	11.73 ^c	3.66

a, b, c, d: Values in the same row with different superscripts differ significantly (P<0.05).

Table 9: Litter weight and weight gain in different groups during suckling period.

Age (day)	Experimental group				MSE
	G1	G2	G3	G4	
Litter weight (g):					
1	474.71 ^{ab}	535.86 ^a	437.43 ^b	461.43 ^{ab}	25.06
7	853.57 ^c	930.00 ^a	881.43 ^b	866.43 ^{bc}	48.95
14	1505.70 ^c	1508.60 ^c	1564.30 ^b	1597.90 ^a	74.12
21	1890.70 ^c	2038.60 ^b	2176.40 ^{ab}	2251.40 ^a	93.66
28	2319.30 ^c	2755.70 ^b	2845.50 ^{ab}	2912.90 ^a	101.39
Litter weight gain (g):					
1-7	378.86 ^b	394.14 ^{ab}	444.00 ^a	405.00 ^{ab}	48.20
8-14	652.14 ^b	578.57 ^c	682.86 ^{ab}	731.43 ^a	57.13
15-21	385.00 ^c	530.00 ^b	612.14 ^{ab}	653.57 ^a	62.22
22-28	428.57 ^c	717.14 ^a	669.10 ^b	661.50 ^b	55.34
1-28	1844.60 ^c	2219.90 ^b	2408.07 ^a	2451.47 ^a	104.68

a, b, c: Values in the same row with different superscripts differ significantly (P<0.05).

Table 10: Average live body weight and gain of bunnies produced by does in different groups during suckling period.

Age (day)	Experimental group				MSE
	G1	G2	G3	G4	
Bunnies weight (g):					
1	66.32 ^b	83.52 ^a	59.18 ^c	63.24 ^{bc}	4.20
7	138.72 ^{ab}	149.66 ^a	127.66 ^{bc}	120.51 ^c	7.02
14	241.76 ^{ab}	253.69 ^a	231.81 ^{bc}	222.95 ^c	10.90
21	324.20 ^b	352.20 ^a	328.70 ^b	322.16 ^b	10.27
28	433.70 ^c	518.22 ^a	452.38 ^b	453.02 ^b	18.46
Bunnies weight gain (g):					
1-7	72.40 ^a	66.14 ^a	68.48 ^a	57.28 ^b	6.53
8-14	103.04	104.04	104.15	102.44	9.12
15-21	82.44 ^b	98.50 ^a	96.89 ^a	99.20 ^a	7.61
22-28	109.51 ^c	166.03 ^a	123.68 ^b	130.86 ^b	16.07
1-28	367.38 ^c	434.70 ^a	393.20 ^b	389.78 ^b	17.39

a, b, c: Values in the same row with different superscripts differ significantly (P<0.05).

Table 11: Economic efficiency for the different groups.

Item	Experimental group				MSE
	G1	G2	G3	G4	
Total feed intake (kg)	13.59	13.60	13.67	12.99	0.19
Feed cost (L.E.)	33.98	33.99	34.16	32.47	0.49
Oil additives (g)	0.00	67.99	68.33	64.95	
Oil cost (L.E.)	0.00	6.12	4.10	4.87	
Total feed cost (L.E.)	33.98 ^b	40.11 ^a	38.26 ^a	37.34 ^a	0.67
TWG of does (g)	1374.90 ^c	1440.70 ^{bc}	1566.70 ^{ab}	1643.10 ^a	34.92
PTWG of does (L.E.)	27.50 ^c	28.81 ^b	31.33 ^{ab}	32.86 ^a	0.70
LW at weaning (g)	2319.30 ^c	2755.70 ^b	2845.50 ^{ab}	2912.90 ^a	101.39
PLW at weaning (L.E.)	46.39 ^c	55.11 ^b	56.91 ^{ab}	58.26 ^a	2.03
TR (L.E.)	73.89 ^c	83.92 ^b	88.24 ^{ab}	91.12 ^a	2.31
NR (L.E.)	39.91 ^c	43.81 ^{bc}	49.98 ^{ab}	53.78 ^a	2.11
NRI %	0.00 ^d	9.77 ^c	25.23 ^b	34.75 ^a	5.50

a, b, c, d: Values in the same row with different superscripts differ significantly (P<0.05).

3.4. Blood parameters:

Blood serum parameters of does shown in Table (6) revealed significant differences (P<0.05) in the concentrations of total protein (TP), albumin (AL), globulin (GL), total lipids (TL), triglycerides (TG), total cholesterol (TC), HDL and LDL. Does in G4 showed significantly (P<0.05) the highest concentration of TP and GL, while the highest AL concentration was detected significantly (P<0.05) in G2. Does in G2 showed significantly (P<0.05) the highest glucose concentration, while does in G1 showed significantly (P<0.05) the highest TL, TG, TC, HDL and LDL concentrations.

Activity of AST and ALT was nearly similar in different groups. concentration of prolactin was insignificantly the highest in G4 and its level in different groups confirmed the differences in milk yield (Table 4). However, the differences in concentration of T3 and T4 were not significant (Table 6).

3.5. Immune response of does and produced bunnies:

Immunoglobulin (IgG and IgM) concentration in blood serum of does and their offspring at 28 days of suckling period are presented in Table (7). Concentrations of IgG and IgM in serum of both does and their offspring were higher in G4, followed by G2 and G3, while they lower in G1, but the differences were not significant. Concentrations of both IgG and IgM increased in serum of offspring than of does. Concentrations of both IgG and IgM in serum of does was associated with globulin concentration in serum of does in all groups (Table 6).

3.6. Litter performance:

Results in Table (7) showed significant differences (P<0.05) in litter size and mortality rate of bunnies among groups during different suckling intervals. Does in G3 showed significantly (P<0.05) the highest litter size (LS) at birth, while those in G4

showed significantly ($P < 0.05$) the highest LS at 7, 14, 21 and 28 days of suckling period. Concerning the variations in mortality rate of bunnies among different groups, G1 showed significantly ($P < 0.05$) the highest mortality rate, followed by G2 and G3, while G4 had the lowest rate. Mortality rate of young rabbit during suckling period decreased with increased milk yield (Table 4) as well as with increasing immunoglobulin concentration in serum (Table 7).

Litter weight (LW) and their gain in different groups are shown in Table (9). Litter weight at birth was significantly ($P < 0.05$) the highest in G2, while G4 had the highest LW at 28 days of age. Litter weight at birth was significantly ($P < 0.05$) the lowest in G3, while G1 had the lowest ($P < 0.05$) LW thereafter until weaning. During the whole suckling period, G3 and G4 recorded significantly ($P < 0.05$) the highest LW gain, followed by G2, while G1 had the lowest gain.

3.7. Growth performance of bunnies of different groups:

Data in Table (10) showed that bunnies of G2 were significantly ($P < 0.05$) the heaviest weights and gains at different times of the suckling period, and this trend may be attributed to the lower litter size of does in this group.

3.8. Economic efficiency:

Results in Table (11) showed that feed cost was nearly similar for different groups. G4 showed significantly ($P < 0.05$) the highest total weight gain (TWG) of does, price of total weight gain (PTWG) of does, litter weight (LW) at weaning, price of litter weight (PLW) at weaning, total revenue (TR) and net revenue (NR), followed by G2 and G3, while G1 had the lowest values. Net revenue improvement for G2, G3 and G4 were 9.77, 25.23 and 34.75% compared with G1, respectively.

4. Discussion

Results of growth performance of doe rabbits agreed with those obtained by Awadalla and Kamal (2000), who found that addition of BS oil to the diet of rabbit improved growth performance. Also, Miraghaee *et al.* (2011) showed that supplementing 1% *Nigella sativa* improved ADG of broiler. Moreover, Nworgu (2007) reported that ADG of broiler chickens was significantly ($P < 0.05$) different, which was least in control as compared to birds served fluted pumpkin leaf extract.

The improvement of reproductive traits with dietary supplementation with PS and BS oils may be attributed to increasing their contents of unsaturated fatty acids mainly oleic and linoleic acids (Nickavar *et al.*, 2003; Alfawaz, 2004).

The significant enhancement in milk production was observed in goats fed diet supplemented with BS seeds (Vihan and Panwar, 1987). Lactating goats

supplemented with garlic, cinnamon or ginger oils, significantly increased ($P < 0.05$) milk yield (Kholif *et al.*, 2012). Also, Pascual *et al.* (2003) reported that during lactation, milk production was greater in animal fat and vegetable oil groups.

Generally, Rashwan (1998) working with doe rabbits, reported that milk production was improved as a result to supplementing doe diet with 12 g fenugreek or aniseed/ kg diet. Improvement of nutrition (by using feed additives, antibiotics, green fodder or natural sources) increase rabbit doe milk (Rashwan and Marai, 2000).

It is of interest to note that ADMY in all treatment groups increased gradually up to 21 days and decreased at 28 days of suckling period thereafter (Table 4). Similar trends of changes in milk yield of rabbits was observed by Omara *et al.* (2004).

Results of milk composition agreed with those obtained by Kowalska (2008), who indicated that milk fat increased from 18.4 to 22.2% in female rabbits receiving 3% fish oil compared to the control group. Kholif *et al.* (2012) found that garlic, cinnamon and ginger oils significantly increased ($P < 0.05$) protein, and solids not fat contents compared with the control. Santos *et al.* (2010) reported that milk lactose percent was increased by essential oils supplementation.

Results of blood parameters agreed with those obtained by Hassan *et al.* (2007) and Al-Beitawi *et al.* (2009), who recorded that *N. sativa* significantly decreased serum levels of TC and TG. Also, Hajati *et al.* (2011) and Miraghaee *et al.* (2011) found that dietary PS oil supplementation decreased TC and TG concentration in plasma and serum of broilers. Moreover, Sayed *et al.* (2007) found that T3 concentration in blood serum of Bouscat doe rabbit increased with fenugreek and aniseed additive.

Immunoglobulin (IgG and IgM) concentration in blood serum of does and their offspring agreed with those obtained by Khattab *et al.* (2011), who found that plasma immunoglobulin concentration of suckling buffalo calves was increased with supplementing 5 ml/head/day of BS oil.

Mortality rate of young rabbits is of vital importance in commercial rabbit farming, since it determines the net income of the rabbitries. Rashwan (1998) working with doe rabbits, reported that pre-weaning mortality was decreased as a results of supplementing doe diet with 12 g fenugreek or aniseed / kg diet. The increase of pre-weaning mortality associated the increase in litter size at birth and reduction of remating interval period, although doe milk yield appeared to be the most important factor in this respect. Thus, all factors which may decrease doe rabbit milk (nutrition, management of the rabbitry, climatic conditions and doe diseases), increase pre-weaning mortality (Rashwan and Marai, 2000). The

high mortality of young rabbits among primiparous rabbits was due to relatively low milk production. For this reason, the lower mortality of young rabbits may be due to higher fat content of milk rather than from the higher milk yield (Kowalska, 2008). Also, Sayed *et al.* (2007) found improvement in LS at 8 week after parturition to supplement the diet of Bouscat does and their offspring with a mixture of fenugreek and aniseed at the level of 0.5%.

Results litter weight agreed with those obtained by Rashwan (1998), who working with doe rabbits, reported that litter gain upon weaning improved as a result of supplementing doe diet with 12 g fenugreek or aniseed / kg diet. Also, Pascual *et al.* (2003) reported that growth of the litter during lactation was greater in group supplemented with animal fat and vegetable oil. Sayed *et al.* (2007) found improve litter weight at 8 week after parturition with a mixture of fenugreek and aniseed.

Growth performance of bunnies agreed with those obtained by Kowalska (2008), who found that fat supplemented diets increased rabbit kid weight at birth, 21 and 35 days of age.

Results of economic efficiency agreed with those obtained by El-Diahy (2004), who found that palm oil supplementation improved economic efficiency of milk production in Friesian cows.

Conclusion

Based on the foregoing results, it could be concluded that doe rabbits fed diets supplemented with a combination of pumpkin and black seeds oils (2.5 g PS oil plus 2.5 g BS oil/kg diet) showed the best results concerning growth performance, milk yield and composition, blood parameters, immune response and reproductive performance of does as well as the best results regarding litter size and weight, mortality rate and growth performance of their offspring and economic efficiency.

Correspondence to:

Prof. Dr. Hamed M.A. Gaafar
Animal Production Research Institute, Egypty.
hamedgaafar@gmail.com

References

1. Rashwan, A.A. and I.F.M. Marai (2000). Mortality in young rabbits: A Review. *World Rabbit Science*, 8(3): 111-124.
2. Nickavar, B.; F. Mojab; K. Javidnia and M.A.R. Amoli (2003). Chemical Composition of the Fixed and Volatile Oils of *Nigella sativa* L. from Iran. 0939D5075/2003/0900D0629 \$ 06.00” 2003 Verlag der Zeitschrift für Naturforschung, Tübingen. <http://www.znaturforsch.com.D>.
3. Omar, R.E.M.; E.A. Mahmoud; M.M. Karousa and Randa S.A. Ismaeil (2002). Effects of additives propolis and nigella sativa seed oil on some behavioural patterns, performance products and blood parameters in Sasso chickens. *Egypt. Poult. Sci.*, 21: 140-151.
4. Meral, I.; N. Donmez; B. Baydas; F. Belge and M. Kanter (2004). Effect of *Nigella sativa* L. on heart rate and some haematological values of alloxan-induced diabetic rabbits. *Scand. J. Lab. Anim. Sci.*, 31(1): 49-53.
5. Alfawaz, M.A. (2004). Chemical composition and oil characteristics of pumpkin (*Cucurbita maxima*) seed kernels. *Res. Bult.*, No. (129), Food Sci. & Agric. Res. Center, King Saud Univ., pp. 5-18.
6. Nakiaie, S.N.; D. Rade; D. Kevin; D. Štrucelj; Z. Mokrovaek and M. Bartoliae (2006). Chemical characteristics of oils from naked and husk seeds of *Cucurbita pepo* L. *Eur. J. Lipid Sci. Technol.*, 108: 936-943.
7. NRC (1977). *Nutrients Requirements of Domestic Animals. Nutrients Requirements of Rabbits*. 2nd Edition. National Research Council, National Academy of Science. Washington, DC. USA.
8. AOAC (1995). *Association of Official Analytical Chemists, 16th ed. Official Methods of Analysis*, Washington, DC, USA.
9. SPSS (2008). *Statistical package for the social sciences, Release 16, SPSS INC, Chicago, USA*.
10. Awadalla, S.A. and Azza M. Kamal (2000). Effect of *Nigella sativa* seeds oil on performance and some biochemical parameters in rabbit serum. *Egyptian Vet. Nutrition J.*, 1(1): 31-42.
11. Miraghaee, S.S.; B. Heidary; H. Almasi; A. Shabani; M. Elahi and M.H.M. Nia (2011). The effects of *Nigella sativa* powder (black seed) and *Echinacea purpurea* (L.) Moench extract on performance, some blood biochemical and hematological parameters in broiler chickens. *African Journal of Biotechnology*, 10(82): 19249-19254.
12. Nworgu, F.C. (2007). Economic importance and growth rate of broiler chickens served fluted pumpkin (*Telfaria occidentalis*) leaves extract. *African Journal of Biotechnology*, 6(2): 167-174.
13. Vihan, V.S. and H.S. Panwar (1987). Protective effect of *Nigella sativa* seeds on CC14-induced hepatotoxicity. *Indian Vet. J.*, 64: 347-349.
14. Kholif, S.M.; T.A. Morsy; M.M. Abdo; O.H. Matloup and A.A. Abu El-Ella (2012). Effect of supplementing lactating goat rations with garlic, cinnamon or ginger oils on milk yield, milk composition and milk fatty acids profile. *J. Life Sci.*, 4(1): 27-34.

15. Pascual, J.J.; C. Cervera; F. Quevedo and J. Fernandez (2003). Evolution of perirenal fat in primipara does fed with three high energy feeds. *World Rabbit Sci.*, 11: 247 (Abstract).
16. Rashwan, A.A. (1998). Effects of dietary additions of aniseed, fenugreek and caraway on reproductive and productive performance of New Zealand White rabbit does. *Egyptian J. Rabbit Sci.*, 8(2): 157-167.
17. Omara, M.E.; A.M.A. Mohi El-Din and M.A.M. Abu El-Hamd (2004). Effect of diets containing different types of silage on productive and reproductive performance of doe rabbits and growth performance of their bunnies. *J. Agric. Sci. Mansoura Univ.*, 29(10): 6301-6312.
18. Kowalska, D. (2008). Effect of essential unsaturated fatty acids in fish oil on litters and composition of milk of rabbit does. 9th World Rabbit Congress – June 10-13, 2008 – Verona – Italy, pp. 214 (Abstract).
19. Santos, M.B.; P.H. Robinson; P. Williams and R. Losa (2010). Effects of addition of an essential oil complex to the diet of lactating dairy cows on whole tract digestion of nutrients and productive performance. *Anim. Feed Sci. Technol.*, 157: 64-71.
20. Hassan, M.S.H.; A.M. AboTaleb; M.M. Wakwak and B.A. Yousef (2007). Productive, physiological and immunological effect of using some natural feed additives in Japanese quail. *Egypt. Poult. Sci.*, 27: 557-581.
21. AL-Beitawi, N.A.; S.S. EL-Ghousein and A.H. Nofal (2009). Replacing bacitracin methylene disalicylate by crushed *Nigella sativa* seeds in broiler rations and its effects on growth, blood constituents and immunity. *Livest. Sci.*, 125: 304-307.
22. Hajati, H.; A. Hasanabadi and P.W. Waldroup (2011). Effects of Dietary Supplementation with pumpkin oil (*Cucurbita pepo*) on performance and blood fat of broiler chickens during finisher period. *American Journal of Animal and Veterinary Sciences*, 6 (1): 40-44.
23. Sayed, M.A.M.; A.A. Azoz; A.A. El-Maqs and A.M. Abdel-Khalek (2007). Some performance aspects of doe rabbits fed diets supplemented with fenugreek and aniseed. EAAP, European Federation of Animal Science. Free Communications on Animal Nutrition. http://www.eaap.org/previous_Annual_Meetings/2007Dublin/sessions/session_20.htm.
24. Khattab, H.M.; A.Z. El-Basiony; S.M. Wakwak and B.A. Marwan (2011). Immune response and productive performance of dairy buffaloes and their offspring supplemented with black seed oil. *Iranian Journal of Applied Animal Science*, 1(4): 227-234.
25. El-Diahy, Y.M.M (2004). Effect of fat supplementation on productive and reproductive performance in lactating Friesian cows. M. Sc. Thesis, Fac. of Agric., Kafr El-Sheikh, Tanta University.

1/25/2014