

## Seasonal changes in physico-chemical properties of River sediments in Agbabu areas, Nigeria: Insight from principal component analysis

Akinyemi, S. A.<sup>1</sup>, Fagbote<sup>2</sup>, E.O., Adebayo<sup>1</sup>, O.F., Aturamu<sup>1</sup>, A. O., Gitari<sup>3</sup>, W. M., Olanipekun<sup>2</sup>, E. O., Ojo<sup>1</sup>, A. O.

<sup>1</sup>Department of Geology and Applied Geophysics, Ekiti State University, Ado Ekiti, Nigeria.

<sup>2</sup>Department of Chemistry, Ekiti State University, Ado-Ekiti, Nigeria.

<sup>3</sup>Department of Ecology and Resources Management, University of Venda, South Africa.

\*\*Corresponding author: Akinyemi, S. A.

E-mail: [segun.akinyemi@eksu.edu.ng](mailto:segun.akinyemi@eksu.edu.ng); Phone no: +2348106648373.

**Abstract:** The paper assesses the seasonal changes in the physico-chemical properties and the textural characteristic of the sediments for the duration of 2008 to 2009. The study appraises variations of physico-chemical parameters such as P, Mg<sup>2+</sup>, Ca<sup>2+</sup>, Na<sup>+</sup>, K<sup>+</sup>, pH, EC, TN, TOC and CEC during the dry and rainy seasons. Texturally, the sediment composed mostly of sand size particles with small aggregates of silt and clay particles. The silt and clay fractions of sediments were enriched during dry season but depleted during the rainy season. The pore water pH is slightly acidic with values varying between 4.42 – 6.74 and 4.36 – 6.20 during the dry and rainy seasons respectively. The mean pH values of 5.58 and 5.44 for the periods of dry and rainy seasons respectively fall below the World Health Organisation (WHO) threshold standards of 6.5 - 8.5. A significant increase in TOC, CEC, TN, Ca<sup>2+</sup>, Mg<sup>2+</sup>, Na<sup>+</sup>, K<sup>+</sup> during the dry season is attributed to the increase in the percentage of fine materials (i.e. clay and silt) and organic matter constituents in the sediments. Principal component analysis (PCA) revealed three extracted components namely comp. 1, comp. 2 and comp. 3, with TOC and the nature of sediment texture controlling the variation in the physico-chemical characteristics. The cluster analysis results corroborated the observed pattern in the principal component analysis.

[Akinyemi, S. A., Fagbote, E.O., Adebayo, O.F., Aturamu, A. O., Gitari, W. M., Olanipekun, E. O., Ojo, A. O.. **Seasonal changes in physico-chemical properties of River sediments in Agbabu areas, Nigeria: Insight from principal component analysis.** *World Rural Observ* 2016;8(3):77-84]. ISSN: 1944-6543 (Print); ISSN: 1944-6551 (Online). <http://www.sciencepub.net/rural>. 11. doi:[10.7537/marswro080316.11](https://doi.org/10.7537/marswro080316.11).

**Keywords:** Seasonal changes, physico-chemical properties, principal component analysis, cluster analysis.

### 1. Introduction

In the study areas, the problem of river water and sediment contamination has been of great concern to the rural dwellers. This is due to extent of deterioration of the environment and water bodies by human activities, mainly from on-loading and off-loading of petroleum products at the southern part of the study area. The indiscriminate discharge of sewage and solid wastes into the surroundings which eventually enter the natural water bodies could adversely affect the aquatic ecosystem. The supply ways of the metals entering the natural waters are governed by a dynamic set of physico-chemical influences while their solubility are mostly controlled by pH, concentration, nature of metal species, organic ligands, the oxidation condition of mineral components and the redox state of the aquatic system (Lala et al., 2008; Koffi et al., 2014). Metals were adsorbed onto both inorganic and organic particulates and are thus incorporated into sediment which led to elevated concentration of heavy metals in bottom sediments (Ochieng et al., 2007; Liu et al., 2009). Physico-chemical properties play an essential role in the evaluation of water quality. The accretion of metals from the water body to the sediment depends

on a number of external environmental factors such as pH, - (DO), - (EC) and the available surface area for adsorption due to the changes in grain size distribution (Davies et al., 2006). Nonetheless, metals are not continually fixed by sediments. Metals incorporated in the sediments may be re-activated and released back to waters. This takes pace through the changes of environmental conditions such as acidification, redox potential, and the organic ligand levels which adversely affect the living organisms (Liu et al., 2009). This research was carried out to illustrate the seasonal changes in the physico-chemical properties of bottom sediments using multivariate statistical analytical techniques. This will in turn reveal the principal factors responsible for seasonal variations in the physico-chemical properties of the studied bottom sediments.

### 2. Materials and methods

#### 2.1 Particle-size distribution

Size distribution analysis was carried out using Bouyoucous hydrometer methods as modified by Day (1965). Sediment samples were deflocculated with sodium hexametaphosphate solution (Calgon 44g/L) and sodium carbonate (8g/L). The pH of the solutions

were adjusted to about 8.3. The particle size distribution was determined using the textural triangular diagram.

### 2.2 Determination of pore water pH

The pH of the pore water in the sediments was determined by using 1:2.5 ratio of sediment to deionised water to form slurry. The measurement of pH was carried out using Corning pH meter Model 7 according to Jackson (1964).

### 2.3 Determination of total organic carbon

The total organic carbon in the sediments was determined by the wet combustion method of Walkey and Black (1934) as modified by Jackson (1964).

### 2.4 Exchangeable cations

Exchangeable  $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Ca}^{2+}$ , and  $\text{Mg}^{2+}$  were determined at neutral pH (i.e. pH 7) with ammonium acetate ( $\text{C}_2\text{H}_3\text{O}_2\text{NH}_4$ ) extraction method proposed by Thomas (1992). The filtrate was analysed for  $\text{Na}^+$  and  $\text{K}^+$  by photoelectric flame photometer, but  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  were determined by ethylene diamine tetra acetic (EDTA) titration method.

### 2.5 Available phosphorus

The available phosphorus in the sediment was measured using Bray and Kurt No. 1 methods as modified by Jackson (1964).

### 2.6 Total nitrogen

Total nitrogen in the sediment samples were determined using macro Kjeldahl's methods as described by Bremner (1965).

### 2.7 Statistical analyses

Descriptive statistical parameters such as mean, standard deviation were calculated along with Pearson's correlation coefficient. Multivariate statistics in term of dendrogram hierarchy cluster analysis (DHCA) and principal component analysis (PCA) were also carried out. PCA is an unverified configuration method with Varimax rotation and Kaiser normalisation. This is commonly used for dimension reduction of data and extraction of a small number of principal components. This was used for discovering the possible comparable pattern in the behaviour of metals and analyzing relationships among the observed various variables (Loska and Wiechula, 2003). Principal component analysis is consequently used to analyze relationships among the observed physico-chemical parameters and identify possible underlying factors for seasonal changes. This was carried out using the SPSS V17.0 for Windows. Dendrogram hierarchy cluster analysis (DHCA) combines all physico-chemical parameters together and could be used to ascertain similarities and differences between them. This could provide details to further verify the results of the PCA analysis. Dendrogram groups the elements as clustering variable

using Euclidian distance and Ward procedure (Laursen et al., 2014).

## 3. Results and discussion

### 3.1 Sediment texture analyses

Textural constituents of sediments varied among the five sampling points in the study areas (Table 1). The studied sediment composed mainly of sand with an admixture of silt and clay. Sand components range between 5.50 – 90.50% followed by clay fraction (6.18 – 59.25%) and silt constituents (3.23 – 46.10%). Sand fraction is higher in most of the studied samples during the rainy season but higher in silt and clay in the dry season. Nonetheless, OLWS4 has higher constituents of silt during rainy season. Different combinations of sediments texture observed are due to the transport of sediments from one place to another which are always associated with tidal currents (Satheshkumar and Khan, 2009). Correlation matrix revealed significant positive correlation of silt and clay components with all considered physico-chemical parameters during dry season but with the exception of pH during rainy season (Table 3).

### 3.2 Physico-chemical parameters of sediments

#### 3.2.1 Pore water pH

Statistical evaluation of the physico-chemical properties of the sediments revealed that the average pH varied between 4.42 – 6.74 during the dry season and 4.36 – 6.20 during the rainy season. Average pH of sediment for dry and rainy seasons were 5.58 and 5.44 respectively, with average pH of 5.51. This is slightly acidic but below WHO threshold standards of 6.5 - 8.5. Slight difference was recorded in pH levels during dry and rainy seasons (Fig. 1 & Table 2). The pH values were lower during the rainy season months than during the dry season. This is similar to research data available in literature (Jamabon, 2011; Daka and Moslen, 2013; Nwadinigwe et al., 2014). Commonly, the pH of bioleaching process of contaminated sediments is dependent on the buffering capacity, which determines metals to be released from sediments (Chen and Kin, 2001; Bartoli et al., 2012). Usually, variations in pH values during different seasons could be attributed to factors like removal of  $\text{CO}_2$  by photosynthesis through bicarbonate degradation, temperature and organic matter decay.

#### 3.2.2 Total organic carbon (TOC)

Average values of TOC ranged from 1.05 to 4.15% during the dry season and 0.07 to 2.24% during the rainy season (Fig. 1). The distribution of TOC closely followed the particle size distribution of the sediment. The sediment with low clay content also have low TOC and as the clay content increased, the TOC content also increased (Saravanakumar et al., 2008 and reference therein). In the present study, lower value of TOC was obtained during rainy season

than during dry season for samples code-named AGM2, OLSW3 and OLSW4. Average TOC of sediment for dry and rainy seasons were 2.84 and 1.20 respectively, with average TOC of 2.02 (Table 2). Correlation analysis revealed significant positive correlation of TOC with pH during dry season (Table 3). There is a significant difference in TOC levels of sediments during the dry and rainy seasons. An increase in TOC during the dry season may be ascribed to the increased percentage of finer materials in the sediments (clayey and silt sediments), temperature and fast organic matter decay.

### 3.2.3 Total nitrogen (TN)

Average TN in the sediments varied between 0.12 – 0.37% during the dry season and 0.03 – 0.21% during the rainy season. The variation in the concentration of TN is influenced by the textural nature of the sediments (i.e. percentage of fines). Table 2 shows lower value of TN during the rainy season than the dry season for samples code-named AGM2, OLSW3 and OLSW4. Average total nitrogen obtained for the dry and rainy seasons were 0.24 and 0.13 respectively, with average TN of 0.18 (Table 2). Total nitrogen showed a significant positive correlation with pH and TOC during the dry season but only correlate with TOC during the rainy season (Table 3). The significant difference in total nitrogen levels during the dry and rainy season is perhaps due to the oxidation of dead plant organic matter in the sediments. The lower value of total nitrogen recorded during the rainy season may be ascribed to low quantity of organic matter and low quantity of finer materials which is caused by the winnowing action of waves. Dye (1978) observed high TN content in finer materials and attributed it to trapping of detritus by finer particles, which usually lead to increase in bacterial population. This may also be a reason for the high level of total nitrogen encountered in the present study.

### 3.2.4 Cation exchange capacity (CEC)

Average cation exchange capacity varied between 10.98 – 24.03 cmol<sub>c</sub>kg<sup>-1</sup> during the dry season and 9.63 – 20.98% during the rainy season. Table 2 shows low cation exchange capacity during the rainy season and high during the dry season for samples code-named AGM2, OLSW3 and OLSW4. The average cation exchange capacity of sediments for the dry and rainy seasons were 18.91 cmol<sub>c</sub>kg<sup>-1</sup> and 16.25 cmol<sub>c</sub>kg<sup>-1</sup> respectively, with average CEC of 17.58 cmol<sub>c</sub>kg<sup>-1</sup> (Table 2). Cation exchange capacity shows strong correlation with all physico-chemical parameters but does not correlate with pH during rainy season (Table 3). There is a significant difference in CEC levels during the dry and rainy seasons. This is ascribed to the differences in the quantity of finer materials and organic matter in the sediments. Cation

exchange capacity commonly increases as grain size decreases, but this relation is not true for many stream sands and silts that contain important percentages of rock fragments. Instead, a minimum CEC is measured in the fine-sand to coarse silt range where the fewest clay aggregates occur (Kennedy, 1965).

### 3.2.5 Total phosphorous (TP)

Total inorganic phosphorous ranged from 5.45 to 23.88 μgg<sup>-1</sup> during the dry season and 4.30 to 24.83 μgg<sup>-1</sup> during the rainy season. Figure 1 shows high value of total phosphorous during the rainy season for samples code-named OLSW1, OLSW2 and OLSW4. Similarly, high total phosphorous was recorded during the dry season for samples code-named AGM2 and OLSW3. The average total phosphorous present in the sediments during the dry and rainy seasons were 16.00 μgg<sup>-1</sup> and 13.23 μgg<sup>-1</sup> respectively, with average total value of 14.61 μgg<sup>-1</sup> (Table 2).

Total phosphorous correlates with all considered physico-chemical parameters during the dry season but do not correlate with only pH during the rainy season (Table 3). The higher values observed during the dry season may be ascribed to dead organic matter from the top layer. Whereas lower values obtained during the rainy season may be related to the removal of top layer of sediments by heavy floods (Saravanakumar et al., 2008).

### 3.2.6 Potassium and sodium

Potassium ranged from 0.68 – 2.30 cmol<sub>c</sub>kg<sup>-1</sup> during dry season and 0.08 – 3.24 cmol<sub>c</sub>kg<sup>-1</sup> during rainy season. The amount of potassium was found to be fluctuating during the period of investigation. The average values of potassium in the sediments during the dry and rainy seasons were 1.50 cmol<sub>c</sub>kg<sup>-1</sup> and 1.32 cmol<sub>c</sub>kg<sup>-1</sup> respectively, with average value of 1.41 cmol<sub>c</sub>kg<sup>-1</sup>. The maximum value of potassium was obtained in sample code-named OLSW4 during the rainy season. Sodium varies between 0.25 – 1.25 cmol<sub>c</sub>kg<sup>-1</sup> during the dry season and 0.08 – 1.27 cmol<sub>c</sub>kg<sup>-1</sup> during the rainy season. Average values of sodium for the dry and rainy seasons are 0.77 cmol<sub>c</sub>kg<sup>-1</sup> and 0.64 cmol<sub>c</sub>kg<sup>-1</sup> respectively, with average value of 0.70 cmol<sub>c</sub>kg<sup>-1</sup>. The maximum value of sodium was observed in the sample code-named OLSW4 during the rainy season (Figure 1). Sodium and potassium ions showed significant positive correlations with pH, TOC, TN and P during the dry season but they did not correlate with pH during the rainy season (Table 3).

### 3.2.7 Calcium, magnesium and manganese

Calcium varied between 2.65 – 7.48 cmol<sub>c</sub>kg<sup>-1</sup> during dry season and 1.02 – 5.49 cmol<sub>c</sub>kg<sup>-1</sup> during rainy season. The average value of calcium in the sediment during the dry and rainy seasons were 5.37 cmol<sub>c</sub>kg<sup>-1</sup> and 3.68 cmol<sub>c</sub>kg<sup>-1</sup> respectively, with

average value of calcium which is  $4.52\text{cmol}_e\text{kg}^{-1}$ . The maximum value of calcium was obtained in samples code-named OLWS3 during dry season. Magnesium range from  $1.55 - 4.38\text{cmol}_e\text{kg}^{-1}$  during dry season and  $1.29 - 5.66\text{cmol}_e\text{kg}^{-1}$  during rainy season. Average magnesium values for the dry and rainy seasons were  $3.04\text{cmol}_e\text{kg}^{-1}$  and  $2.82\text{cmol}_e\text{kg}^{-1}$  respectively, with average value of  $2.93\text{cmol}_e\text{kg}^{-1}$ .

The maximum concentration of magnesium was obtained in sample code-named AGM2 during the rainy season (Figure 1). Calcium and magnesium ions have significant positive correlation with all physico-chemical parameters during the dry season but they did not correlate with only pH during the rainy season. Manganese ranged between  $0.07 - 0.30\text{cmol}_e\text{kg}^{-1}$  during the dry season and  $0.02 - 2.62\text{cmol}_e\text{kg}^{-1}$  during the rainy season. Average manganese values for the dry and rainy seasons were  $0.19\text{cmol}_e\text{kg}^{-1}$  and  $0.54\text{cmol}_e\text{kg}^{-1}$  respectively, with average value of  $0.37\text{cmol}_e\text{kg}^{-1}$ . The concentration maxima for manganese was obtained in the sample code-named OLSW3 during rainy season (Figure 1). Manganese ions strongly correlate with all physico-chemical parameters during the dry season but did not correlate with only pH during the rainy season (Table 3).

#### 4. Principal Component Analysis of seasonal variation in sediments

The results of output of principal component analysis of sediments collected during the dry and

rainy seasons using SPSS version 17 is shown in Table 4. Two components were extracted during the dry season which suggests insufficient physical mixing of sediments. Component 1 contributes 83.86% of the data total variation and comprises of strong positive loadings of silt,  $\text{Mg}^{2+}$ ,  $\text{Ca}^{2+}$ ,  $\text{Mn}^{2+}$ ,  $\text{K}^+$ , TOC, TN, CEC and clay which suggest surface area as the controlling factor. Component 2 contains 15.04% of the data total variation and shows strong positive loadings of pH and sand materials which indicate the pH of the interstitial pore as the controlling factor. Three components were extracted from sediment samples collected during the rainy season. Components 1 showed strong loadings of P,  $\text{K}^+$ , TN, TOC, silt and clay and its represents 64.86% of the total data variation indicating organic matter constituents as the principal factor. Components 2 contains strong loadings P,  $\text{K}^+$ ,  $\text{Na}^+$ ,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$  and CEC and contributes about 18.78% to the total data variation suggesting surface area of the finer component of river sediments.

Components 3 showed positive loadings of P,  $\text{Na}^+$ ,  $\text{Mn}^{2+}$ ,  $\text{Ca}^{2+}$ , TN and CEC and represents 9.97% of the total data variations. This indicated that the data variations are partly controlled by organic matter and surface area. Dendrogram cluster analysis corroborated the principal component analysis results by showing two cluster groups; Cluster 1 consists of  $\text{Mg}^{2+}$ ,  $\text{Ca}^{2+}$ ,  $\text{Na}^+$ ,  $\text{K}^+$ , P, TN, TOC, CEC, silt and clay whereas cluster group 2 comprises of pH and sand (Figure 2).

Table 1: Seasonal variations of sediment composition at the studied areas ( $n = 3$ )

Dry season							
Sample	Sand (%)		Silt (%)		Clay (%)		Textural class
	Mean	Stdev.	Mean	Stdev.	Mean	Stdev.	
AGM2	11.50	0.29	30.00	0.00	58.25	0.25	Silty clay
OLSW1	59.00	0.00	3.75	0.25	37.50	0.29	Sandy clay
OLSW2	59.75	0.25	4.00	0.00	36.50	0.29	Sandy clay
OLSW3	5.50	0.29	35.50	0.29	58.75	0.47	Clay
OLSW4	5.50	0.29	35.00	0.00	59.25	0.03	Clay
OKCtr	NA						
Rainy season							
Sample	Sand (%)		Silt (%)		Clay (%)		Textural class
	Mean	Stdev.	Mean	Stdev.	Mean	Stdev.	
AGM2	75.95	0.10	4.10	0.06	20.20	0.08	Silty clay
OLSW1	60.20	0.12	3.23	0.22	36.78	0.13	Sandy clay
OLSW2	59.50	0.15	3.43	0.10	36.75	0.19	Sandy clay
OLSW3	75.54	0.24	4.13	0.05	20.08	0.04	Clay
OLSW4	11.45	0.05	46.10	0.04	42.53	0.03	Clay
OKCtr	90.50	0.29	3.98	0.06	6.18	0.10	NA

Table 2: Average concentration of physico-chemical parameters for dry and rainy seasons in sediment of the study area

Variables	Mean for dry season	Mean for rainy season	Average for dry and rainy season
pH	5.58	5.44	5.51
TOC (%)	2.84	1.20	2.02
TN (%)	0.24	0.13	0.18
P ( $\mu\text{g/g}$ )	16.00	13.23	14.61
CEC( $\text{cmol}_c\text{kg}^{-1}$ )	18.91	16.25	17.58
$\text{K}^+$ ( $\text{cmol}_c\text{kg}^{-1}$ )	1.50	1.32	1.41
$\text{Na}^+$ ( $\text{cmol}_c\text{kg}^{-1}$ )	0.77	0.64	0.70
$\text{Ca}^{2+}$ ( $\text{cmol}_c\text{kg}^{-1}$ )	5.37	3.68	4.52
$\text{Mg}^{2+}$ ( $\text{cmol}_c\text{kg}^{-1}$ )	3.04	2.82	2.93
$\text{Mn}^{2+}$ ( $\text{cmol}_c\text{kg}^{-1}$ )	0.19	0.54	0.37

Table 3: Pearson's correlation matrix of sediment samples during dry and rainy seasons

	pH	TOC	TN	P	$\text{K}^+$	$\text{Na}^+$	$\text{Ca}^{2+}$	$\text{Mg}^{2+}$	$\text{Mn}^{2+}$	CEC	Sand	Silt	Clay
pH	1												
TOC	0.276	1											
TN	0.346	.980**	1										
P	0.259	.999**	.974**	1									
$\text{K}^+$	0.331	.961**	.963**	.955**	1								
$\text{Na}^+$	0.221	.978**	.966**	.975**	.990**	1							
$\text{Ca}^{2+}$	0.371	.994**	.989**	.991**	.957**	.963**	1						
$\text{Mg}^{2+}$	0.381	.988**	.992**	.983**	.982**	.979**	.994**	1					
$\text{Mn}^{2+}$	0.294	.989**	.975**	.984**	.986**	.993**	.981**	.990**	1				
CEC	0.522	.960**	.968**	.953**	.937**	.924**	.982**	.979**	.958**	1			
Sand	.730*	-0.452	-0.363	-0.47	-0.36	-0.478	-0.36	-0.34	-0.421	-0.195	1		
Silt	0.144	.991**	.963**	.992**	.949**	.981**	.971**	.966**	.979**	.915**	-0.566	1	
Clay	0.64	.915**	.931**	.907**	.902**	.871*	.951**	.949**	.911**	.987**	-0.055	.852*	1
pH	1												
TOC	-0.678	1											
TN	-0.711	.948**	1										
P	-0.978**	.738*	0.712	1									
$\text{K}^+$	-0.725	.808*	0.647	.841*	1								
$\text{Na}^+$	-0.982**	0.686	0.674	.992**	.778*	1							
$\text{Ca}^{2+}$	-0.833*	0.65	0.555	.912**	.824*	.919**	1						
$\text{Mg}^{2+}$	-0.379	0.434	0.265	0.54	.753*	0.497	.750*	1					
$\text{Mn}^{2+}$	-0.643	0.072	0.236	0.54	0.023	0.634	0.516	0.007	1				
CEC	-0.786*	.735*	0.682	.870*	.854*	.842*	.918**	.834*	0.372	1			
Sand	0.551	-.900**	-.816*	-0.597	-.751*	-0.523	-0.41	-0.207	0.186	-0.47	1		
Silt	-0.65	.732*	0.639	0.685	.809*	0.606	0.448	0.265	-0.133	0.488	-.900**	1	
Clay	-0.268	.855*	.804*	0.316	0.471	0.267	0.24	0.079	-0.216	0.313	-.848*	0.531	1

Correlation(*r*) are significant at the 0.05 level (2-tailed).

Table 4: Principal Component Analysis of sediments showing variations in physico-chemical parameters during dry and rainy seasons

Variables	Rainy season			Communalities	Dry season		
	Comp. 1	Comp. 2	Comp. 3		Comp. 1	Comp. 2	Communalities
Sand	-0.97			0.98		0.92	0.99
Clay	0.92			0.85	0.94		0.99
TOC	0.87			0.96	0.99		0.99
TN	0.83		0.42	0.89	0.99		0.98
Silt	0.79			0.74	0.98		0.99
Mg <sup>2+</sup>		0.98		0.97	0.99		0.99
CEC		0.78	0.48	0.94	0.97		0.99
K <sup>+</sup>	0.58	0.77		0.97	0.98		0.96
Ca <sup>2+</sup>		0.72	0.61	0.95	0.99		0.99
Mn <sup>2+</sup>			0.96	0.98	0.99		0.99
pH	-0.42		-0.81	0.97		0.94	0.99
Na <sup>+</sup>		0.481	0.78	0.99	0.98		0.98
P	0.44	0.53	0.71	0.98	0.99		0.99
EV	8.43	2.44	1.30		10.90	1.96	
VAR (%)	64.86	18.78	9.97		83.86	15.04	
CVAR (%)	64.86	83.64	93.61		83.86	98.90	

EV = Eigen value, VAR (%) = variance percentage and CVAR (%) = cumulative variance.





Fig. 1: Seasonal changes of selected physico-chemical parameter levels in the sediments through dry and rainy seasons.

## Dendrogram using Ward Method



## Dendrogram using Ward Method




Figure 2: Dendrogram cluster analysis of sediments during dry and rainy seasons.

## 5. Conclusions

Investigative study of the seasonal changes on the physico-chemical characteristics of the sediments in the studied areas revealed that the sediments were mostly composed of sand sized particles with finer materials (i.e. silt and clay). The sand constituents of investigated sediments were higher during the rainy season but the sediments were enriched in silt and clay fractions during the dry season. The pH values of

studied sediments were slightly acidic but fall below WHO threshold standards of 6.5 - 8.5. Insignificant differences in pH levels were observed in the sediments during the dry and rainy seasons.

A significant increase in TOC, CEC, TN, Ca<sup>2+</sup>, Mg<sup>2+</sup>, Na<sup>+</sup>, K<sup>+</sup> during the dry season is attributed to increase in the percentage of finer materials (i.e. clay and silt) and organic matter in the sediments. Principal component analysis revealed that the TOC and

percentage of fine materials are controlling the varying level of physico-chemical parameters. The cluster groups verified the observed pattern in the principal component analysis.

## 6. Acknowledgements

We acknowledge Mr. Uyi Hanson, Rivers State University of Science and Technology, Port Harcourt and Miss Kate Iwuzor of Fugro Laboratories, Port Harcourt for their technical assistance.

## References

- Bremmer, J. M. (1965). Total Nitrogen in: Methods of Soil Analysis. Part 2 (C A Black *et al* ed.) American Society of Agronomy Inc. Madison, Wisconsin U.S.A, pp 1149-1178.
- Daka, E. R. and Moslen, M. (2013). Spatial and Temporal Variation of Physico-Chemical Parameters of Sediment from Azuabie Creek of the Upper Bonny Estuary, Niger Delta. *Research Journal of Environmental and Earth Sciences*, 5(4): 219-228.
- Davies, O. A., Allison, M. E. & Uyi, H. S. (2006). Bioaccumulation of heavy metals in water, sediment and periwinkle (*Tympanoonusfuccatus* var *radula*) from the Elechi creek, Niger Delta. *African journal of Biotechnology*, 5: 968-973.
- Day, P. R. (1965). Particle Fractionation and Particle Size Analysis in: Methods of Soil Analysis, Part 1 (C A Black *et al* ed.) American Society of Agronomy Inc. Madison, Wisconsin U.S.A, pp 545-567.
- Dye, A. H. (1978). An ecophysiological study of the meiofauna of the Swartkops estuary, I. The sampling sites: Physical and chemical features. *Zool. Afr.*, 13: 1-18.
- Jackson, M. L. (1964). *Soil Chemical Analyses*. Englewood Cliffs. New York, Printice Hall.
- Jamabon, A. (2011). Sediment characteristics of the mangrove swamps of the upper Bonny River, Niger Delta, Nigeria. *New Clues in Sciences*, 1: 74-79.
- Kennedy, V. C. (1965). Mineralogy and Cation-Exchange Capacity of Sediments from Selected Streams. Geological survey professional paper, 433-D: D1-D28.
- Koffi, K. M., SafiatouCoulibaly, S., Atse, B. C., Kouamelan, E. P. (2014). Survey of heavy metals concentrations in water and sediments of the estuary Bietri bay, Ebrie lagoon, Cote D'ivoire. *International Journal of Research in Earth & Environmental Sciences*, 1 (3): 1-10.
- Lalah, J. O., Ochieng, E. Z. & Wandiga, S. O. (2008). Sources of heavy metal input into Winam Gulf, Kenya. *Bulletin of Environmental Contamination and Toxicology*, 81: 27-284.
- Laursen, J., Milman, N., Pind, N., Pedersen, H., Mulvad, G. (2014). The association between content of the elements S, Cl, K, Fe, Cu, Zn and Br in normal and cirrhotic liver tissue from Danes and Greenlandic Inuit examined by dual hierarchical clustering analysis. *J. Trace Elem. Med. Biol.*, 28: 50-55.
- Liu, C., Xu, J., Liu, C., Zhang, P., & Dai, M. (2009). Heavy Metals in the Surface Sediments in Lanzhou Reach of Yellow River, China. *Bulletin of Environmental Contamination and Toxicology*, 82: 26-30.
- Loska, K., Wiechula, D. (2003). Application of principal component analysis for the estimation of source of heavy metal contamination in surface sediments from the Rybnik Reservoir. *Chemosphere*, 51: 723-733.
- Nwadinigwe, C. A., Udo, G. J., Nwadinigwe, A. O. (2014). Seasonal Variations of Heavy Metals Concentrations in Sediment Samples Around Major Tributaries in Ibeno Coastal Area, Niger Delta, Nigeria. *International Journal of Scientific & Technology Research*, volume 3 issue 11: 254-265.
- Ochieng, E. Z., Lalah, J.O. & Wandiga, S.O. (2007). Analysis of heavy metals in water and surface sediment in five Rift Valley Lakes in Kenya for assessment of recent increase in anthropogenic activities. *Bulletin of Environmental Contamination and Toxicology*, 79: 570-576.
- Saravanakumar, A., Rajkumar, M., Serebiah, J. S. and Thivakaran, G. A. (2008). Seasonal variations in physico-chemical characteristics of water, sediment and soil texture in arid zone mangroves of Kachchh-Gujarat. *Journal of Environmental Biology*, 29(5): 725-732.
- Satheeshkumar, P., Khan, B. A. (2009). Seasonal Variations in Physico-Chemical Parameters of Water and Sediment Characteristics of Pondicherry Mangroves. *African Journal of Basic & Applied Sciences*, 1 (1-2): 36-43.
- Thomas, G W. (1992). Exchangeable Cations. In: A L Page, R. H. Miler and D. R. Keeney (Editors). *Methods of Soil Analysis*, Agronomy Series 9, ASA and SSSA, Madison, WI pp 159-161. Walkey, A and I A Black (1934) Determination of Organic Carbon in Soil. *Soil Science*, 37: 29-238.