

Discussion of Stem Cell Totipotent and Differentiation

Ma Hongbao *, Yang Yan *, Margaret Ma **

* Brookdale Hospital, Brooklyn, New York 11212, USA, ma8080@gmail.com

** Boston, Massachusetts 02138, USA

Abstract: The totipotent of stem cell is the most important property for the stem cell to keep the life continuously. All the cells can develop into an entire alive animal body that is same as the baby comes from the fertilized egg – the zygote. It is important to reverse the non-totipotent cell to the totipotent, which is the lifeline reverse that make the animal growth from old to younger.

[Ma H, Yang Y, Ma M. **Discussion of Stem Cell Totipotent and Differentiation.** *Stem Cell* 2011;2(1):20-24] (ISSN 1545-4570). <http://www.sciencepub.net/stem>. 3

Key words: DNA; life; stem cell; totipotency

The totipotent of stem cell is the most important property for the stem cell to keep the life continuously. The totipotency is the ability that a cell can divide and produce a new organism. A single cell has an integrate genes to differentiate to a whole living body (Baud, 2005). Totipotent cells can be any cell in a body (Cantley, 2005). Normally, the differentiation is one direction where an undifferentiated cell (especially stem cell) can differentiate into differentiated cells but differentiated cells cannot reverse to the immatured cells. However, under certain condition, this direction can reverse, especially in plant and low class of the life species. A plant cutting or callus can grow to an entire plant and this appears everywhere in the earth. Many plants reproduce for next generation through schedule and it is widely used in the agriculture.

The Pluripotent stem cells cannot develop into a fetal or adult organism alone because they are lack of the potential to contribute to extraembryonic tissue, such as the placenta. After the embryonic development stage is over, the stem cells no longer have this unlimited potential to develop into all cell types, the pluripotency is lost and they can only become certain types of cells.

Induced pluripotent stem cells (iPSCs) are genetically reprogrammed adult cells that exhibit a pluripotent stem cell-like state similar to embryonic stem cells. While these artificially generated cells are not known to exist in the human body, they show qualities remarkably similar to those of embryonic stem cells (ESCs); thus, iPSCs are an invaluable resource for drug discovery, cell therapy, and basic research.

An alive animal body begins from a zygote. The fertilized egg zygote is a totipotent cell, which is the beginning of an animal body, including human. A human begins a zygote that a sperm fertilizes an egg and creates the single totipotent cell. In the beginning

hours after fertilization of the human, this cell divides into the identical totipotent cells, which can later develop into any of the 3 germ layers of a human (endoderm, mesoderm, or ectoderm) and into cells of the cytotrophoblast layer or syncytiotrophoblast layer of the placenta. After reaching the 16-cell stage, the totipotent cells of the morula differentiate into cells. The differentiated cells will become either the blastocyst's inner cell mass or the outer trophoblasts. Approximately 4 days after the fertilization and a few cell divisions, these totipotent cells will be specialized, then lose the totipotent ability.

Totipotent is the most important point for the stem cells that initiates the whole creature body. In creature development, the egg cell in a lady and the sperm from a man fuses jointly to become a free chamber zygote. The zygote divides many times and forms cells that are the precursors to the trillions of cells as the bricks of the creature body.

Totipotent stem cell can differentiate into all the human body's cells. In most animals, the only true totipotent stem cell is the fertilized egg and its immediate descendants. A totipotent stem cell can potentially generate a complete organism. In some cases, cells can regain totipotency. A plant cutting or callus can be used to grow an entire plant. The plant is different from animal in this point. The totipotency regaining is a potential ability for the life to reverse the lifeline.

Differentiation results from differential gene expression. In order to clone an animal, such as a sheep or a mouse or a rat, udder cells are removed from a ewe and starved for one week to cause G0 arrest. Nuclei from arrested ewe udder cells are fused with enucleated eggs from a ewe, and then stimulated to re-enter the cell cycle. After a few rounds of cell division, the embryo is transplanted into a surrogate sheep mother. The sheep that is born is genetically identical to the ewe from that the nucleus obtains.

Clone technique is an important technique in modern biotechnology.

Cellular determination is important in the animal development. Cellular determination results from the asymmetric segregation of cellular determinants. However, normally determination is the result of inductive signaling between cells. Asymmetric segregation of cellular determinants is caused from the asymmetric localization of cytoplasmic molecules within the cell before dividing. During cell division procedure, one daughter cell receives more localized molecules and the other daughter cell may receive less of these molecules, which results in two different daughter cells taking on different cell fates based on differences in gene expression. The different gene differentiation comes from different gene expression factors that make the genes' differentiate under different condition. The localized cytoplasmic determinants are often transcription factors or mRNAs encoding by the transcription factors.

The field of stem cell biology has undergone tremendous expansion over the past years. Scientific investigation has continued to expand our understanding of these complex cells at a rapidly increasing rate. This understanding has produced a vast array of potential clinical applications (Hemmat, Lieberman et al. 2010).

Primordial germ cells (PGCs), the precursors of sperm and eggs, are the route to totipotency and require establishment of a unique epigenome in this lineage. The genetic program for PGC specification in the mouse also initiates epigenetic reprogramming that continues when PGCs migrate into the developing gonads. Among these later events is active and genome-wide DNA demethylation, which is linked to extensive chromatin remodeling (Surani and Hajkova 2010).

In many tissues, mammalian aging is associated with a decline in the replicative and functional capacity of somatic stem cells and other self-renewing compartments. Understanding the basis of this decline is a major goal of aging research (Sharpless 2010).

The octamer-binding transcription factor 4 (Oct-4) plays important role in totipotent cells differentiation. Oct-4 expressed in totipotent embryonic cells and germ cells. As totipotent cells differentiate somatic cells, Oct-4 gene is downregulated. Oct-4 expression is maintained after postgastrulation in primordial germ cells. Oct-4 is necessary for embryonic cells to keep the totipotent status keep from their differentiation to somatic cells. (Pesce M, Schöler HR., 2000).

Example of the totipotent protocols - Proliferation of totipotent hematopoietic stem cells in

vitro with retention of long-term competitive in vivo reconstituting ability (Christopher, 1992):

Recombinant Tkneo9 virus at a titer of 1×10^6 per ml is generated from a T-2 producer line maintained in 10% calf serum/Dulbecco's modified Eagle's medium. Marrow cells from adult male (C57BL/6J x C3H/HeJ) F1 (B6C3F1) mice injected i.v. 4 days earlier with 5-fluorouracil (5-FU, 150 mg/kg of body weight) are infected with Tkneo9 virus using a supernatant infection protocol in which $3-5 \times 10^6$ marrow cells are cultured for 24 hr in 100-mm Petri dishes in 10 ml of virus supernatant containing Polybrene at 4 μ g/ml, 5% pokeweed mitogen-stimulated spleen cell-conditioned medium, and 10% agar-stimulated human leukocyte-conditioned medium as described. Cells are then recovered by gentle agitation and scraping of dishes with a rubber policeman, essential Eagle medium and resuspended in LTC medium [a medium/10% horse serum/10% fetal calf serum/ 10^{-6} M sodium hydrocortisone hemisuccinate/ 10^{-4} M 2-mercaptoethanol]. Aliquots of 3×10^6 cells are overlaid on previously established 3-wk-old long-term B6C3F1 female marrow adherent layers that had been irradiated with 15 Gy [250 kilovolt peak (kVp) x-rays] to inactivate persisting hematopoietic cells (Fraser, 1990). LTC are maintained by weekly removal of half of the medium and nonadherent cells and restoration of the volume with fresh LTC medium. To assay cells in LTC for repopulating cells, adherent layers are removed with a rubber policeman, suspended by passage through a 21-gauge needle and then combined with the nonadherent cells. Cells are washed once in 2% fetal calf serum/a-minimal essential medium, and aliquots from individual culture flasks are then injected i.v. into irradiated (8 Gy, 250 kVp x-rays) female recipients. In some cases, 2×10^5 female B6C3F1 marrow cells that had been previously subjected to two cycles of transplantation and regeneration are also injected to allow quantitative measurements of CRU in the retrovirally marked test cells to be obtained. For cultures that are used to assess recovery of repopulating cells in the nonadherent fraction over time, all of the medium and nonadherent cells are removed weekly and replaced either with fresh medium alone or with LTC medium containing 25 units per ml of recombinant mouse interleukin-3 (IL-3). The nonadherent cells removed after 3, 5, 6, and 7 weeks of culture are then injected into irradiated recipients. Hematopoietic Tissue Analysis. Recipients are sacrificed either 5 wk or between 5 and 7 months after transplantation of cultured cells. DNA is routinely extracted from marrow, spleen, and thymus. In some cases, DNA is also extracted from lymph nodes and from various subpopulations of marrow and spleen. Highly

enriched (>90%) mast cell populations are generated by culturing marrow or spleen cells for 3 wk in WEHI-3B-supplemented medium (as a source of IL-3), and highly enriched (>95% Mac-i-positive) macrophage populations are generated by culturing marrow or spleen cells for 48 hr in medium supplemented with 10% WEHI-3B-conditioned medium and then for 7-10 days in medium supplemented with 35% human leukocyte-conditioned medium. Highly enriched (>90% Thy-i-positive) T-cell populations are obtained by elution of nonadherent cells after loading of a half-spleen equivalent onto a 3-ml nylon wool column and incubating it for 1 hr at 37°C. Nylon wool adherent cells are then released by gentle agitation for 2-3 min in phosphate-buffered saline/10 mM EDTA and a highly enriched (>90% B220-positive) population of B lymphocytes subsequently isolated from this fraction by panning for 1 hr at 37°C in dishes precoated with unpurified rabbit anti-mouse immunoglobulin (<10⁸ cells per dish) and selective removal of the adherent cells. Southern Blot Analysis. High-molecular-weight DNA is digested with HindIII or EcoRI, which cuts once in the proviral genome and releases a fragment unique to the integration site. Ten micrograms of digested DNA (5 µg for male control lanes) is electrophoresed and analyzed by Southern blotting as described with a 32P-labeled probe specific for the neor gene sequence in the provirus from plasmid pMC1. HindIII-digested blots are stripped and reprobed with a Y chromosome-specific fragment from plasmid pY2 (Thomas, 1987).

Animal clone as an exciting but criticized topic technique in the life science is extracting the whole human community's attention. Even with ethnic and safety concern, nobody can prevent the development of the animal clone science (Ma, 2004). Science development will be benefit to all the human society (Ma).

All the stem cell research work must be controlled under the common ethic sense. The social spirit should be cared even in the scientific society.

Life is unique in the known universe, which is in a diversity form from bacteria to human. The life organisms exist in ever where in the earth. Life is a physical and chemical process. From ontology aspect, the world is timeless and the life exists forever as any other body in the nature. The nature of life is that life is a process of negative entropy, evolution, autopoiesis (auto-organizing), adaptation, emergence and living hierarchy. Up to now, there is no scientific evidence to show that life body and non-life body obey the same natural laws. But, all the researches are made by the methods of biology, biochemistry and molecular biology, etc. It is very possible that the life and non-life are essential different in the biophysics, i. e. the

quantum level. In the future, it is possible to make artificial life by either biological method or electronic technique (Ma and Cherng, 2005). The characterizations of stem cell will offer important information to answer what life is, the nature of life, if a life body can live forever or not, if a life procedure (lifeline) can be reversed or not (from old condition to younger condition), etc. It is believed people will find the way to live forever –as the immortal living body in the future, which gets eternal finally. When the eternal time comes, there will be no die. However, the so called immortal (eternal) here, it is biological immortal, not the physical immortal. Even the biological immortal achieves, the living body still can be die by disease, or killed, etc. For example, the big war still can kill people as many as it will, even to destroy all the living things in the earth. Especial, people's war ability increases very fast and the mass destroying weapons including nuclear weapons are made more in the future.

As the nature will, to live eternally is an extracting dream in all the human history. Stem cell is the original of life and all cells come from stem cells. Germline stem cell (GSC) is the cell in the earliest of the cell stage. It is possible to inject the GSC into adult human body to get the eternal life. This article is to try to describe the stem cell and to explore the possibility of the eternal life with the stem cell strategy (Ma, 2007). As the earth living beings, it is important to say that science is very important for the earth civilization, but to control the world in a peace environment is also important. Stem cell research can offer the scientific knowledge to let people live longer biologically, but cannot keep people from the war and cannot keep people from the evil.

Correspondence to:

Ma Hongbao, PhD
Brookdale University Hospital and Medical Center
Brooklyn, New York 11212, USA
Email: ma8080@gmail.com

References

1. Christopher C. Fraser, Stephen J. Szilvassy, Connie J. Eaves, And R. Keith Humphries. Proliferation of totipotent hematopoietic stem cells in vitro with retention of long-term competitive in vivo reconstituting ability. Proc. Nati. Acad. Sci. USA. Vol. 89, pp. 1968-1972, 1992.
2. Fraser, C. C., Eaves, C. J., Szilvassy, S. J. & Humphries, R. K. (1990). Blood 76, 1071-1076.
3. Thomas, K. R. & Capecchi, M. R. (1987) Cell 51, 503-512.

4. Baud L, Haymann JP, Bellocq A, Fouqueray B. Contribution of stem cells to renal repair after ischemia/reperfusion. *Bull Acad Natl Med.* 2005;189(4):635-43.
5. Bavister BD, Wolf DP, Brenner CA. Challenges of primate embryonic stem cell research. *Cloning Stem Cells* 2005;7(2):82-94.
6. Bernard Lo, Patricia Zettler, Marcelle I. Cedars, Elena Gates, Arnold R. Kriegstein, Michelle Oberman, Renee Reijo Pera, Richard M. Wagner, Mary T. Wuerth, Leslie E. Wolf, Keith R. Yamamoto. A New Era in the Ethics of Human Embryonic Stem Cell Research. *Stem Cells.* <http://www.StemCells.com>. <http://stemcells.alphamedpress.org/cgi/reprint/2005-0324v1.pdf>. 2005.
7. Bhatt RI, Brown MD, Hart CA, Gilmore P, Ramani VAC, George NJ, Clarke NW. Novel method for the isolation and characterisation of the putative prostatic stem cell. *Cytometry A.* 2003;54(2):89-99.
8. Cantley LG. Adult stem cells in the repair of the injured renal tubule. *Nat Clin Pract Nephrol.* 2005;1(1):22-32.
9. Condorelli G, Peschle C. Stem cells for cardiac repair: state of the art. *Front Biosci* 2005;10:3143-50.
10. Daar AS, Sheremeta L. The science of stem cells: ethical, legal and social issues. *Exp Clin Transplant.* 2003;1(2):139-46.
11. Duffield JS, Bonventre JV. Kidney tubular epithelium is restored without replacement with bone marrow-derived cells during repair after ischemic injury. *Kidney Int.* 2005;68(5):1956-61.
12. Duffield JS, Park KM, Hsiao LL, Kelley VR, Scadden DT, Ichimura T, Bonventre JV. Restoration of tubular epithelial cells during repair of the postischemic kidney occurs independently of bone marrow-derived stem cells. *J Clin Invest.* 2005;115(7):1743-55.
13. Herrera MB, Bussolati B, Bruno S, Fonsato V, Romanazzi GM, Camussi G. Mesenchymal stem cells contribute to the renal repair of acute tubular epithelial injury. *Int J Mol Med.* 2004;14(6):1035-41.
14. Hishikawa K, Fujita T. Stem cells and kidney disease. *Hypertens Res.* 2006;29(10):745-9.
15. http://stemcells.nih.gov/staticresources/research/protocols/BresaGen_hESC_manual_2.1.pdf
16. Humphreys BD, Duffield JD, Bonventre JV. Renal stem cells in recovery from acute kidney injury. *Minerva Urol Nefrol.* 2006;58(1):13-21.
17. Kashofer K, Bonnet D. Gene Therapy Progress and Prospects: Stem cell plasticity. *Gene Ther.* 2005 (Epub ahead of print).
18. Kimberly Kasow. OPBMT2 Protocol: Allogeneic Hematopoietic Stem Cell Transplantation for Children Affected with Malignant Osteopetrosis - A Pilot Study. http://www.stjude.org/protocols/0.2881.450_2331_17072.00.html. 2007.
19. Kwang-Soo Kim. Stem cell research continues in Korea beyond the Hwang scandal. *Stem Cells.* <http://www.StemCells.com>. <http://stemcells.alphamedpress.org/cgi/reprint/2007-0089v1.pdf>. 2007.
20. Lin F, Cordes K, Li L, Hood L, Couser WG, Shankland SJ, Igarashi P. Hematopoietic stem cells contribute to the regeneration of renal tubules after renal ischemia-reperfusion injury in mice. *J Am Soc Nephrol.* 2003;14(5):1188-99.
21. Lin F. Stem cells in kidney regeneration following acute renal injury. *Pediatr Res.* 2006;59(4 Pt 2):74R-8R.
22. Ma H. Technique of Animal Clone. *Nature and Science* 2004;2(1):29-35.
23. Ma H. Gene Transfer Technique. *Nature and Science.* 2005;3(1):25-31.
24. Ma H, Cherng S. Nature of Life. *Life Science Journal.* 2005;2(1):7-15.
25. Ma H, Cherng S. Eternal Life and Stem Cell. *Nature and Science.* 2007;5(1):81-96.
26. Mitalipova et al. *Stem Cells.* 2003;21(5):521-6.
27. Morigi M, Benigni A, Remuzzi G, Imberti B. The regenerative potential of stem cells in acute renal failure. *Cell Transplant.* 2006;15 Suppl 1:S111-7.
28. Nigel Hawkes. Scientists find the secret of eternal life for stem cells. <http://www.timesonline.co.uk/tol/news/uk/article137674.ece>
29. Oliver JA. Adult renal stem cells and renal repair. *Curr Opin Nephrol Hypertens.* 2004;13(1):17-22.
30. Paul Woodard. SCDHAP Protocol: ematopoietic Stem Cell Transplantation (HSCT) for Patients with Sickle Cell Disease and Prior Stroke or Abnormal Transcranial Doppler Ultrasound (TCD) using Reduced Conditioning and T-Cell-Depleted Hematopoietic Stem Cells from Partially Matched Family Donors - Phase I Study. http://www.stjude.org/protocols/0.2081.450_2327_18472.00.html. 2007.
31. Renee Madden. SCT521 (COG # ASCT0521) Protocol: Soluble Tumor Necrosis Factor Receptor: Enbrel (Etanercept) for the Treatment of Acute Non-Infectious Pulmonary Dysfunction (Idiopathic Pneumonia Syndrome) Following Allogeneic Stem Cell Transplantation. http://www.stjude.org/protocols/0.2881.450_2333_5873.00.html. 2007.
32. Stedman's Medical Dictionary. The American Heritage®. Houghton Mifflin Company.

- <http://dictionary.reference.com/search?q=stem%20cell>. 2002.
33. Williams D. Stem cells in medical technology. *Med Device Technol* 2005;16(3):9-11.
 34. Wing Leung. INFT2 Protocol: HLA - Nonidentical Stem Cell and Natural Killer Cell Transplantation for Children Less than 2 Years of Age with Hematologic Malignancies. http://www.stjude.org/protocols/0,2881,450_2330_11129,00.html. 2007.
 35. Yamashita S, Maeshima A, Nojima Y. Involvement of renal progenitor tubular cells in epithelial-to-mesenchymal transition in fibrotic rat kidneys. *J Am Soc Nephrol*. 2005;16(7):2044-51.
 36. Hemmat, S., D. M. Lieberman and S. P. Most (2010). "An introduction to stem cell biology." *Facial Plast Surg* **26**(5): 343-349.
 37. Sharpless, N. E. (2010). "Hot topics in stem cells and self-renewal: 2010." *Aging Cell* **9**(4): 457-461.
 38. Surani, M. A. and P. Hajkova (2010). "Epigenetic reprogramming of mouse germ cells toward totipotency." *Cold Spring Harb Symp Quant Biol* **75**: 211-218.
 39. [Pesce M](#), [Schöler HR](#). Oct-4: control of totipotency and germline determination. *Mol Reprod Dev*. 2000 Apr;55(4):452-7.

11/7/2010