

Immunomodulatory and prophylactic properties of some selected Nigerian higher fungi

Segun Gbolagade Jonathan * Olayinka Oluyemi Oluranti and Odunayo Joseph Olawuyi

Department of Botany and Microbiology, University of Ibadan, Ibadan, Nigeria
sg.jonathan@mail.ui.edu.ng

Abstract: The prophylactic potentials of five Nigerian higher fungi – *Pleurotus tuber-regium*, *Pleurotus pulmonarius*, *Termitomyces robustus*, *Fomes lignosus*, *Lentinus subnudus* and their combination in equal proportion (mixture extract) were investigated. The ethanolic fungi extracts were tested on sets of albino rats (*Rattus norvegicus*) infected with selected bacteria (*Escherichia coli*, *Staphylococcus aureus*, *Salmonella typhi*, *Pseudomonas aeruginosa*, *Streptococcus faecalis* and *Bacillus subtilis*). Extracts were orally administered at 1.0ml concentration against 0.1ml/cfu microbial suspension intraperitoneally. Survival rates of the experimental rats were monitored. Mixture of these extracts exhibited significant level of prophylactic effect against all the bacteria in the host organisms.

[Segun Gbolagade Jonathan, Olayinka Oluyemi Oluranti and Odunayo Joseph Olawuyi. **Immunomodulatory and prophylactic properties of some selected Nigerian higher fungi.** *Stem Cell* 2017;8(1):94-99]. ISSN: 1945-4570 (print); ISSN: 1945-4732 (online). <http://www.sciencepub.net/stem>. 16. doi:[10.7537/marsscj080117.16](https://doi.org/10.7537/marsscj080117.16).

Keywords: Prophylactic, Extracts, Survival rates, Infection, Bacteria.

1. Introduction

Pleurotus tuber-regium is a tropical mushroom and an edible higher fungus which produces a sclerotium or underground tuber. A sclerotium is a dark brown, yam-like material specially produced by some specific higher fungi (Oso, 1977; Jonathan 2002; Jonathan *et al.*, 2008). It could be used in combinations with other plants parts to treat headache, stomach ailments, colds, fever, asthma, smallpox and high blood pressure (Oso, 1977). The fruitbodies and sclerotium of this basidiomycete have been found to possess antimicrobial potential against some disease causing microorganisms (Jonathan and Fasidi, 2003). Also, *Pleurotus pulmonarius* (oyster mushroom) has been produced using several agricultural residues (Bononi *et al.*, 1995). Banana leaf, mixed with sugarcane bagasse or corn cob are examples of the substrates used for the production of *P. pulmonarius* (Chang *et al.*, 1981) while cassava residues with sugarcane bagasse are used for the production of *P. oestreatus* (Felinto, 1999).

Fomes is a genus of polypore higher fungus which belongs to the order Aphyllophorales, and family polyporaceae (Alexopoulos *et al.*, 1996). The basidiocarps of *Fomes* may generally be recognised by conspicuous fruitbodies which are perennial with attractive yellow or brownish colouration. The sporocarp are fleshy when young and becomes woody on aging (Jonathan, 2002). *Fomes lignosus* had also been reported to be utilized in Nigeria folk medicine due to its inhibitory potentials against some microbes implicated in the pathogenesis of skin infections, food poisoning, gastro-intestinal tract and urino-genital tract infections (Olawuyi *et al.*, 2010).

Termitomyces is a genus of basidiomycete fungi belonging to the family Lyophyllaceae (Kirk *et al.*, 2008). They usually grow in close association with termite nests. Despite their spore, transfer is mainly by shedding from higher fungi which protrude from the termite mounds (Mueller *et al.*, 2005). *Termitomyces titanicus* of West Africa is one of the largest higher fungi in the world.

Lentinus is a genus whose plants are tough and pliant, becoming hard when old unless very watery. *Lentinus subnudus* belongs to the family polyporaceae which is the largest and the most diverse group of poroid aphyllophorales (Pegler, 1983). Though, grows naturally on dead woods, logs and trunks, they can also be cultivated in logs of *Sporandias mombin* and unfermented composts (Fasidi and Kadiri, 1993; Gbolagade *et al.*, 2006).

Mushrooms are generally known for their nutritional and medicinal properties. This study was aimed at examining the immunomodulatory and prophylactic potentials of these higher fungi found in Nigeria (Aina *et al.*, 2012).

2. Materials and methods

2.1 Higher fungi and extracts preparation

Fruit bodies of *Pleurotus tuber-regium*, *P. pulmonarius*, *Fomes lignosus*, *Termitomyces robustus* and *Lentinus subnudus* were cut into bits, dried at 40°C and milled into powder using grinding machine. Ethanol was used as solvent for the extraction of the higher fungi using the methods described by Jonathan and Fasidi (2003).

2.2 Experimental rats

Albino rats (*Rattus norvegicus*) were used for the study and were about 12 weeks old with average weight of 132 grammes.

2.3 Prophylactic experiment

One millilitre (1mL) of the extract was administered orally into each experimental animal for 7 consecutive days. The test bacteria (*E. coli*, *S. aureus*, *S. typhi*, *P. aeruginosa*, *S. feacalis*, and *B. subtilis*) were homogenised in peptone water and 0.1ml of the microbial suspension were intraperitoneally administered into each animal post-higher fungi extract administration. This was done thrice at intervals of 2 days. Infection was then monitored for about 10 days. Survival rate of the experimental rats was based on the number that survived the infection. Dead animals were dissected immediately to remove the livers and kidneys, while the survived ones were later sacrificed. The excised

organs were weighted and kept in formalin (10%) for histological studies.

2.4 Histological studies

The livers and kidneys kept in 10% formalin were histologically examined for pathological changes (Wiess *et al.*, 2000, Mark Source, 2002).

2.5 Analysis of Data

The results of this study were subjected to the analysis of variance (ANOVA) and Duncan's Multiple Range Tests (DMRT) ($p < 0.05$).

3. Results and Discussion

The five higher fungi in this study were found to exert prophylactic effects against the infected organisms in the test animals, though at different levels. The mixture extract and *Pleurotus pulmonarius* were most effective against *E. coli*. Survival rates of experimental rats were 100% after the bacterial infection (Table 1).

Table 1: Survival rates of *Rattus norvegicus* used for prophylactic administration of the higher fungi against *E. coli*.

Day	FOM	PT	PS	Term	Lent	Mixt	Control A	Control B
1	91.67 ^a	75.00 ^b	100.00 ^a	100.00 ^a	100.00 ^a	100.00 ^a	75.00 ^b	100.00 ^a
5	75.00 ^b	25.00 ^c	91.67 ^a	75.00 ^b	75.00 ^b	100.00 ^a	0.00	100.00 ^a
10	75.00 ^b	16.67 ^c	91.67 ^a	75.00 ^b	75.00 ^b	100.00 ^a	0.00	100.00 ^a

Values with the same letter(s) in each row are not significantly different by Duncan's multiple range test ($P < 0.05$). Each is a mean of three replicates.

Key:

FOM = *Fomes lignosus*, PT = *Pleurotus tuber-regium*, PS = *Pleurotus pulmonarius*, Term = *Termitomyces robustus*, Lent = *Lentinus subnudus*, Mix = Mixture extract, Control A = Infected rats without extract, Control B = Uninfected rats.

Lentinus subnudus and *Pleurotus tuber-regium* demonstrated the best prophylactic activity against *S. aureus* with survival rates of 100% and 91% respectively (Table 2). *F. lignosus*, *T. robustus* and the mixture extracts were very effective against *S. typhi*. Virtually all the animals survived the infection. *L. subnudus* also had an appreciable effect against the organism (Table 3). Both *Pleurotus* species (*Pleurotus tuber-regium* and *P. pulmonarius*) had the same

effect. The prophylactic effects against *P. aeruginosa* shown in Table 4 indicates *P. tuber-regium* as the best extract with a 100% animal survival throughout the experimental period. Table 5 shows that all the higher fungi extracts were uniformly active against *S. Feacalis*, while the mixture extract and *F. lignosus* were the most active extracts against *B. subtilis* (Table 6). Survival rates of the animals were 100% and 91% respectively.

Table 2: Survival rates of *Rattus norvegicus* used for prophylactic administration of the higher fungi against *S. aureus*

Day	FOM	PT	PS	Term	Lent	Mix	Control A	Control B
1	83.33 ^{ab}	100.00 ^a	75.00 ^b	75.00 ^b	100.00 ^a	83.33 ^{ab}	50.00 ^c	100.00 ^a
5	66.67 ^b	91.67 ^a	50.00 ^c	75.00 ^b	100.00 ^a	75.00 ^b	0.00	100.00 ^a
10	66.67 ^b	91.67 ^a	25.00 ^c	75.00 ^b	100.00 ^a	75.00 ^b	0.00	100.00 ^a

Table 3: Survival rates of *Rattus norvegicus* used for prophylactic administration of the higher fungi against *S. typhi*

Day	FOM	PT	PS	Term	Lent	Mix	Control A	Control B
1	100.00 ^a	75.00 ^b	75.00 ^b	100.00 ^a	100.00 ^a	100.00 ^a	75.00 ^b	100.00 ^a
5	91.67 ^a	66.67 ^b	66.67 ^b	100.00 ^a	75.00 ^b	100.00 ^a	0.00	100.00 ^a
10	91.67 ^a	66.67 ^b	66.67 ^b	100.00 ^a	75.00 ^b	100.00 ^a	0.00	100.00 ^a

Table 4: Survival rates of *Rattus norvegicus* used for prophylactic administration of the higher fungi against *P. aeruginosa*

Day	FOM	PT	PS	Term	Lent	Mix	Control A	Control B
1	100.00 ^a	100.00 ^a	83.33 ^b	83.33 ^b	75.00 ^b	83.33 ^b	75.00 ^b	100.00 ^a
5	75.00 ^b	100.00 ^a	66.67 ^b	50.00 ^c	50.00 ^c	75.00 ^b	0.00	100.00 ^a
10	66.67 ^b	100.00 ^a	66.67 ^b	50.00 ^c	50.00 ^c	75.00 ^b	0.00	100.00 ^a

Table 5: Survival rates of *Rattus norvegicus* used for prophylactic administration of the higher fungi against *S. feacalis*

Day	FOM	PT	PS	Term	Lent	Mix	Control A	Control B
1	75.00 ^b	75.00 ^b	100.00 ^a	75.00 ^b	100.00 ^a	83.33 ^{ab}	50.00 ^c	100.00 ^a
5	66.67 ^b	75.00 ^b	75.00 ^b	75.00 ^b	75.00 ^b	75.00 ^b	0.00	100.00 ^a
10	66.67 ^b	75.00 ^b	75.00 ^b	75.00 ^b	75.00 ^b	75.00 ^b	0.00	100.00 ^a

The histological findings on the tissues of rats infected post-fungi extract administration are shown in Tables 7 and 8. These further confirmed the prophylactic effect of the higher fungi extracts. No visible lesions were seen on both liver and kidney tissues of rats administered with the extracts 7 days before infection and also the control uninfected rats. The control infected rats (without extracts) showed some abnormalities in the kidney (marked portal

congestion, diffuse tubular necrosis and numerous tubules with protein casts in the tubular lumen) and the liver (marked bile duct proliferation, mild cellular infiltration). The pathological changes were prevented in the experimental rats by the protective and preventive potentials of the higher fungi extracts (Plates 1 and 2).





**A** No visible lesions**B** X400 There is marked portal congestion (arrows) and mild portal fibrosis (stars).**Plate 1: Photomicrograph of the liver of rats infected with *Escherichia coli* with no visible lesions (A) and the kidney with marked portal congestion (B)**

Table 6: Survival rates of *Rattus norvegicus* used for prophylactic administration of the higher fungi against *Bacillus subtilis*

Day	FOM	PT	PS	Term	Lent	Mix	Control A	Control B
1	100.00 ^a	75.00 ^b	100.00 ^a	100.00 ^a	100.00 ^a	100.00 ^a	75.00 ^b	100.00 ^a
5	91.67 ^a	75.00 ^b	0.00	75.00 ^b	75.00 ^b	100.00 ^a	0.00	100.00 ^a
10	91.67 ^a	75.00 ^b	0.00	75.00 ^b	75.00 ^b	100.00 ^a	0.00	100.00 ^a

**A** There is very mild cellular infiltration at the portal area.**B** No visible lesions seen.**Plate 2: Photomicrograph of rats' livers infected with *Bacillus subtilis* (A) Mild cellular infiltration (B) The kidney of infected rates with no visible lesions.****Table 7: Histological findings on tissues of rats infected with *E. coli*, those treated with *Fomes lignosus* and control**

Group	Organ	Findings
Rats administered with <i>F. lignosus</i> extract 7 days before infection	Liver	No visible lesions seen
	Kidney	No visible lesions seen
Control (uninfected)	Liver	No visible lesions seen
	Kidney	No visible lesions seen.
Control (infected)	Liver	No visible lesions
	Kidney	Marked portal congestion

Table 8: Histological findings on tissues of rats infected with *Bacillus subtilis*, those treated with Mixture extract and control

Group	Organ	Findings
Rats administered with Mixture extract 7 days before infection	Liver	No visible lesions seen
	Kidney	No lesions developed
Control (uninfected)	Liver	No visible lesions seen
	Kidney	No visible lesions seen.
Control (infected)	Liver	Mild cellular infiltration at the portal area.
	Kidney	No visible lesions.

On the overall, the mixture extract was found to have the best prophylactic effect against the bacterial infections in the experimental rats. The survival rates of the control (uninfected) rats remain the same (100%) throughout the period of experiment. For the control infected rates, no animal survived after the 4th day of infection.

Wang and Wang (1997) obtained similar results when glucans extracted from different polysaccharides were used against *Aeromonas hydrophila* and *Edwardsiella tarda* infections in tilapia. The survival rates of the fish increased significantly. Typically for fungi, the most active components of their immunomodulators are the glucan structures. The immunomodulatory properties of these mushrooms are due to the presence of glucans in their cell wall (Benedict 1972; Kupra *et al.*, 1979; Suzuki *et al.*, 19990; Brandt *et al.*, 2000). According to Medzhitove *et al.* (1997), the glucan structures are generally indicated as pathogen associated molecular patterns. Betaglucan is one of the most effective immunomodulators and are recommended by Claypton (2004), because of its ability to activate macrophage. An example of betaglucan is lentinan obtained from *Lentinus edodes* which stimulates natural killer cell, T-cell and macrophage dependent responses (Nanba *et al.*, 1987). A mushroom compound (active hexose) which is a product of the enzymatic modification of several types of medicinal higher fungi has the ability to increase natural killer cell and macrophage activity. Moreover, several clinical tested health supplements from *Fomes lignosus* have been considered as best natural products that can be consumed to regulate healthy diet and also boost the immune system. *Grifola frondosa* (maitake) enhance the activities of natural killer cells, T-cells and macrophages. In Asia, *Trametes versicolor* is a popular higher fungi that has been used traditionally because it stimulates macrophages. *Agaricus blazei* also has the ability of increasing T-cell activity thereby stimulating the immune system. The beta glucans associated with this mushroom are the active polysaccharides (having 1, 6-betaglucan as the most potent immunomodulator). In one animal study, *Cordyceps sinensis* was discovered to increase antibody thereby enhancing immunity. Also, in an in-vitro study, a cordyceps polysaccharide was found to increase cytokines (Wang *et al.*, 1994; Kurashige *et al.*, 1977; Wassei and Weis, 1999).

Stamets (2001), also reported that compounds with antibacterial properties are prepared from mushroom such as *Fomitopsis*, *Piptoporus*, *Ganoderma*, *Inonotus* and *Trametes* species. They are also useful in preventing and treating viruses such

as poxyiridae and orthopox viruses, flu viruses including bird flu, SARS and hepatitis as well as infections from *Mycobacterium tuberculosis*, *S. aureus* and *E. coli*.

Generally, mushrooms have been suggested to have immunological properties, giving protection against a number of infectious diseases (Hobbs, 1986; Chilara, 1992; Mizuno *et al.*; Jonathan and Awotona, 2010).

The findings of Medzhitov *et al.* (1997) confirmed that the mixture of mushroom polysaccharides gives the maximum response in the immune system by increasing the number and enhancing the activity of killer, T and NK (natural killer) lymphocytes.

4.0 Conclusion

It is clear from the study that preventive abilities were exhibited against the test pathogenic bacteria. This could be attributed to the prophylactic effects of the higher fungi extracts. The strength for the prophylactic activity was obtained within the seven days pre-infection fungi-extract administration. This was evidenced in the survival rates (100%) of the experimental rats used for the study. The ability of the extracts to exert prophylactic effect within a week at low concentration has to do with the immune-enhancing properties of the higher fungi. Therefore, they are recommended as good prophylaxis for prevention and protection against infections, especially the blend or mixture of the higher fungi.

Correspondence to:

Dr. Segun. Gbolagade Jonathan
Department of Botany and Microbiology,
University of Ibadan
E-mail: sg.jonathan@mail.ui.edu.ng
Tel: +2348164746758

References

1. Aina DA, Oloke JK, Jonathan SG and Olawuyi OJ(2012). Comparative assessment of mycelial biomass and exo-polysaccharide production of wild type and mutant strains of *Schizophyllum commune* grown in submerged liquid medium *Nature and Science* 10(10):82-89.
2. Alexopoulos, C.J., Mims, and Blackwell M. 1996. *Introductory Mycology*. 3rd Edition. John Wiley and Sons, New York..
3. Benedict, R.G and Braddy L.R. 1972. Antimicrobiology activities of higher fungi metabolites. *JA Pharm. Sci.* 61pp 1820-1822.
4. Bononi, V.L., Capatari, M; Maziero, R, Trifem, S.F.B. 1995. *Cultivo de cogumelos comstivels Sao Paulo* ; Icon, 206p.

5. Brandt, C.R., Piriano F. 2000. Higher fungi antivirals. *Recent Res. Dev. for Antimicrob. Agents and Chemoth.* 4: 11 – 26.
6. Chang, S.T. Lau O.W., Cho, K.Y. 1981. The cultivation and nutritional value of *P. sajorcaju*. *European J. of Appl. Microb. and Biotechn.*, 12: 58 – 62.
7. Chilara G. 1992. Immunopharmacology of lentinan, a polysaccharide isolated from *Lentinus edodes*: its application as a host defense potentiator. *Intl. J. of Oriental Med.* 17: 55 – 77.
8. Clayton, P. (2004). Bird ‘Flu’ and Glucan. *The Nutrit. Pract.* (6) 3. 01189798686.
9. Fasidi, I.O., and Kadiri, M. 1993. Effect of sporophore maturity on chemical composition of *Volvariella esculenta* (Mass) Singer, a Nigerian edible higher fungi. *Die Nahrung.* 37(3) 269 – 273.
10. Felinto, A.S. 1999. Cultivo de cogumelos comestiveis do genero *Pleurotus* spp em residues agroindustriais. *Picaciraba: USP/ESALQ*, 123p. (Dissertacao-Mestrado).
11. Gbolagade J.S, Fasidi,I.O, Ajayi E.J and Sobowale A.A (2006). Effect of physico-chemical factors and semi-synthetic media on vegetative growth of *Lentinus subnudus* (Berk.), an edible mushroom from Nigeria. *Food Chemistry.* 99:742-747.
12. Hobbs C. 1986. *Medicinal higher fungi*. Loveland (C.O); Interweave Press. Jenkins, D.T. 1986. *Amanita of N. America Mad River*. Eureka, CA.
13. Jonathan S.G. 2002. Vegetative growth requirements and antimicrobial activities of some higher fungi in Nigeria. Ph.D thesis, University of Ibadan.
14. Jonathan, S.G. and Fasidi I.O. 2003. Requirements for vegetative growth of *Tricholoma lobayensis* (Heim) A. Nigerian Edible Fungus, *Advances in Food Sc.* 25(3) 91 – 95.
15. Jonathan, S.G. Kigigha, L.T. and Ohimain E. 2008. Evaluation of the inhibitory potentials of eight edible higher Nigerian fungi against pathogenic microorganisms. *Afr. Journal of Biomed Sc.* 11: 195 – 200.
16. Jonathan, S.G and Awotona F.E 2010. Studies on antimicrobial potentials of three Ganoderma species. *African J. of Biomed. Res.* 13(2) 119-125.
17. Kirk, P.M., Cannon, P.F., Minter, D.W., Stalpers, J.A. 2008. *Dictionary of Fungi* (10th ed). Wallingford UK: CABI p. 682. ISBN 0 – 8. 5199-826-7.
18. Kupra J., Anke T., Oberwinkler G., Schramn G., Steglich, W. 1979. Antibiotics from Basidiomycetes vii. *Crinipellis stipitaria* (Fr) Pat. *J. of Antibiotics.* 32: 130-5.
19. Medzhitov R., Janeway Ch. A. 1997. Innate immunity: Impact on the adaptive immune response. *Curr. Opinion in Immunology.* 9, 4-9.
20. Mizuno, T., Saito H., Nishitoba, T., Kawagishi H. 1995. Antitumor active substances from higher fungi. *Food Reviews Int.* 11: 23-61.
21. Mizuno, T., Zhuang C. 2000. Houbitake, *Pleurotus pulmonarius* antitumor activity and Utilization. *Food Reviews Int.* 11:185-187.
22. Mueller, U.G., Gerardo, N.M., Aaran, D.K., Six D.L., Schultz, T.R. 2005. “The evolution of Agriculture in insects.” *Ann. Rev. of Ecol., Evol. and Systematics* 36: 563.
23. Nanba N., Mori K., Toyomasu T, Kuroda H, 1987. Antitumor action of shiitake (*Lentinus edodes*) fruit bodies administered to mice. *Chem. Pharm. Bull.* 35: 2453-2458.
24. Olawuyi O.J. Jonathan S.G., Aina D.A, and Baysah G.I. 2010. Studies on antibacterial activity of *Fomes lignosus* – a Nigerian mushroom. *Nig J. of Microb.* 24(1): 2153 – 2159.
25. Oso, B.A. 1977. *Pleurotus tuber-regium* from Nigeria. *Mycologia.* 67(2): 271-277.
26. Pegler, D.N. 1983. The Genus *Lentinus*. Kew. Bull. Add. Series 10.
27. Stamets, P. (2001). Novel Anti-virals from Higher fungi *Herbal Gram* 51: 24, 27.
28. Sturion, G.L., Oettener M. 1996. Composicao quimica de cogumelos comestiveis (*Pleurotus* spp) orginado de cultivos em differences substrates. *Ciencia e Tecnologia de Alimentos.* 15 : 189 – 193.