

**Morphological and Leaf Epidermal features of *Capsicum annum* and *Capsicum frutescens* (Solanaceae)**

C.U Nwachukwu

Department of Biology, Alvan Ikoku College of Education. Owerri, Imo State  
[nwachukwucu2005@yahoo.co.uk](mailto:nwachukwucu2005@yahoo.co.uk)

FN MBAGWU

Department of Plant Science and Biotechnology, Imo State University Owerri

Onyeji Augustina Nwamaka

Department of Plant Science and Biotechnology, Imo State University Owerri

**ABSTRACT:** Studies on the morphological (vegetative and floral) and leaf epidermal features of *Capsicum annum* and *Capsicum frutescens* found in different parts of Imo State were investigated. This was with the main aim of evaluating their reliability as aid, in determining intraspecific relationships among these taxa. Morphological features among the two taxa studied include variations and similarities in habit. The study showed annual herb in *Capsicum annum* and perennial herb in *Capsicum frutescens*. Similarly the height of *Capsicum annum* is 60 cm, while *Capsicum frutescens* is 72 cm.

Furthermore, the two taxa shared common attributes as revealed in the stem, leaf and the floral morphology. The stem type, colour and bark showed that the two taxa are erect-branched from base, green and smooth respectively. The floral morphology further strengthens the Intraspecific relationships among these two taxa. The floral result revealed that the flower type, symmetry, arrangement, pedicel, calyx and corolla colour and shapes are all the same in the two taxa studied. The leaf epidermal characters in the two taxa studied did not show much variation except in the number of stomata: 21 and 44 and 18 and 60 on the upper and lower surfaces of *Capsicum annum* and *Capsicum frutescens* respectively. The other attributes of stomata type, number of subsidiary cells, shape of epidermal cells and presence of trichome are the same for the two taxa hence this study confirms the values of morphological and leaf epidermal features in the systematics and biological consideration of *Capsicum annum* and *Capsicum frutescens*. [The Journal of American Science. 2009;5(1):50-58]. (ISSN: 1545-1003).

**Keywords:** Morphology; Leaf epidermis; *Capsicum*; Solanaceae; Systematics

**INTRODUCTION:**

The genus *Capsicum* belongs to the family Solanaceae (Night shade). Members of the Solanaceae family are mostly herbs or under herbs while some others are climbers. The family contains about 90 genera and nearly 3000 species (Vidyarth and Tripatha 2002, Stern 2000). The genus *Capsicum* is further classified into the division Magnoliophyta, class magnoliopsida, order solanates and family Solanaceae (Heiser and Smith 1953). *Capsicum* is a crop that is widely cultivated because of its spicy nature and nutritional value. The crop accounts for a large portion of vitamins A and C in many Nigerian diets. *Capsicum annum* and *Capsicum frutescens* are the most common species in Nigeria. Heiser and Smith(1953) distinguished two *Capsicum* species cultivated as vegetables while varieties are all forms of either *Capsicum annum* or *Capsicum frutescens*. *Capsicum annum* is not known in a wild state and species commonly cultivated are *Capsicum annum* known as sweet pepper, bell pepper, cherry pepper and

green pepper (Messraen 1992, Gill, Hs. Thakur P.C and Thakur T.C 1973.). *Capsicum frutescens* on the other hand occurs in the wild though became domesticated in many parts of the tropics. Species commonly cultivated as *Capsicum frutescens* are known as bird eye pepper, red pepper and Tobacco pepper (Heiser and Smith 1953).

The economic importance has been discussed. In West Africa and in Nigeria in particular, *Capsicum annum* and *Capsicum frutescens* are third among the cultivated vegetables being utilized in the dry state as spice, *Capsicum* content, an alkaloid that is a digestive stimulant is used as ointment for arthritis and neuropathic pains (Stern 2000). *Capsicum* species are rich in Vitamin A potency which is responsible for red colour in mature fruit, as well as ab out 50 – 280 Mg/100g of Vitamin C. *Capsicum annum* and *Capsicum frutescens* are further used as pungent spices for domestic culinary purposes and by food manufacturing industries for seasoning of processed foods in the preparation of curry powder, hot sauce and in pickling (Tindall 1986). *Capsicum* species are mostly herbs with branched top roots. The stem is herbaceous, erect and hairy; leaves are alternate, opposite in flora region, simple and estipulate. The placentation is axile, ovules are numerous, style single terminating in a bilabed stigma. The fruit is a berry and the seeds are minute endospermic with a straight or curved embryo (Esua 1977).

The use of morphological and leaf epidermal features has been found to be of immense interest in taxonomy. An excellent review of the application of morphological features in systematic studies is shown in the works of Okwulehi and Okoli, 1999, Chakrabarty and Gupta, 1981 and Olowokudejo, 1990: Edeoga and Eboka 2000, Edeoga and ikem 2001 and Stern 2000. Furthermore, the use of leaf epidermal features (epidermal cell, stomata and trichome) in systematics has become popular and distinctive and have been used as a great taxonomic tool at the levels of family, genus and species. The works of Paliwal (1967), Shah and Gopal (1972). Gill and Karatela (1982), Edeoga (1991) Edeoga and Osawe (1996) are typical examples. This study assesses the relevance of morphological and leaf epidermal features of *Capsicum annum* and *Capsicum frutescens* as well as to evaluate the reliability of these characters in the systematic consideration of the *Capsicum* species studied.

### **Materials And Methods**

The laboratory work was carried out at the Plant Science and Biotechnology Department of University of Nigeria Nsukka between September and November, 2006. These studies were made on mature living fresh materials of *Capsicum annum* and *Capsicum frutescens* collected from the garden behind the Education Trust Fund Block 11, the staff nursery/primary school garden and the garden beside Saint Joseph Chaplaincy all within Imo State University, Owerri, Imo State.

For Morphological studies Twenty mature leaves of each taxa from the middle portion of the plants were collected, length and width of the leaves were measured using a zoom meter rule. The length of the leaf was obtained by spreading the middle leaflet on a flat surface on the laboratory bench, while for the width the same media leaflet was chosen and measured to ensure uniformity Olowokudejo (1990). The seed number per pod was obtained by counting the number of seeds in the biggest pod of each taxa to ensure consistency. Photographs of the fresh materials were taking using ordinary camera and characters of the two taxa were divided into vegetative and floral morphology and tabulated.

For the leaf epidermal features, twenty samples of each species of *Capsicum* studied were examined using light microscope. An area of about 1cm square was removed from a central standard position, always midway between the base and Apex of the mature and fresh leaves of the two

taxa studied. Epidermal preparations were made by boiling the collected materials of each of the two taxa in different test tubes containing 70% ethanol for 10min. These were allowed to cool and latter bleached in 8% sodium hydrochlorite (Naocl) for 5min.


Epidermal peals were stained with 1% ethanol safranin and temporarily mounted in aqueousglycerol solution (Cutler, 1978). Photo micrographs of the epidermal features were taken from the slides using a Leitz wetzler ortholux microscope fitted with vivitar v-335 camera.

## Results

The morphological (vegetative and floral) and leaf epidermal features of *Capsicum annum* and *Capsicum frutescens* investigated are summarized in Table 1, 2 and 3 for the morphology and Table 4 for the leaf epidermis and illustrated in (Figs. 1a - b and 2a - b). The vegetative results of the two taxa studied showed that the habit and height of *Capsicum annum* is annual herb and 60cm while the habit and height of *Capsicum frutescens* is perennial herb and 72cm respectively. The habit still confirm the herbecous nature of these taxa. The attributes of stem type, colour and bark are similar to both taxa. Similarly, the leaf shape, leaf apex, leaf base and leaf type equally reveals similarity in both taxa except for leaf arrangement which is alternate in *Capsicum annum* and opposite in *Capsicum frutescens* (Table 2).

Furthermore, the floral result of the two taxa studied showed that the attributes of flower type,floral symmetry, pedicel, calyx colour, shape, corolla colour, fruit type are the same in the taxa. The difference in the floral result is observed in the flower arrangement: opposite in *Capsicum annum* and alternate in *Capsicum frutescens*,: Corolla fusion: fused in *Capsicum annum* and free in *Capsicum frutescens* and fruit shape indicates ovoid in *Capsicum annum* and linear in *Capsicum frutescens* (Table 3).


The result of the leaf epidermal features of *Capsicum annum* and *Capsicum frutescens* investigated are summarized in Table 4. The shape of the epidermal cells of the upper and lower surfaces is irregular and sinuous in both taxa studied. The two *Capsicum* taxa were amphistomatic having stomata at both the adaxial (upper) and abaxial (lower) surfaces of the leaf. The distribution of stomata on both the upper and the lower surfaces of the leaves of the taxa studied revealed anomocytic type of stomatal arrangement. The stomatal index ranged from 18% - 21% in the upper surfaces of the two taxa studied respectively and from 44% in *Capsicum annum*, 60% in *Capsicum frutescens* in the lower surfaces respectively. This implies that the stomatal density is therefore highest on the lower epidermis and lowest on the upper epidermis. There were absence of trichomes in the *Capsicum* taxa investigated.




*Fig 1a: Habit of Capsicum anum*



*Fig 1b: Habit of Capsicum frutescens*



2a




2b

Leaf epidermal features of lower leaf epidermis of *Capsicum annum* and *Capsicum frutescens*


S – Stoma Ec. – epidermal cell

A - *Capsicum annum* with much stomata and broad epidermal cell shape

B - *Capsicum Frutescens* with much stomata and small epidermal cell shape



**3a**



**3b**

Leaf epidermal features of upper leaf epidermis of *Capsicum annuum* and *Capsicum frutescens*

S – Stoma E.c – epidermal cell

A – *Capsicum annuum* with scattered and fewer stomata and broad epidermal cell shape

B – *Capsicum frutescens* with scattered and fewer stomata and small epidermal cell shape

Table 1 Vegetative characters of the two *Capsicum* species studied

Character	<i>Capsicum annum</i>	<i>Capsicum frutescens</i>
Habit	Annual herb	Perennial herb
Height	60cm	72cm
Stem type	Erect branched from base	Erect branched from base
Colour	Green	Green
Bark	Smooth	Smooth
Leaf texture	Smooth	Smooth
Leaf arrangement	Alternate	Opposite
Leaf shape	Ovate	Ovate
Leaf apex	Mucronate	Mucronate
Leaf base	Round	Round
Leaf type	Simple	Simple
Length(cm)	9.9± 1.13	5.35±0.15
Width (cm)	4.9±0.8	4.2±1.1

Table 2. Floral morphological characters of the two *Capsicum* species studied

Character	<i>Capsicum annum</i>	<i>Capsicum frutescens</i>
Flower type	Auxillary cyme	Auxillary cyme
Floral symmetry	Actinomorphic	Actinomorphic
Arrangement	Opposite	Alternate
Pedicel	Erect	Erect
Calyx colour	Greenish white	Greenish white
Shape	Elliptic	Elliptic
Corolla colour	White	White
Corolla free/fused	Fused	Free
Fruit type	Pod	Pod
Fruit shape	Ovoid	Linear
Seed number per pod	4 seeded	6-8 seeded
Pod length(cm)	1.5±0.11	5.35±0.15
Pod width(cm)	2.65±0.25	4.15±0.35

Table 3 list of *Capsicum* species from which fresh materials were collected

Collection number	Taxa	Locality	Collector(s)	Place of deposition
001	<i>C .annum</i>	Staff nursery/primary school garden Garden behind education trust fund block II Garden beside St Joseph catholic chaplaincy	Onyeji Augustina Nwamaka Ojimba Chioma  Akpaka Doris	IMSUH
002	<i>C.frutescens</i>	Staff nursery/primary school garden Garden behind education trust fund block II Garden beside St Joseph catholic chaplaincy	Iwuchukwu Ikechukwu  Iwu Jane  Ochiji Chidimma	IMSUH

IMSUH: Imo State University Herbarium

Table 4 Epidermal characteristics of the *Capsicum* species studied

Characters	<i>C. annum</i>		<i>C. frutescens</i>	
	Upper surface	Lower surface	Upper surface	Lower surface
Shape of epidermal cell	Irregular and sinuous	Irregular and sinuous	Irregular and sinuous	Irregular and sinuous
Stomatal type	Anomocytic	Anomocytic	Anomocytic	Anomocytic
% stomatal index	21	60	18	44
Number of subsidiary cells	None	None	None	None
Trichomes	Absent	Absent	Absent	Absent

### Discussion

The result of the morphological and leaf epidermal features of *Capsicum annum* and *Capsicum frutescens* studied showed some characteristic that could be used for taxonomic decision. Morphologically, the vegetative features of habit and height of *Capsicum annum* distinguish it from *Capsicum frutescens*. The observation is in line with earlier works of Okwulehi and Okoli, 1999, and Edeoga and Eboka 2000, who used comparative morphology of different species in establishing relation among various taxa. The result of leaf arrangement showed alternate shape in *Capsicum annum* and opposite in *Capsicum frutescens*. This observation is supported by the works of Edeoga and Eboka 2000, in *Dissotis* (Okeke and Nwachukwu 2001) in Euphorbiaceae but not in the genus *Capsicum*.

The result of the leaf epidermal studied indicates that the shape of epidermal cells of the *Capsicum* species studied are irregular and sinuous. Similarly, the two taxa are amphistomatic with more stomata on the lower surface (abaxial) than the upper surface (adaxial). The percentage stomatal index of the two taxa was highest on the lower epidermis 44% - 60% compared to the upper epidermis 18% - 21%. The stomatal index result is not strange since Olowokudejo (1990) found stomatal index valuable and very reliable in distinguishing between the leaves of medicinal species of *Ocimum* from non medicinal ones. The absence of trichome in the two taxa studied may not be of taxonomic importance and hence not strange since Esua (1977) insisted that much reliability is not always accorded to trichomes alone for taxonomic conclusion due to their similarities in different species. The anomocytic type of stomata found in the two *Capsicum* species indicates that the species are phylogenetically related (Nwachukwu and Mbagwu 2006). The morphological (vegetative and floral) and leaf epidermal features of the taxa studied confirm the intraspecific relationships between *Capsicum annum* and *Capsicum frutescens* in their stem, leaf, flower, fruit, seed, stomatal type and index attributes hence the differences in leaf and flower arrangement, fusion of corolla and fruit shape are not enough to separate the two taxa studied. These distinguishing morphological and leaf epidermal features observed in these investigations are of systematic value because they are reasonably constant in the taxa studied. Olowokudejo (1990) made similar observation in the genus *Anonna*. The purpose of this study is to show that application of morphological features has proven to be of immense assistance in interpreting problems related to plant classification. Thus the necessity of

including the results from the morphological and leaf epidermis with data derived from other botanical disciplines remain vital when formulating conclusions on the systematic of the *Capsicum* species.

#### References

1. Chakrabarty T. and Gupta 1981. Morphological studies on three herbaceous species of railway tract. *Proc. Int Aced. Sci. Pl.* 90:305-315.
2. Cutler D.F 1978. Applied Plant Anatomy. Longmans, London and New York. Pp. 2 - 9.
3. Edeoga H.O 1991. Taxonomic studies on certain species of *Dioscorea* L. (Dioscoreaceae) in Eastern Nigeria. Phd Thesis, University of Porthacourt, Nigeria.
4. Edeoga H.O and Eboka A.U 2000. Morphology of the leaf epidermis and systematics in some *Dissotis* species Benth (Melastomataceae). *Global J. pure and applied sci.* 6:371-374.
5. Edeoga H.O and Ikem C.L 2001. Comparative Morphology of the leaf epidermis in three species of *Boehavia* L Nyctaginaceae. *J.Pl. Anat Morph*, 1:14-21.
6. Edeoga H.O and Osawe P.I 1996 Cuticular studies on some Nigerian species of *senna* Tourn. Ex Mill (Syn *Cassia* ex.L) Leguminosae-Caesalpinoideae. *Acta Phyto tax Geobot* 47:41-46.
7. Esua K. 1977. Anatomy of seed plant (2nd Edition) Wiley, London, Pp: 83-97.
8. Gill, L.S and Karatela Y.Y 1982. Epidermal structure and stomatal ontogeny in some Nigerian Cucurbitaceae. *Willdenowia* 12:303-310.
9. Heiser, C.B and Swith P.C 1953 .The cultivated *Capsicum*. *Eco. Bot* 1:214-227.
10. Messiaen C.M 1992. The Tropical Vegetable Garden. Macmillian (London) Pp 235.
11. Nwachukwu, C.U and Mbagwu, F.N (2006). Anatomical Studies on the petiole of some species of *Indigofera*. *Medwell, Agricultural journal* 1(2);55-58
11. Okeke S.E and Nwachukwu C.U 2001. Characterization of *Maesobotra barteri* var. *barteri*. *Nig. J. Bot.* 13:70-80.
12. Okwulehi I.C and Okoli B.E 1999. Morphological and Palynological studies in some species of *Corchorus* L. *Tiliaceae*, *New Botanist*: 25:87-101.
13. Olowokudejo J.D 1990. Comparative Morphology of leaf epidermis in the genus *Annona* (Annonaceae) in West Africa, *phytomorphol*, 40:407-422.
14. Shah G.L and Gopal B.V 1972. Some observation on the diversity of stomata and trichomes in some species of *Dioscorae*. *Ann. Bot.* 36:997-1004.
15. Stern K.R 2000. Introductory plant biology, Mac Graw Hill Company Inc. United Sataes of America. Pp 630.
16. Tindall H.D 1986. Vegetables in the Tropics (afford) Pp 35-38.
17. Vidyard, R.D and Tripathi S.C 2002. A text Book of Botany. S. Chand and Company Ltd. 7361, Ram Nagar, New Delhi 1054 Pp.