

The Antischistosomal Activity of *Fasciola gigantica* and *Schistosoma mansoni* Eggs is Influenced by Saponin Extracted from *Atriplex nummularia*

Amany Sayed Maghraby^a, Manal Abdel Aziz Hamed^{*b}, Hanan Farouk Aly^b, and Sanaa Ahmed Ali^b

^a Laboratory of Immunology and Infectious Diseases, Center of Excellence for Advanced Sciences, Therapeutic Chemistry Department, ^b Therapeutic Chemistry Department, National Research Center, Dokki, Cairo, Egypt.

*manal_hamed@yahoo.com

Abstract: The objective of the present study was to evaluate the antischistosomal, biochemical and humoral immune response of *Fasciola gigantica* and *Schistosoma mansoni* eggs homogenate influenced with or without saponin extracted from *Atriplex nummularia*. The work was extended to study the histopathological picture of the liver before and after challenge. Total worms reduction recorded 57.14, 80.95 and 42.85% in immunized mice with *Fasciola* egg homogenate (50µg/100µl PBS/mouse), *Fasciola* egg homogenate influenced by saponin (50µg/100µl PBS/mouse) and saponin alone (50µg/100µl PBS/mouse), respectively. Immunized groups with *Schistosoma* egg antigen and *Schistosoma* egg antigen influenced by saponin showed reduction in total worms by 47.61, 52.38%, respectively. In conclusion, immunization with *Fasciola gigantica* egg homogenate possesses promising antischistosomal properties with an immunomodulatory response to saponin. Heterologous homogenate had antischistosomal activity more than homologous homogenate. In addition, heterologous homogenate influenced with saponin had more antischistosomal activity than its homologous homogenate. Moreover, *Fasciola gigantica* egg homogenate had an immunoprophylactic effects by increasing the IgM and IgG levels against *Schistosoma* egg antigen. [Journal of American Science. 2010;6(8):368-381]. (ISSN: 1545-1003).

Keywords: *Fasciola gigantica*- *Atriplex nummularia*- saponin- *Schistosoma mansoni*- egg antigen-immunoprophylactic

1. Introduction

Vaccine strategies represent an essential component for the future control of schistosomiasis, as an adjunct to chemotherapy (McManus and Loukas, 2008). Immunization of *Schistosoma mansoni* (*S. mansoni*) infected mice by *Fasciola* egg antigens (SEA) showed cross-reactivity that protect against infection by reduction fluke burden and egg production (Hassan *et al.*, 2008a). Hassan *et al.* (2008b) demonstrated a structural homology in *Fasciola gigantica* excretory-secretory (E/S) and egg antigens. This homology was resided in the components of the similar molecular weights between both antigens. *Fasciola* / *Schistosoma* defined a cross-reactive antigen from *F. hepatica* worms that designated as FhSmIII. An antiserum of this antigen was developed and used as a probe to detect its presence (or its determinants) in different extracts of parasitic trematodes. In this manner, it was possible to demonstrate that FhSmIII was found in *S. mansoni*, *S. bovis* and *Paragonimus westermani* (Hillyer, 1984).

S. mansoni egg antigen (SEA) was found to contain numerous protein, polysaccharide, glycoprotein (Hassanein *et al.*, 1999) and enhances microsomal and mitochondrial enzymes including urea cycle enzymes (Kamel *et al.*, 1998). Vaccination

of mice with egg antigen showed delayed ovulation of ova in the liver and stools (Dunn *et al.*, 1988). In addition, Hillyer *et al.*, (1988) noticed that SEA appears to be good candidate in developing a screening assay for the immunodiagnosis of schistosomiasis. Infection of mice with a multiple doses of SEA down regulated egg deposition before *S. mansoni* infection (Botros *et al.*, 1999).

Saponin was constituted as one of the most distributed classes of secondary metabolite from plant or animal domains (Al-Habori and Rahman, 1998). They have broad spectrum of biological activity; cytotoxicity, antitumor, antioxidant and anti-inflammatory (Yanamandra *et al.*, 2003). Triterpenoid and steroid saponin have been found to be immunomodulator and detrimental to several infectious protozoan such as *Plasmodium* and *Leishmaniaia spp.* (Plock *et al.*, 2001). The addition of a new immunomodulator to the adjuvant-adaptation (ADAD) system with fatty acid binding proteins increased the protection against *F. hepatica* (Lopez-Aban *et al.*, 2008).

Saponin had a depressive effect against cancer cells by imitated specific cytokine inhibitors (Francis *et al.*, 2001) and detected specific toxic activity against macrophage (Kuroda *et al.*, 2001). Previous investigation on saponin revealed its role as

immunomodulator against *S. mansoni* infection (Maghraby *et al.*, 2007).

The objective of this study was to evaluate the influence of saponin on the anti-schistosomal activity of *F. gigantica* and *S. mansoni* egg antigens before and post challenge with 100 cercariae of *S. mansoni* through determination of certain biochemical and humoral immune responses. The histopathological analysis of the liver was also done to support our findings.

2. Materials and Methods

Female Swiss albino mice CD-1 strain (18 – 22g) were obtained from the Animal House, National Research Centre, Cairo, Egypt. They lived in controlled environment of constant temperature and humidity with freely access of water and food. *Atriplex nummularia* (family: *Chenopodaceae*) collected from Marsa Matroh Desert, Cairo, Egypt. Extraction of plant saponin was carried out according to Maghraby *et al.* (2007).

Antigens preparation

Cercariae, worm and egg antigens of *S. mansoni* as well as *F. gigantica* eggs used for enzyme linked immunosorbent assay (ELISA) technique were obtained from Theodore Bilharz Research Institute (Giza, Egypt). Immunization protocol was done according to Maghraby *et al.* (2007).

Ethics

Anesthetic procedures complied with the ethical guidelines approved by the Ethical Committee of the Federal Legislation and National Institutes of Health Guidelines in USA were approved by the Medical Ethical Committee of the National Research Centre in Egypt.

Immunization with *Schistosoma mansoni* eggs antigen

Mice were divided into eleven groups (5 mice/ group). Group 1, served as normal control group subcutaneously injected with 100µl of phosphate buffer saline (PBS). Groups 2-4 received 50 µg protein/ 100µl PBS of *S. mansoni* egg homogenate (SEA), saponin (SAP), SEA mixed with saponin (SEA+SAP), respectively and sacrificed 15 days post the 1st immunization. Groups 5-7 immunized with 50 µg protein/ 100µl PBS of SEA, SAP and SEA+SAP. At 15th day post the 1st immunization each group was booster by 2nd immunization with 50 µg / 100µl PBS of SEA, SAP,

SEA+SAP, respectively and sacrificed 15 days later. Group 8 served as positive control infected mice with 100 *S. mansoni* cercariae by tail immersion technique (Oliver and Stirewalt, 1952). Groups 9-11 immunized at 0 time with 50 µg / 100µl PBS of SEA, SAP and SEA+SAP, respectively. At 15 days post the 2nd immunization, mice were challenged with 100 *S. mansoni* cercariae and sacrificed after two months of infection.

Immunization with *Fasciola gigantica* eggs antigen

Vaccination regimens and animal grouping was carried out as described above, whereas *Fasciola* egg homogenate (FEA) and *Fasciola* egg homogenate influenced with saponin (FEA+SAP) were replaced all SEA groups.

Parasitological studies

Worms were recovered by liver perfusion as described by Smithers and Terry (1965). The percent of reduction in worm burden was calculated by the method of Tendler *et al.* (1968) as follows

$$P = (C - V) / C \times 100$$

Where, P was the percentage of protection; C was the mean number of parasite recovered from infected mice; and V was the mean number of parasite recovered from vaccinated mice. The relative sex ratio (RSR) was calculated by the method of Fallon *et al.* (1994) according to the formula:

$$RSR = [\text{Male/ female ratio in treated group}] / [\text{Male / female ratio in untreated group}]$$

The ratio of untreated groups was standardized as 1.

Enzyme linked Immunosorbent Assay (ELISA)

The assay was performed according to Hillyer *et al.* (1979). This assay was used for determination of IgM and IgG levels in immunized mice sera before and after infection with *S. mansoni* cercariae. Plates were coated with cercarial antigen preparation (CAP), soluble worm antigen preparation (SWAP), soluble egg antigens (SEA) and incubated at room temperature overnight. Orthophenylene diamin dihydrochloride (OPD) was used as a substrate. The reaction was read at 490 nm using Micro Well Plate Reader.

Liver tissue homogenate and biochemical analysis

The liver was removed, blotted, weight and homogenized in 4.5 volume of 0.1 hecadedecyle trimethyle ammonium bromide (CTB) for urea cycle enzymes determination. For estimation of succinate and lactate dehydrogenase enzymes and antioxidant

parameters, liver tissue was homogenized in 0.9 N NaCl (1:10 w/v).

Protein was estimated by the method of Bradford (1976). Lipid peroxides was determined as malondialdehyde. Its concentration was calculated using the extinction coefficient value $1.56 < 10^5 M^{-1} \text{ cm}^{-1}$ and read at 535 nm by the method of Buege and Aust (1978). Glutathione was estimated by the method of Moron *et al.* (1979) using dithiobis-2-nitrobenzoic acid (DTNB) in PBS. Catalase activity was assayed spectrophotometrically according to Nelson and Kiesow (1972). Superoxide dismutase (SOD) was estimated by the method of Nishikimi *et al.* (1972). Urea cycle enzymes; ornithine aminotransferase (OAT), argininosuccinate synthetase (ASS), argininosuccinate layase (ASL) and arginase (Arg.) were employed according to the method of Linton and Campell (1962). Estimation of lactate dehydrogenase (LDH) was carried out according to the method of Babson and Babson (1973). Succinate dehydrogenase (SD) was measured colorimetrically at 490 nm (Shelton and Rice, 1957).

Liver histopathology

Sections of liver tissue from each experimental group were fixed in 10% buffered formalin solution for histopathological studies. Paraffine embedded sections (5 μm thick) were taken after fixation. Slides were stained using haematoxylin and eosin (H&E) according to the method by Hirsch *et al.* (1997).

Statistical analysis

Statistics is carried out by one way analysis of variance (ANOVA), Costat Computer Program accompanied by least significance difference between groups at $P < 0.05$.

Antishistosomal activity and immunological determination will be carried out by independent *t*-test (Ronald *et al.*, 1983) and Graph pad instat software.

3. Results

Immunized groups with *Fasciola* egg homogenate (FEA) recorded reduction in total, male and female worms by 57.14, 42.10 and 46.15%, respectively with a relative sex ratio of 1.07. *Fasciola* egg homogenate influenced by saponin (FEA+SAP) showed reduction by 80.95, 73.68 and 76.92% as compared to infected group with relative sex ratio of 1.14. On the other hand, mice immunized by SAP showed diminution in total, male and female worms by 42.85, 31.57 and 15.38 %, respectively. Immunized groups with *Schistosoma* egg antigen (SEA) recorded reduction by 47.61, 26.31 and

38.46% in total, male and female worms, respectively. Attenuation in total, male and female worms post immunization with SEA+SAP was 52.38, 36.84 and 38.46 %. The relative sex ratio recorded 0.80, 1.19 and 1.02 for the last three antigens, respectively (Table 1).

Immunoprophylactic effect of *Fasciola gigantica* egg homogenate and *Schistosoma* egg antigen

Post 2nd immunization, the IgM level against SEA showed significant elevation in immunized group with FEA+SAP (Fig.1). Post 1st and 2nd immunization with SEA+SAP, the level of IgM against CAP showed a significant stimulation. SEA and SEA+SAP showed significant stimulation against SWAP. After challenge with *S.mansoni* cercariae, the level of IgM in immunized mice by SEA against SWAP showed a significant elevation, but insignificant stimulation was recorded with SEA+SAP as compared to the infected group (Fig.1).

Post first and second immunization with FEA and FEA+SAP as well as SEA, the levels of IgG showed significant elevation in immunized mice sera against CAP and SWAP. Post 2nd immunization with SEA+SAP, IgG level against CAP was significantly elevated. IgG level in immunized mice sera of SEA+SAP against SEA showed a significant elevation as compared to the infected group (Fig.2).

Urea cycle enzyme activities

After 1st immunization with saponin, FEA and FEA+SAP, significant increase in urea cycle enzymes; OAT and ASL were recorded. Significant decrease was observed in ASS and arginase as compared to normal group. After 2nd immunization with SAP, FEA and FEA+SAP, significant increase in OAT and significant decrease in ASS were noticed. On the contrary, ASS, ASL and arginase enzyme activities showed significant increase in all immunized infected groups (Table 2). Significant increase in all urea cycle enzyme activities after 1st and 2nd immunization with SEA+SAP was recorded. A significant decrease by 29.46% was observed in ASS after 1st immunization with SAP. Arginase enzyme activity showed significant decrease after 1st immunization with SAP and significant increase after 2nd immunization with SEA+SAP compared to normal mice. Meanwhile OAT exhibited significant increase in SAP vaccinated group and insignificant change in SEA and SEA+SAP groups (Table 3).

Glycolytic enzyme activities

Significant decrease was noticed in glycolytic enzymes (SD and LDH) after 1st and 2nd immunization with SAP, FEA and FEA+SAP (Table 3). Total protein recorded significant increase after 1st

immunization with FEA while significant decrease was noticed with saponin. After 2nd immunization with SAP, FEA and FEA+SAP a significant increase in total protein was recorded. The glycolytic enzyme; SD showed significant reduction after 1st immunization with SAP, SEA+SAP. LDH showed a significant reduction after 1st immunization with SAP. In 2nd immunization, significant decrease was noticed.

Antioxidant parameters

S. mansoni infected mice and all immunized groups recorded significant increase in lipid peroxides and superoxide dismutase levels, while significant decrease in glutathione and catalase levels were observed. In post challenged groups, the antioxidant levels recorded significant improvement in mice immunized by saponin. In addition, FEA and

FEA+SAP recorded more potent effect than SEA and SEA+SAP (Table 4).

Histopathological studies

Figures (3 and 4) showed histological profile of liver sections, arrow indicated normal hepatocytes (A). Immunized mice liver showed normal hepatic lobular structure and normal distribution of central vein and portal spaces with very mild lymphoid infiltration in few portal spaces (C, D, E, F, I, J, K, L). Liver of infected mice showed distribution of foci with acute inflammation destroying hepatic trabecular structure, portal space with biliary duct hyperplasia and chronic inflammatory infiltrations (B). Post challenged mice revealed remarkably reduced of granuloma size and count with minimal hepatic infiltrations (G, H, M, N).

Table 1: Reduction of total, male and female *S. mansoni* worms and relative sex ratio in immunized groups post challenge infection.

Groups	Total worm (TW)	Male (M)	Female (F)	%R (TW)	%R (M)	%R (F)	RSR
Infected	42.00 ± 4.52	19.00 ± 1.26	13.00 ± 1.82	-	-	-	1
<i>Schistosoma</i> eggs	22.00 ± 1.59	14.00 ± 1.41*	8.00 ± 1.16*	47.61	26.31	38.46	1.19
Saponin	24.00 ± 1.41*	13.00 ± 1.41*	11.00 ± 1.32*	42.85	31.57	15.38	0.80
<i>Schistosoma</i> eggs + saponin	20.00 ± 1.41*	12.00 ± 1.62*	8.00 ± 1.67*	52.38	36.84	38.46	1.02
<i>Fasciola</i> eggs	18.00 ± 1.85*	11.00 ± 1.41*	7.00 ± 1.72*	57.14	42.10	46.15	1.07
<i>Fasciola</i> eggs + saponin	8.00 ± 1.62*	5.00 ± 1.72*	3.00 ± 1.11*	80.95	73.68	76.92	1.14

- Values represented mean ±SD of 5 mice in each group.
- (*) Level of significance as compared to infected group at P<0.01
- % R is the percentage of reduction as compared to infected group.
- RSR is the relative sex ratio of each group, where $RSR = (\text{male} / \text{female})^{\text{treated}} / (\text{male} / \text{female})^{\text{infected}}$

Table (2): Change in urea cycle enzyme activities in immunized groups with *F. gigantica* and *S. mansoni* eggs homogenate influenced with and without saponin before and after challenge with *S. mansoni***Part A**

Groups	Immunized groups with <i>F. gigantica</i> eggs homogenate				Immunized groups with <i>S. mansoni</i> eggs homogenate			
	OAT	ASS	ASL	Arg.	OAT	ASS	ASL	Arg.
F / S 1st	6.09 ± 0.37 ^c	1.11 ± 0.08 ^d	27.72 ± 0.48 ^a	43.93 ± 0.74 ^{ef}	7.23 ± 0.66 ^{bc}	1.750 ± 0.23 ^{ab}	29.35 ± 0.90 ^b	47.43 ± 2.08 ^{bc}
SAP + (F/S 1st)	6.70 ± 0.24 ^b	1.19 ± 0.01 ^{bc}	23.97 ± 0.82 ^c	45.22 ± 1.17 ^{cd}	8.58 ± 0.67 ^{ab}	2.13 ± 0.26 ^a	31.24 ± 1.78 ^{ab}	47.85 ± 2.11 ^{bc}
F / S 2nd	7.31 ± 0.25 ^a	1.19 ± 0.03 ^{cd}	23.91 ± 0.46 ^c	44.40 ± 1.42 ^{de}	8.98 ± 0.69 ^a	1.83 ± 0.36 ^{ab}	30.15 ± 1.83 ^{ab}	50.33 ± 0.59 ^b
SAP + (F/S 2nd)	4.90 ± 0.30 ^e	1.95 ± 0.05 ^a	26.75 ± 0.36 ^b	46.67 ± 0.61 ^b	9.60 ± 0.49 ^a	2.04 ± 0.12 ^a	33.15 ± 0.85 ^a	55.98 ± 1.77 ^a
F / S + Infection	1.94 ± 0.103 ^h	0.79 ± 0.07 ^f	19.45 ± 0.98 ^e	22.43 ± 2.31 ^h	3.98 ± 0.89 ^e	1.06 ± 0.29 ^{bc}	21.36 ± 1.48 ^e	34.65 ± 3.90 ^e
SAP + F / S + Infection	3.1 ± 0.188 ^g	0.77 ± 0.06 ^g	20.35 ± 1.23 ^d	31.10 ± 1.52 ^g	5.20 ± 1.82 ^d	1.17 ± 0.11 ^{bc}	24.40 ± 3.55 ^{cde}	44.58 ± 3.80 ^{cd}

Part B

Common groups	OAT	ASS	ASL	Arg.
SAP 1st	6.65 ± 0.41 ^{bc}	0.91 ± 0.08 ^e	26.53 ± 0.39 ^b	42.93 ± 1.05 ^f
SAP 2nd	5.87 ± 0.19 ^d	1.20 ± 0.03 ^{bcd}	24.30 ± 1.02 ^c	46.75 ± 1.69 ^{bc}
Infection	3.50 ± 0.14 ^g	0.57 ± 0.10 ^h	2.04 ± 0.25 ^g	4.79 ± 0.71 ⁱ
SAP + Infection	7.48 ± 0.31 ^a	0.72 ± 0.02 ^g	16.61 ± 1.37 ^f	30.61 ± 1.73 ^g
Control	4.60 ± 0.26 ^f	1.29 ± 0.04 ^b	24.73 ± 0.26 ^b	49.39 ± 1.05 ^a

- Data are mean ± SD of five mice in each group.
- Values are expressed as: nmol/mg protein for lipid peroxide and catalase, µmol/g protein for superoxide dismutase, µg/mg protein for glutathione.
- OAT: ornithine aminotransferase, ASS: argininosuccinate synthetase, ASL: argininosuccinate layase, Arg: arginase, SAP: saponin, *F*: *Fasciola gigantica*, *S*: *Schistosoma mansoni*, 1st: first immunization, 2nd: second immunization.
- Unshared letters between groups are the significance values at p< 0.0001.
- Statistical analysis is carried out by one way analysis of variance (ANOVA), Costat Computer Program.

Table 3: Change in glycolytic enzyme activities and total protein content in immunized groups with *F. gigantica* and *S. mansoni* eggs homogenate influenced with and without saponin before and after challenge with *S. mansoni* infection.

Part A.						
Groups	Immunized groups with <i>F. gigantica</i> eggs homogenate			Immunized groups with <i>S. mansoni</i> eggs homogenate		
	SD	LDH	Total protein	SD	LDH	Total protein
FEA / SEA 1st	0.29 ± 0.01 ^e	96.27 ± 3.92 ^d	44.25 ± 0.95 ^e	1.17 ± 0.11 ^{ab}	145.25 ± 2.22 ^b	40.18 ± 0.78 ^{ef}
SAP+ (FEA/ SEA 1st)	0.32 ± 0.02 ^d	113.43 ± 2.33 ^b	38.25 ± 0.95 ^g	0.97 ± 0.05 ^{ab}	148.150 ± 1.39 ^{ab}	40.95 ± 0.98 ^e
FEA / SEA 2nd	0.32 ± 0.02 ^d	94.83 ± 2.88 ^j	70.50 ± 1.30 ^c	1.25 ± 0.09 ^{ab}	150.25 ± 6.65 ^{ab}	47.60 ± 2.28 ^d
SAP + (FEA / SEA 2nd)	0.24 ± 0.01 ^e	102.75 ± 2.22 ^c	80.25 ± 1.25 ^b	1.39 ± 0.06 ^a	155.40 ± 5.49 ^a	59.15 ± 3.00 ^b
FEA / SEA + Infection	1.65 ± 0.01 ^a	79.37 ± 3.00 ^h	29.75 ± 0.50 ^j	1.42 ± 0.07 ^a	91.41 ± 5.40 ^d	35.88 ± 4.09 ^f
SAP + FEA / SEA+ Infection	0.75 ± 0.01 ^c	84.58 ± 4.09 ^f	38.00 ± 0.82 ^g	1.39 ± 0.13 ^a	123.70 ± 9.96 ^c	45.01 ± 4.5 ^d
Part B.						
Common groups	SD	LDH	Total protein			
SAP 1st	0.27 ± 0.01 ^e	92.62 ± 1.36 ^e	36.75 ± 2.36 ^h			
SAP 2nd	0.29 ± 0.01 ^e	93.92 ± 3.87 ^e	54.50 ± 1.29 ^d			
Infection	0.10 ± 0.02 ^f	60.75 ± 0.47 ⁱ	83.25 ± 4.28 ^a			
SAP + Infection	1.03 ± 0.06 ^b	83.08 ± 1.47 ^g	35.75 ± 0.95 ⁱ			
Control	1.66 ± 0.04 ^a	143.93 ± 3.45 ^a	39.75 ± 1.71 ^f			

- Data are mean ± SD of five mice in each group.
- Values are expressed as µl mol/mg protein / min and total protein in mg protein / ml.
- SD: succinate dehydrogenase, LDH: lactate dehydrogenase, SAP: saponin, *F. Fasciola gigantica*, *S. Schistosoma mansoni*, 1st: first immunization, 2nd: second immunization.
- Unshared letters between groups are the significance values at p< 0.0001
- Statistical analysis is carried out by one way analysis of variance (ANOVA), Costat Computer Program.

Table 4: Antioxidant parameters in immunized groups with *F. gigantica* and *S. mansoni* eggs homogenate influenced with and without saponin before and after challenge with *S. mansoni* infection.

Part A.								
Groups	Immunized groups with <i>F. gigantica</i> eggs homogenate				Immunized groups with <i>S. mansoni</i> eggs homogenate			
	LPO	GSH	Cat	SOD	LPO	GSH	Cat	SOD
FEA / SEA1st	0.56 ± 0.01 ^{fg}	125.83 ± 3.11 ^b	22.82 ± 0.50 ^c	469.96 ± 2.72 ^h	0.56 ± 0.01 ^b	125.83 ± 3.11 ^b	22.82 ± 0.51 ^c	469.96 ± 2.72 ^h
SAP + (FEA / SEA 1st)	0.78 ± 0.05 ^c	117.67 ± 2.06	3.31 ± 0.03 ^c	10.66 ± 0.26 ^e	0.80 ± 0.02 ^e	113.04 ± 1.58	17.73 ± 0.98 ⁱ	481.78 ± 1.91 ^e
FEA / SEA 2nd	0.58 ± 0.02 ^e	120.51 ± 1.36 ^d	22.99 ± 0.59 ^c	474.02 ± 1.64 ^g	0.61 ± 0.02 ^d	117.82 ± 1.65 ^e	22.50 ± 0.03 ^d	476.75 ± 2.62 ^f
SAP + (FEA / SEA 2nd)	0.51 ± 0.02 ^g	125.66 ± 2.26 ^b	23.41 ± 0.56 ^b	466.68 ± 1.56 ⁱ	0.55 ± 0.02 ^f	122.20 ± 1.14 ^c	23.19 ± 0.02 ^b	470.46 ± 1.52 ^{gh}
FEA / SEA + Infection	0.75 ± 0.02 ^c	113.20 ± 2.32 ^f	20.21 ± 0.79 ^f	571.15 ± 6.08 ^c	0.78 ± 0.01 ^c	94.59 ± 1.43 ^j	20.05 ± 0.08 ^g	579.79 ± 2.33 ^c
SAP + FEA / SEA + Infection	0.56 ± 0.01 ^{fg}	125.83 ± 3.11 ^b	22.82 ± 0.50 ^c	469.96 ± 2.72 ^h	0.56 ± 0.01 ^b	125.83 ± 3.11 ^b	22.82 ± 0.51 ^c	469.96 ± 2.72 ^h
Part B.								
Common groups	LPO	GSH	Cat	SOD				
SAP 1st	0.65 ± 0.01 ^d	116.41 ± 3.48 ^f	21.63 ± 0.44 ^e	481.56 ± 4.10 ^e				
SAP 2nd	0.54 ± 0.02 ^g	123.73 ± 1.86 ^c	23.47 ± 0.53 ^b	468.88 ± 2.76 ^h				
Infection	0.93 ± 0.01 ^a	72.06 ± 1.55 ⁱ	15.81 ± 0.27 ^h	710.18 ± 6.71 ^a				
SAP + Infection	0.85 ± 0.02 ^b	97.87 ± 1.54 ^h	17.98 ± 0.33 ^g	620.32 ± 6.86 ^b				
Control	0.49 ± 0.02 ^h	129.18 ± 2.20 ^a	23.90 ± 0.18 ^a	463.25 ± 2.35 ^j				

- Data are mean ± SD of five mice in each group.
- Values are expressed as: nmol/mg protein for lipid peroxide and catalase, µmol/g protein for superoxide dismutase, µg/mg protein for glutathione
- SOD: superoxide dismutase, GSH: glutathione, LPO: lipid peroxides, Cat: catalase, SAP: saponin, *F*: *Fasciola gigantica*, *S*: *Schistosoma mansoni*, 1st: first immunization, 2nd: second immunization.
- Unshared letters between groups are the significance values at p<0.0001
- Statistical analysis is carried by one way analysis of variance (ANOVA), Costat Computer Program.




Fig. 1: Detection of IgM by ELISA in mice sera post the first immunization (black bars) and second immunization (grey bars) with *S. mansoni* or *F. gigantica* homogenates influenced with or without saponin and post challenge (white bars) with 100 *S. mansoni* cercariae. Plates were coated with CAP (a), SWAP (b), or SEA (c).




Fig. 2: Detection of IgG by ELISA in mice sera post the first immunization (black bars) and second immunization (grey bars) with *S. mansoni* or *F. gigantica* homogenates influenced with or without saponin and post challenge (white bars) with 100 *S. mansoni* cercariae. Plates were coated with CAP (a), SWAP (b), or SEA (c).




Fig. 3: Representative histological analysis photos for liver sections of different mice groups. (A) Normal structure of liver control mice [100x]. (B) infected group (positive control) [100x]. (C) FEA immunized group after 1st immunization [200x]. (D) FEA immunized group after 2nd immunization [200x]. (E) FEA and SAP immunized group after 1st immunization [200x]. (F) FEA +SAP immunized group after 2nd immunization [200x]. (G) FEA infected group [100x]. (H) infected FEA +SAP group [200x].




Fig. 4: Representative histological analysis photos for liver sections of different mice groups. (I) SEA group after 1st immunization [200x]. (J) SEA group after 2nd immunization [100x]. (K) SEA +SAP group after 1st immunization [200x]. (L) SEA +SAP group after 2nd immunization [100x]. (M) SEA-infected group [100x]. (N) SEA + saponin infected group [100x].

4. Discussion

Recent studies revealed that FEA enhances antibodies production supporting the hypothesis of immunomodulatory effect caused by vaccines (Schuster et al., 2007). Fasciola derived 12kd antigen developed antibodies that protected against *S. mansoni*, proofing its cross- reactivity (Cervi et al., 2004). Almeida et al. (2003) stated that *S. mansoni* (Sm14) induced high levels of protection against both *S. mansoni* and *F. hepatica*. Hillyer (2005) demonstrated that some vaccine candidates have anti-fecundity, anti-pathology, and anti-embryonation effects. El- Sayed et al. (2000); Noureldin et al. (2000) and Ruangittichai et al. (2006) reported that all cross-reactivity is ascertained by a significant protection of 57% against challenge. Our results recorded significant reductions ranged from 47.61: 80.95% in total worm of immunized mice with *Schistosoma*, *Fasciola* antigens and their combination with saponin. Yabe et al. (2008) and Maghraby et al. (2009) investigated interactions among *F. gigantica* and *Schistosoma* spp. In addition, Barduagni et al. (2008) found patients co-infected by *Fasciola* spp. and *S. mansoni*, while Abane et al. (2000) found that *F. hepatica* fatty acid binding protein (FABP) had been used in vaccination against *S. bovis*. López-Abán et al. (2008) reported that addition of a new immunomodulator to FABPs increase the protection against *F. hepatica*.

Our results showed that immunized mice with FEA+SAP and FEA caused a significant improvement in urea cycle enzymes, SD, LDH and total protein content. After challenged with *S. mansoni* infection, a significant reduction in urea cycle enzymes was observed. These results could be correlated to Mousa et al. (1975) and Senft (1976) who reported that bilharzial infection resulted in defect in protein metabolism through a defect in absorption of amino acid or defect in enzymes synthesis and derangement of many metabolic pathways which may involve detoxification mechanism in urea cycle regulation. Immunization with SEA or SEA+SAP showed a significant increase in OAT, ASS, ASL and an insignificant change was noticed in arginase, LDH and total protein content, while significant reduction was observed in SD enzyme activity. This can be explained on the basis that these antigen or metabolic product act on gene expression as a signal, so the transcription of DNA specific sequence into messenger RNA either be repressed or activated. Gene repression or activation is an effective way of changing enzyme activity (Hoek et al., 1997). Saponin showed a definite reactive immunomodulatory action and has a diuretic action causing increasing in OAT (Zhang et al., 1990; Plohmmon et al., 1997; Yamanandra et al., 2003). Al-

Habori and Raman (1998); Bogoiavlenskii et al. (1999) and Francis et al. (2001) showed that saponin modulate serum glycoprotein.

Pascal et al. (2000) and Hamed (2006) reported that oxidative stress due to schistosomiasis causes an elevation in lipid peroxides. In addition, Hamed (2006) showed that liver GSH was drastically depleted in *S. mansoni* infected mice and Gharib et al. (1999) attributed this decrease to the increased cytotoxicity with H₂O₂ produced as a result of inhibition of glutathione reductase that keeps glutathione in the reduced state. *S. mansoni* infection impairs the antioxidant system since the level of GSH depletion was used as an index of oxidative stress and a sign that hepatic cells are utilizing more antioxidant defenses (Ip, 2000). Therefore, all antigens used improved the level of antioxidants due to the reduction of schistosomal toxins evolved by worms which was confirmed through the observed reduction in worm burden. Catalase activity in our results recorded significant decrease after infection. Hahn et al. (2001) attributed this decrease to the oxidative stress which leads to peroxide radicals that detoxified by catalase and thus results in decline in its activity. Our results were in accordance with Son et al. (2007) who attributed the increase in superoxide dismutase to the enhancement of its mRNA expression as a result of exposure to superoxide and hydroxyl radicals.

Histopathological examination revealed that hepatic cells, central vein and portal triad were normal in immunized groups. Fibrous perihepatitis and hepatic parenchyma showed severe fibrosis and severe diffuse inflammatory infiltration, causing the distortion of hepatic lobules (Ali and Hamed 2006; EL-Banhawy et al., 2007). The current study showed that *S. mansoni* egg granuloma sizes were reduced after immunization with all antigens used. Saponin was maintaining liver architecture that in turn preventing the development of malignancies (Haridas et al., 2001).

In conclusion, *Fasciola* eggs, as a heterologous antigen, exerted more antischistosomal activity than the homologous *Schistosoma* one. In addition, combination of saponin enhanced the protective immunity of *Fasciola* eggs homogenate with more potent effect than *Schistosoma* eggs antigen.

Acknowledgment

The authors would like to thank Ass.Prof. Dr Kamel, H. Shaker, Natural Products Chemistry Department, National Research Center, Cairo, Egypt for supplying the extract of saponin.

This work was financially supported by grant no. 7020501, National Research Center, Cairo, Egypt.

*Corresponding author

Manal Abdel Aziz Hamed

Therapeutic Chemistry Department, National Research Center, Dokki, Cairo, Egypt.
manal_hamed@yahoo.com

5. References

- Abane, J. L., Oleaga, A., Ramajo V., Casanueva, P. Arellano, J.L., Hillyer, G.V., Muro, A. (2000) Vaccination of mice against schistosoma bovis with a recombinant fatty acid binding protein from Fasciola hepatica. *Veterinary Parasitology* 91, 33-42.
- Al-Habori, M., Raman, A. (1998) Antidiabetic and hypocholesterolaemic effects of fenugreek. *Phytotherapy Res* 12, 233-299.
- Ali, S. A., Hamed, M. A. (2006) Effect of *Ailanthus altissima* and *Zizyphus spina Christi* on bilharzial infestation in mice: histological and histopathological Studies. *Journal of Applied Science* 6, 1437-1446.
- Almeida, M. S., Torloni, H., Lee-Ho, P., Vilar, M. M., Thaumaturgo, N., Simpson, A., Tendler, M. J. (2003) Vaccination against Fasciola hepatica infection using a Schistosoma mansoni defined recombinant antigen, Sm14. *Parasit Immunology* 25, 135-137.
- Babson, A. I., Babson, S. R. (1973) Kinetic colorimetric measurement of serum lactate dehydrogenase activity. *Clinical Chemistry* 19, 766-769.
- [Barduagni, P.](#), [Hassanein, Y.](#), [Mohamed, M.](#), [Wakeel, A. E.](#), [Sayed, M. E.](#), [Hallaj, Z.](#), [Curtale, F.](#) (2008) Use of triclabendazole for treatment of patients co-infected by Fasciola spp. and Schistosoma mansoni in Behera Governorate, Egypt. [Parasitology Research](#) 102, 631-633.
- Bogoiavlenskii, A. P., Berezin, V. E., Ogneva, A.V., Tolmacheva, V.P., Dige, I. E., Khudiakova, S. S. (1999) Immunostimulating activity of a saponin-containing extract of saponaria officinalis. *Voprivrusol* 44, 229-232.
- Botros, S. S., Doughty, B.L., Shaken, Z.A., Aki, M.M., Sharmy, R., Diab, T.M., Hassanein, H.I. (1999) Efficacy of an antipathology vaccine in murine schistosomiasis administered with and without chemotherapy. *International Journal of Immunopharmacology* 18, 707-718.
- Bradford, M. M. (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry* 72, 248-254.
- Buege, J. A., Aust, S. D. (1978) Microsomal lipid peroxidation. *Methods in Enzymology* 52, 302-310.
- Cervi, L., Borgonovo, E., Egea, M., Chiapello, L., Masih, D. (2004) Immunization of rats against Fasciola hepatica using crud antigens conjugated with Freund's adjuvant or oligodeoxy nucleotides. *Veterinary Immunology and Immunopathology* 97, 97-104.
- Dunn, D.W., Hillyer, G.V., Vazque, Z.G. (1988) Schistosoma mansoni cationic egg antigens (CEF6): Immunoserology with Oxamniquine-treated patients and involvement of CEF6 in the circumoval precipitin reaction. *American Journal of Tropical Medicine and Hygiene* 38, 508-514.
- El-Banhawey, M. A., Ashry, M. A., EL-Ansary, A. K., Ali, S. A. (2007) Effect of Curcuma longa or Parziquantel. On Schistosoma mansoni infected mice liver: Histological and Histochemical Studies. *Indian Journal of Experimental Biology* 45, 877-889.
- El-Sayad, M. H., Allam, A. F., Shchab, A. Y. (2000) The circumoval precipitin reaction in fasciola. Detection of shell pores by scanning electron microscopy. *Journal Egyptian Society of Parsitology* 30, 655-660.
- Fallon, P. G., Smith, P., Nicholls, T., Modha, J., Doenhoff, M. J. (1994) Praziquantel induced exposure of Schistosoma mansoni alkaline phosphatase drug-antibody synergy, which acts preferentially against female worm. *Parasitology* 16, 529- 535.
- Francis, G., Makkar, H. P. S., Becker, K. (2001) Effect of Quillaja saponins on growth metabolism, egg production and muscle cholesterol in individually reared Nile tilapia. *Comparative Biochemical Physiology* 129, 105-144.
- Gharib, B. Abd-Allah, O.M., Dessein, H., De-Reggi, M. (1999) Development of eosinophil peroxidase activity and concomitant alteration of antioxidant defenses in the liver of mice infected with Schistosoma mansoni. *Journal of Hepatology* 30, 594-602.
- Hahn, U. K., Bender, R. C., Bayne, C. J. (2001) Killing of Schistosoma mansoni sperocytes by hemocytes from resistant Biomphalaria glabrata: role of reactive oxygen species. *Journal of Parasitology* 87, 292- 299.
- Hamed, M. A. (2006) Excretory-secretory product of Fasciola hepatica worm protects against Schistosoma mansoni infection in mice. *Indian Journal of Experimental Biology* 44, 554-561.
- Haridas, V., Arntzen, C. J., Gutterman, J. U. (2001) [Avicins, a family of triterpenoid saponins from Acacia victoriae \(Benthams\), inhibit activation of nuclear factor-kappaB by inhibiting both its nuclear localization and ability to bind DNA.](#) *Proceeding of the National Academy of Sciences USA* 98, 11557-1156.
- Hassan, A. A., Abou-Zinadah, N. Y., Shalaby, H. A., Morsy, T. A. (2008a) Efficacy of some Fasciola gigantica antigens. *Journal of Egyptian Society of Parasitology* 38, 243-254.
- Hassan, A. A., El-Bahy, M. M., Abou-Zinadah, N.Y., Shalaby, H. A. (2008b) The diagnostic efficacy of Fasciola gigantica coproantigen in naturally infected cattle and buffaloes. *Journal of Egyptian Society of Parasitology* 38, 115-130.
- Hassanein, H., Kamel, M., Badawy, A., El-Ghorab, N., Abdeen, H., Zada, S., El-Ahwany E., Doughty B. (1999) Antimiracidial effect of recombinant glutathione S. transferase-26 and soluble egg antigen on immune responses in murine schistosoma mansoni. *Acta pathologica, microbiologica et immunologica Scandinavica* 107, 723-736.
- Hillyer, G.V. (1979) Can we vaccinate against schistosomiasis? *Federation Proceedings* 35, 2568- 2571.
- [Hillyer, G. V.](#) (1984) Immunity of schistosomes using heterologous trematode antigens, a review. *Veterinary Parasitology* 14, 263-283.
- Hillyer, G. V. (2005) Fasciola antigens as vaccines against fascioliasis and schistosomiasis. *Journal of Helminthology* 79, 241-247.
- Hillyer, G. V., Garcia R. M. I., Alicea H., Hernandez A. (1988) Successful Vaccination against murine Schistosoma mansoni infection with a purified 12kd Fasciola hepatica cross-reactive antigen. *American Journal of Tropical Medicine and Hygiene* 38, 103-110.
- Hirsch, C., Zouain, C. S., Alves, J. B., Goes, A.M. (1997) Induction of protective immunity and modulation of granulomatous hypersensitivity in mice using PIII, an anionic fraction of Schistosoma mansoni adult worm. *Parasitology* 115, 21-28.
- Hoek, R. M., Van Kestern, R. E., Smith, A. B., de-Jong-Brink, M., Geraerts, W. P. (1997) Altered gene expression in the brain caused by a trematode parasite: Neuropeptide enes are preferentially affected during parasitosis. *Proceeding National Academic Science* 94, 14072-14076.
- Ip, S., Yiu, H. Y., Kos, K. M. (2000) Differential effect of schisandrin B and dimethyl diphenyl bicarboxylate (DDB) on hepatic mitochondrial glutathione redox status in carbon tetrachloride intoxicated mice. *Molecular and Cellular Biochemistry* 205, 111- 114.
- Kamel, M., Abou-Shousha, T.S., Maghawry, D., Charmy, R., Hassanein, H.I. (1998) The immune responses associated with Schistosoma mansoni SEA. Induced granulomatus hyporesponsiveness in two different strains of mice. Internal conference on schistosomiasis, March, 15-19, organized by the schistosomiasis Research Project, Ministry of Health and Population, Cairo, Egypt.

32. Kuroda, M., Mimaki, Y., Hasegawa, F., Yokosuka, A., Sashida, Y., Sakagami, H. (2001) Steroidal glycosides from the bulbs of *Camassia leichtlinii* and their cytotoxic activities. *Chemical and Pharmaceutical Bulletin* 49, 726-731.
33. Linton, S. N., Campbell, J. W. (1962) studies of urea cycle enzymes in the terrestrial snail *Otala lactea*. *Archives of Biochemistry and Biophysics* 97, 360-369.
34. [López-Abán, J.](#), [Nogal-Ruiz, J. J.](#), [Vicente, B.](#), [Morrondo, P.](#), [Diez-Baños, P.](#), [Hillyer, G. V.](#), [Martínez-Fernández, A. R.](#), [Feliciano, A. S.](#), [Muro, A.](#) (2008) The addition of a new immunomodulator with the adjuvant adaptation ADAD system using fatty acid binding proteins increases the protection against *Fasciola hepatica*. *Veterinary Parasitology* 6, 176-181.
35. Maghraby, A.S., Shaker, K.H., Gaber, H. M. (2009) Cross-Reactivity of *Schistosoma mansoni* – *Fasciola gigantica* influenced by Saponin. *Zeitschrift für Naturforschung* 64c, 288-296.
36. Maghraby, A. S., Shaker, K. H., Zahran, H. G., El-Sherbiny, M. (2007) In vivo, the immunological effects of *Fasciola gigantica* worms homogenate mixed with saponin on mice infected with *Schistosoma mansoni*. *Journal of Medical Sciences* 7, 724-731
37. McManus, D.P., Loukas, A. (2008) Current status of vaccines for schistosomiasis. *Clinical Microbiology Reviews* 4, 225-42.
38. Moron, M. S., Depierre, J.W., Mannervik, B. (1979) Level of glutathione, glutathione reductase and glutathione-S-transferase activities in rat lung and liver. *Biochimica et Biophysica Acta* 582, 67-78.
39. Mousa, W., Kheir-Eldin, A. A., Elsehly, A., Mousa, A. H. (1975) Lipid pattern in bilharziasis hepatic fibrosis. *Egyptian Journal of Bilharziasis* 2, 83-93.
40. Nelson, D. P., Kiesow, L. A. (1972) Enthalpy of decomposition of hydrogen peroxide by catalase at 25 (with molar extinction coefficients of H₂O₂ solution in the (U.V.)). *Analytical Biochemistry* 49, 474-478.
41. Nishikimi, M., Rae, N. A., Yagi, K. (1972) The occurrence of superoxide anion in the action of reduced phenazine methosulphate and molecular oxygen. *Biochemical and Biophysical Research Communication* 46, 849- 853.
42. Noureldin, M. S., El-Shinnawy, H., Abou Elenin, (2000) Serum pretreatment with *Schistosoma mansoni* antigens for serological diagnosis. *Journal of Egyptian Society of Parasitology* 30, 157-168.
43. Oliver, L., Stirewalt, M. A. (1952) An efficient method for the exposure of mice to cercariae of *Schistosoma mansoni*. *Journal of Parasitology* 39, 19- 23.
44. Pascal, M., Abd-Allah, O.M., El-Wali, N.E., Mergani, A., Qurashi, M.A., Magzoub, M., De-Reggi, M., Gharib, B. (2000) Hyaluronate levels and markers of oxidative stress in the serum Sudanese subjects at risk of infection with *Schistosoma mansoni*. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 94, 66-70.
45. Plock, A., Sokolowska, K.W., Presber, W. (2001) Application of flow cytometry and microscopical methods to characterize the effect of herbal drugs on *Leishmania* Spp. *Experimental Parasitology* 97, 141-153.
46. Plohmann, B., Bader, G., Hiller, K., Franz, G. (1997) Immunomodulatory and antitumoral effects of triterpenoid saponins. *Pharmazie* 52, 953-957.
47. Ronald T, Chapman S, Hall L (1983) *Statistics in Research Development*, pp. 264-300, The Chaucer Press Ltd. Bungay, Suffolk, N.Y. and London.
48. Ruangsittichai, J., Viyanant, V., Vichasri-Grans, S., Sobhon, P., Tesana, S., Upatham, E. S., Hofmann, A., Korge, G., Grams, R. (2006) *Opisthorchis viverrini*: identification of a glycine-tyrosine rich eggshell protein and its potential as a diagnostic tool for human opisthorchiasis. *International Journal for Parasitology* 36, 1329-1339.
49. Schuster, R. K., Heidrich, J., Paul, A., Nockler, K. (2007) Liver flukes in dogs and treatment with praziquantel. *Veterinary Parasitology* 150, 362-365.
50. Senft, A. W. (1976) Studies in arginine metabolism by *Schistosoma mansoni* or *Schistosoma haematobium*. *Comparative Biochemical Physiology* 21, 299-306.
51. Shelton, E., Rice, M.E. (1957). Comparison of the reduction of two tetrazolium salts with succinoxidase activity of tissue homogenates. *Journal of National Cancer Institute* 18, 117-125.
52. Smithers, L. E., Terry, R. J. (1969) The immunology of schistosomiasis. *Advances in Parasitology* 7, 41-93.
53. [Son, I. S.](#), [Kim, J. H.](#), [Sohn, H. Y.](#), [Son, K. H.](#), [Kim, J. S.](#), [Kwon, C. S.](#) (2007) Antioxidative and hypolipidemic effects of diosgenin, a steroidal saponin of yam (*Dioscorea* spp.), on high-cholesterol fed rats. [Bioscience Biotechnology and Biochemistry](#) 71, 3063-3071.
54. Tendler, M., Pinto, R. M., Oliveira, L. A., Gebara, G., Katz, N. (1968) *Schistosoma mansoni* vaccination with adult worm antigens. *International Journal of Parasitology* 16, 347 –352.
55. [Yabe, J.](#), [Phiri, I. K.](#), [Phiri, A. M.](#), [Chembensofu, M.](#), [Dorny, P.](#), [Vercruyse, J.](#) (2008) Concurrent infections of *Fasciola*, *Schistosoma* and *Amphistomum* spp. in cattle from Kafue and Zambezi river basins of Zambia. [Journal of Helminthology](#) 82, 373-376.
56. Yanamandra, N., Berhow, M.A., Konduri, S., Dinh, D.H., Oliver, W.C. Nicolson, G.L., Rao, J.S. (2003) Triterpenoids from *Glycyne max* decrease invasiveness and induce caspase-mediated cell death in human SNB19 glioma cells. *Clinical and Experimental Metastasis* 20, 375-383
57. Zhang, C. Yang, X., Xu, L. (1990) Immunomodulatory action of the total saponin of *Gynostemma Pentaphylla*. *Chung Hsi-chih Ho Tsa Chin* 10, 96-98.

7/7/2010