

**Synthesis, docking and anticancer activity of some new thienopyrimidine and thienooxazine derivatives**Eman Abd EL-maboud Fayed <sup>1\*</sup>, Taghreed Zakaria Shaver<sup>2</sup><sup>1</sup> Department of Organic Chemistry, Faculty of Pharmacy (Girls), Al-Azhar University, Cairo, Egypt<sup>2</sup> Department of Pharmaceutical Chemistry, Faculty of Pharmacy (Girls), Al-Azhar University, Cairo, Egypt[\\*alfayed\\_e@yahoo.com](mailto:*alfayed_e@yahoo.com)



**Abstract:** Synthesis of certain new fused 1,3-pyrimidine, 1,3-oxazine and 1,4-oxazepine derivatives, to produce new compounds of possible anticancer activity. Various series of 2-aryl-4,5,6,7-tetrahydrobenzo[*b*]thieno[2,3-*d*]pyrimidin-4(3*H*)-one **5 a-g** and 2-aryl-5,6,7,8-tetrahydrobenzo[*b*]thieno[3,2-*e*][1,3]oxazin-4(*H*)-one **8 a-d** were synthesized. Most of the synthesized compounds exhibited antitumor activity against human breast cancer cell line (MCF-7) and against human liver carcinoma (HepG-2) in vitro. All tested compound showed the anticancer activity against HepG-2 cells with IC<sub>50</sub> ranging from 6.04 to 43.5 µg.

[Fayed EA, Shaver TZ. **Synthesis, docking and anticancer activity of some new thienopyrimidine and thienooxazine derivatives.** *Nat Sci* 2014;12(12):171-181]. (ISSN: 1545-0740). <http://www.sciencepub.net/nature>. 23

**Keywords:** 1,3-pyrimidine; 1,3-oxazine and 1,4-oxazepine derivatives; anticancer activity

**1. Introduction**

Fused thieno[2,3-*d*]pyrimidine and its analogue oxazine derivatives has been widely reported over the last years. They were shown to possess antiviral <sup>1,2</sup> and antibacterial activity. <sup>3-6</sup> Their remarkable medicinal properties cover also anti-inflammatory <sup>7-9</sup> and antihistaminic <sup>10</sup> action. Moreover, among other thieno[2,3-*d*]pyrimidin-4-ones of these family, the compound 1 was recently identified as inhibitor of tumor cells proliferation. <sup>11</sup> Diversification of this scaffold using its cyclohexane moiety afforded another potent derivative 2 displaying IC 50 of 91 nM in the p21-deficient cell line <sup>12</sup> (**Chart 1**). The same core compounds are found to be phosphodiesterase PDE9 inhibitors, for example, **3**. <sup>13,14</sup> Such inhibitors can be applied for treating memory deficits that are associated with aging and neurodegenerative disorders such as Alzheimer's disease. <sup>15</sup>

**Chart 1****2. Material and Methods****Chemistry**

All melting point were taken on Electrothermal LA9000 series, Digital Melting point Apparatus were uncorrected. IR Spectra were determined using KBr disc technique on Nikolet IR 200 FT IR Spectrophotometer at Pharmaceutical Analytical Unit, Faculty of Pharmacy, Al-Azhar University, and values are represented in (cm<sup>-1</sup>). The <sup>1</sup>H-NMR Spectra was recorded on Varian Gemini EM-300MHz, NMR Spectrometer at Research Service Unit, Faculty of Science, Cairo University. DMSO-*d*<sub>6</sub> was used as solvents; Chemical shifts were measured in δ ppm, relative to TMS as internal standard. Mass Spectrum were recorded at 70 ev on DI-50 unit of Shimadzu GC/ MS- QP5050A Spectrometer at Regional Center for Mycology and Biotechnology (RCMB), At-Al-Azhar University represented as m/z (relative abundance %)( formula). Element Analysis (C,H,N) were carried out at Regional Center for Mycology and Biotechnology, Al-Azhar University, the values were found to be within ± 0.4 % of the theoretical ones unless otherwise indicated. Progress of the reaction was monitored by TLC using TLC sheets precoated with UV fluorescent silica gel Merck 60 F254 plates and was visualized using UV lamp.

**2-Cyano-2-cyclohexylideneacetamide (1)**

Prepared using reported procedure.<sup>16</sup>

**2-Amino-4,5,6,7-tetrahydro[1]benzothiophene-3-carboxamide (2)**

Prepared using reported procedure.<sup>17,18</sup>

**2-Styryl-5,6,7,8-tetrahydrobenzo[b]thieno[2,3-d]-pyrimidin-4(3H)-one (3)**

A mixture of compound **2** (1.96, 0.01 mol), cinnamaldehyde (1.32 g, 0.01 mol) in *N,N*-dimethylformamide (10 ml) in presence of KOH was heated under reflux for 6 h. The reaction mixture was cooled, poured onto ice, filtered, washed with water, then dried and crystallized from ethanol to give compound **3**; yield 80%; m.p.: 292-294°C, IR: 3166 (NH); 3035 (Ar-H); 2922, 2847 (aliphatic C-H); 1652 (C=O); 748, 681 (phenyl), <sup>1</sup>HNMR: 1.79 (m, 4H, 2CH<sub>2</sub> at positions 6, 7); 2.73 (s, 2H, CH<sub>2</sub> at positions 5); 2.88 (s, 2H, CH<sub>2</sub> at positions 8); 6.95-7.00 (d, 2H, CH=CH); 7.21-7.90 (m, 5H, Ar-H); 12.30 (s, 1H, NH exchangeable by D<sub>2</sub>O), MS (m/z %): 309 (21.56) (M+1), 308 (84.76) (M<sup>+</sup>), 179 (40.85) (C<sub>9</sub>H<sub>9</sub>NOS<sup>1+</sup>), 77 (48.76) (C<sub>6</sub>H<sub>5</sub><sup>1+</sup>), 43 (100) (HNCO<sup>1+</sup> &/or CH<sub>3</sub>CO<sup>1+</sup>). Anal. Calcd for C<sub>18</sub>H<sub>16</sub>N<sub>2</sub>OS; C, 70.10; H, 5.23; N, 9.08; Found: C, 70.34, H, 5.34, N 9.21.

**2-(4-Morpholinobenzylideneamino)-4,5,6,7-tetrahydrobenzo[b]thiophene-3-carboxamide (4a) and 2-(4-(piperazin-1-yl)benzylideneamino)-4,5,6,7-tetrahydrobenzo[b]thiophene-3-carboxamide (4b)**

**General procedure:**

A mixture of compound **2** (1.96, 0.01 mol), respective aldehyde (0.01 mol) in *N,N*-dimethylformamide (10 ml) in presence of KOH was stirred at room temperature for 2 hours. The product precipitated was filtered, washed with water, then dried and crystallized from ethanol to give compound **4 a& b**; For **4 a** yield 80%; m.p.: 243-244°C; IR: 3339, 3174 (NH<sub>2</sub>); 3050 (Ar-H); 2916, 2846 (aliphatic C-H); 1650 (C=O); 812 (p-substituted phenyl), <sup>1</sup>HNMR: 1.70-1.76 (m, 4H, 2CH<sub>2</sub> at positions 5, 6); 2.64 (s, 2H, CH<sub>2</sub> at positions 4); 2.75 (s, 2H, CH<sub>2</sub> at positions 7); 3.71-3.75 (m, 8H, morpholino CH); 7.02-7.05 (d, 2H, Ar-H); 7.30 (s, 2H, NH<sub>2</sub> exchangeable by D<sub>2</sub>O); 7.73-7.76 (d, 2H, Ar-H), 8.36 (s, 1H, N=CH), MS (m/z %): 370 (17.89) (M+1), 369 (77.81) (M<sup>+</sup>), 352 (100) (C<sub>19</sub>H<sub>18</sub>N<sub>3</sub>O<sub>2</sub>S<sup>1+</sup>), 191 (77.25) (C<sub>10</sub>H<sub>7</sub>O<sub>2</sub>S<sup>1+</sup>), 77 (13.14) (C<sub>6</sub>H<sub>5</sub><sup>1+</sup>). Anal. Calcd for C<sub>20</sub>H<sub>23</sub>N<sub>3</sub>O<sub>2</sub>S; C, 65.01; H, 6.27; N, 11.37; Found: C, 65.28, H, 6.31, N, 11.52.

For **4 b** yield 75%; m.p.: 244-245°C; IR: 3320, 3186 (2 NH); 3073 (Ar-H); 2930, 2850 (aliphatic C-H); 1660 (C=O); 885 (p-substituted phenyl), <sup>1</sup>HNMR: 1.70 (m, 4H, 2CH<sub>2</sub> at positions 5, 6); 2.26 (s, 1H, NH (piperazino) exchangeable by D<sub>2</sub>O); 2.66 (s, 2H, CH<sub>2</sub> at positions 4); 2.72 (s, 2H, CH<sub>2</sub> at positions 7); 2.88 (s, 4H, 2CH<sub>2</sub>NH (piperazinyl CH)); 3.30 (s, 4H,

2CH<sub>2</sub>N (piperazinyl CH)); 6.91-8.27 (m, 4H, Ar-H); 8.36 (s, 1H, N=CH); 8.55 (s, 1H, NH<sub>2</sub> exchangeable by D<sub>2</sub>O). Anal. Calcd for C<sub>20</sub>H<sub>24</sub>N<sub>4</sub>OS; C, 65.19; H, 6.56; N, 15.20; Found: C, 65.32, H, 6.63, N, 15.37.

**2-Aryl-4,5,6,7-tetrahydrobenzo[b]thieno[2,3-d]-pyrimidin-4(3H)-one 5 a-g**

**General procedure:**

The products were prepared by 2 methods.

**Method 1:**

The products were prepared starting from compounds **4** (0.01 mol) by refluxing in DMF (10 ml) in the presence of KOH for 5 hours. The product formed was filtered while hot dried and crystallized from ethanol yielded **5 b & c**.

**Method 2:**

A mixture of compound **2** (1.96, 0.01 mol), respective aldehyde (0.01 mol) in *N,N*-dimethylformamide (DMF) (10 ml) in presence of KOH was refluxed for 6 hours. The product precipitated was filtered, washed with water, then dried and crystallized from ethanol to give compound **5 a, d- g**.

**2-(4-(Piperidin-1-yl)phenyl)-4,5,6,7-tetrahydrobenzo[b]thieno[2,3-d]pyrimidin-4(3H)-one 5 a**

Compound **5 a** was prepared as described from **IV** and 4-(piperidin-1-yl)benzaldehyde. Yield 77 %; m.p.: >300 °C; IR: 3180 (NH); 3071 (Ar-H); 2927, 2847 (aliphatic C-H); 1657(C=O); 860 (p- substituted phenyl); <sup>1</sup>HNMR: 1.58 (s, 6H, 3CH<sub>2</sub> (piperidinyl CH)); 1.79 (s, 4H, 2CH<sub>2</sub>); 2.72 (s, 2H, CH<sub>2</sub>); 2.88 (s, 2H, CH<sub>2</sub>); 3.3 (s, 4H, 2CH<sub>2</sub> (piperidinyl CH)); 6.95-6.98 (d, 2H, Ar-H, J=9Hz); 7.99-8.02 (d, 2H, Ar-H, J=9Hz); 12.14 (s, 1H, NH exchangeable by D<sub>2</sub>O); MS m/z: 365 (1.32) (M<sup>+</sup>), 57 (100) (C<sub>2</sub>H<sub>5</sub>S<sup>1+</sup> &/or C<sub>2</sub>H<sub>3</sub>NO<sup>1+</sup>), 43 (89.81) (HNCO<sup>1+</sup> &/or CH<sub>3</sub>CO<sup>1+</sup>). Anal. Calcd for C<sub>21</sub>H<sub>23</sub>N<sub>3</sub>O<sub>3</sub>S; C, 69.01; H, 6.34; N, 11.50 Found: C, 69.17; H, 6.42; N, 11.67.

**2-(4-(Piperazin-1-yl)phenyl)-4,5,6,7-tetrahydrobenzo[b]thieno[2,3-d]pyrimidin-4(3H)-one 5 b**

Compound **5 b** was prepared as described from **IV** and 4-(piperazin-1-yl)benzaldehyde. Yield 75 %; m.p.: >300 °C; IR: 3275 (br 2NH); 3050 (Ar-H); 2923, 2834 (aliphatic C-H); 1646 (C=O); 816 (p-substituted phenyl); <sup>1</sup>HNMR: 1.78 (s, 4H, 2CH<sub>2</sub>); 2.10 (s, 2H, CH<sub>2</sub>); 2.49 (s, 2H, CH<sub>2</sub>); 2.73 (s, 4H, 2CH<sub>2</sub>-NH piperazinyl CH); 2.89 (s, 4H, 2CH<sub>2</sub>-N piperazinyl CH); 7.04-7.07 (d, 2H, Ar-H J= 9 HZ); 8.07-8.10 (d, 2H, Ar-H J= 9 HZ); 8.54 (s, 2H, 2NH exchangeable by D<sub>2</sub>O); MS m/z: 366 (0.5) (M<sup>+</sup>), 307 (12.90) (C<sub>18</sub>H<sub>15</sub>N<sub>2</sub>OS<sup>1+</sup> &/or C<sub>16</sub>H<sub>11</sub>N<sub>4</sub>OS<sup>1+</sup>), 77 (37.93) (C<sub>6</sub>H<sub>5</sub><sup>1+</sup>), 43 (100) (HNCO<sup>1+</sup> &/or CH<sub>3</sub>CO<sup>1+</sup>). Anal. Calcd for C<sub>20</sub>H<sub>22</sub>N<sub>4</sub>OS; C, 65.55; H, 6.05; N, 15.29; Found: C, 65.78; H, 6.11; N, 15.41.

**2-(4-Morpholinophenyl)phenyl)-4,5,6,7-tetrahydrobenzo[b]thieno[2,3-d]pyrimidin-4(3H)-one 5 c**

Compound **5 c** was prepared as described from **IV** and 4-morpholinobenzaldehyde. Yield 76 %; m.p.: >300 °C, IR: 3318 (NH); 3056 (Ar-H); 2931, 2839 (aliphatic C-H); 1657 (C=O); 869 (*p*-substituted phenyl); <sup>1</sup>HNMR: 1.78 (s, 4H, 2CH<sub>2</sub>); 2.49 (s, 4H, 2CH<sub>2</sub>); 2.74 (s, 4H, 2CH<sub>2</sub>-O morpholino CH); 2.89 (s, 4H, 2CH<sub>2</sub>-N morpholino CH); 7.00-7.03 (d, 2H, Ar-H J= 9 Hz); 8.04-8.07 (d, 2H, Ar-H J= 9 Hz); 12.18 (s, 1H, NH exchangeable by D<sub>2</sub>O). Anal. Calcd for C<sub>20</sub>H<sub>21</sub>N<sub>3</sub>O<sub>2</sub>S; C, 65.37; H, 5.76; N, 11.44; Found: C, 65.49; H, 5.82; N, 11.57.

**2-(2-Tosylphenyl)-4,5,6,7-tetrahydrobenzo[*b*]-thieno[2,3-*d*]pyrimidin-4(3*H*)-one 5 d**

Compound **5 d** was prepared as described in method 2 from **III** and 2-tosylbenzaldehyde. Yield 67 %; m.p.: >300 °C, IR: 3327 (NH); 3046 (Ar-H); 2933, 2870 (aliphatic C-H); 1646 (C=O); 840, 668 (*p*- & *o*-substituted phenyl); <sup>1</sup>HNMR: 1.80 (s, 4H, 2CH<sub>2</sub>); 2.49 (s, 3H, CH<sub>3</sub>); 2.75 (s, 2H, CH<sub>2</sub>); 2.90 (s, 2H, CH<sub>2</sub>); 6.92-8.13 (m, 8H, Ar-H); 12.25 (s, 1H, NH exchangeable by D<sub>2</sub>O); MS m/z: 453 (0.5) (M+1), 298 (65.61) (C<sub>16</sub>H<sub>14</sub>N<sub>2</sub>O<sub>2</sub>S<sup>1+</sup>), 179 (100) (C<sub>9</sub>H<sub>9</sub>NOS<sup>1+</sup>), 151 (68.33) (C<sub>8</sub>H<sub>9</sub>NS<sup>1+</sup>), 43 (19.93) (HNCO<sup>1+</sup>&/or CH<sub>3</sub>CO<sup>1+</sup>). Anal. Calcd for C<sub>23</sub>H<sub>20</sub>N<sub>2</sub>O<sub>4</sub>S<sub>2</sub>; C, 61.04; H, 4.45; N, 6.19; Found: C, 61.25; H, 4.46; N, 6.32.

**2-(2-(4-Chlorophenylsulfonyl)phenyl)-4,5,6,7-tetrahydrobenzo[*b*]thieno[2,3-*d*]pyrimidin-4(3*H*)-one 5 e**

Compound **5 e** was prepared as described in method 2 from **III** and 2-(4-chlorophenylsulfonyl)benzaldehyde. Yield 87 %; m.p.: >300 °C, IR: 3198 (NH); 3059 (Ar-H); 2931, 2860 (aliphatic C-H); 1650 (C=O); 850, 653 (*p*- & *o*-substituted phenyl); <sup>1</sup>HNMR: 1.71-1.91 (m, 4H, 2CH<sub>2</sub>); 2.62 (s, 2H, CH<sub>2</sub>); 2.69 (s, 2H, CH<sub>2</sub>); 6.77-8.15 (m, 8H, Ar-H); 8.64 (s, 1H, NH exchangeable by D<sub>2</sub>O); MS m/z: MS m/z: 474 (1.50) (M+2), 300 (70.63) (C<sub>16</sub>H<sub>16</sub>N<sub>2</sub>O<sub>2</sub>S<sup>1+</sup>), 298 (41.24) (C<sub>16</sub>H<sub>14</sub>N<sub>2</sub>O<sub>2</sub>S<sup>1+</sup>), 179 (100) (C<sub>9</sub>H<sub>9</sub>NOS<sup>1+</sup>), 151 (44.90) (C<sub>8</sub>H<sub>9</sub>NS<sup>1+</sup>). Anal. Calcd for C<sub>22</sub>H<sub>17</sub>ClN<sub>2</sub>O<sub>4</sub>S<sub>2</sub>; C, 55.87; H, 3.62; N, 5.92; Found: C, 56.04; H, 3.60; N, 6.01.

**2-(4-(4-Chlorophenylsulfonyl)phenyl)-4,5,6,7-tetrahydrobenzo[*b*]thieno[2,3-*d*]pyrimidin-4(3*H*)-one 5 f**

Compound **5 f** was prepared as described in method 2 from **III** and 4-(4-chlorophenylsulfonyl)benzaldehyde. Yield 85 %; m.p.: >300 °C, IR: 3282 (NH); 3050 (Ar-H); 2955, 2850 (aliphatic C-H); 1663 (C=O); 820 (*p*-substituted phenyl); <sup>1</sup>HNMR: 1.78 (s, 4H, 2CH<sub>2</sub>); 2.73 (s, 2H, CH<sub>2</sub>); 2.88 (s, 2H, CH<sub>2</sub>); 6.85-8.16 (m, 8H, Ar-H); 8.41 (s, 1H, NH exchangeable by D<sub>2</sub>O); MS m/z: 474 (2.61) (M+2), 472 (5.05) (M<sup>+</sup>), 298 (100) (C<sub>16</sub>H<sub>14</sub>N<sub>2</sub>O<sub>2</sub>S<sup>1+</sup>), 297 (35.82) (C<sub>17</sub>H<sub>15</sub>NO<sub>2</sub>S<sup>1+</sup>), 270 (84.74) (C<sub>15</sub>H<sub>14</sub>N<sub>2</sub>OS<sup>1+</sup>), 151 (32.29) (C<sub>8</sub>H<sub>9</sub>NS<sup>1+</sup>),

120 (70.17) (C<sub>7</sub>H<sub>4</sub>S<sup>1+</sup>), 43 (65.92) (HNCO<sup>1+</sup>&/or CH<sub>3</sub>CO<sup>1+</sup>). Anal. Calcd for C<sub>22</sub>H<sub>17</sub>ClN<sub>2</sub>O<sub>4</sub>S<sub>2</sub>; C, 55.87; H, 3.62; N, 5.92; Found: C, 55.84; H, 3.68; N, 5.99.

**2-(2-(4-Bromophenylsulfonyl)phenyl)-4,5,6,7-tetrahydrobenzo[*b*]thieno[2,3-*d*]pyrimidin-4(3*H*)-one 5 g**

Compound **5 g** was prepared as described in method 2 from **III** and 2-formylphenyl 4-bromobenzenesulfonate. Yield 75 %; m.p.: >300 °C, IR: 3186 (NH); 3073 (Ar-H); 2930, 2850 (aliphatic C-H); 1660 (C=O); 885, 650 (*o*-, *p*-substituted phenyl); <sup>1</sup>HNMR: 1.78 (s, 4H, 2CH<sub>2</sub>); 2.73 (s, 2H, CH<sub>2</sub>); 2.89 (s, 2H, CH<sub>2</sub>); 6.65-8.23 (m, 8H, Ar-H); 9.50 (s, 1H, NH exchangeable by D<sub>2</sub>O). Anal. Calcd for C<sub>22</sub>H<sub>17</sub>BrN<sub>2</sub>O<sub>4</sub>S<sub>2</sub>; C, 51.07; H, 3.31; N, 5.41; Found: C, 51.24; H, 3.38; N, 5.39.

**2-Hydroxy-4,5,6,7-tetrahydro[1]benzothiophene-3-carboxamide (6)**

A mixture of compound **2** (1.96 g, 0.005 mol) was dissolved in a mixture of water (30 ml) and hydrochloric acid (4.5 ml of 37%w/v soln). A cold solution of sodium nitrite (0.35gm, 0.005 mol) in water (1ml) was added dropwise with stirring to the cooled amine solution (0-5°C) to yield the diazonium salt. Stirring was continued for 15 min to ensure complete reaction. Hydrolysis was done easily by adding an equal amount of water, heating in a warm water bath at 50 °C for 30 min. After evolution of N<sub>2</sub> ceased, the formed precipitate was filtered, dried and crystallized from absolute ethanol to yield the target hydroxy compound. m.p.: 179-180°C, IR: 3404 (OH); 2929, 2880 (aliphatic C-H); 1664 (C=O), <sup>1</sup>HNMR: 1.79 (s, 4H, 2CH<sub>2</sub>); 2.88 (s, 4H, 2CH<sub>2</sub>); 14.96 (br. s, 1H, OH, exchangeable by D<sub>2</sub>O). MS m/z: 196 (5.62) (M-1); 179 (65.67) (C<sub>10</sub>H<sub>11</sub>OS<sup>1+</sup>); 151 (100) (C<sub>8</sub>H<sub>9</sub>NS<sup>1+</sup>). Anal. Calcd for C<sub>9</sub>H<sub>11</sub>NO<sub>2</sub>S; C, 54.80; H, 5.62; N, 7.10; Found: C, 54.97; H, 5.68; N, 7.24.

**6,7,8,9-Tetrahydrobenzo[*b*]thieno[3,2-*f*][1,4]oxazepine-2,5-dione (7)**

Chloroacetyl chloride (1.13gm, 0.8ml, 0.01mol) was added to a stirred solution of compound **6** (1.97, 0.01 mol) in *N,N*-dimethyl formamide (DMF) (20 ml), then the mixture was heated under reflux for 5 hours. The mixture poured onto ice cold water. The solid was then filtered dried and crystallized from ethanol. m.p.: 166-168°C, IR: 3185 (NH); 2926, 2870 (aliphatic C-H); 1662 (C=O), <sup>1</sup>HNMR: 1.81 (s, 4H, 2CH<sub>2</sub>); 2.90 (s, 4H, 2CH<sub>2</sub>); 3.81 (CH<sub>2</sub>-NH); 14.96 (br. s, 1H, NH, exchangeable by D<sub>2</sub>O). MS m/z: 237 (9.29) (M<sup>+</sup>); 206 (100) (C<sub>10</sub>H<sub>8</sub>NO<sub>2</sub>S<sup>1+</sup>); 179 (95.84) (C<sub>10</sub>H<sub>11</sub>OS<sup>1+</sup>); 151 (78.40) (C<sub>8</sub>H<sub>9</sub>NS<sup>1+</sup>). Anal. Calcd for C<sub>11</sub>H<sub>11</sub>NO<sub>3</sub>S; C, 55.78; H, 4.67; N, 5.90; Found: C, 55.84; H, 4.71; N, 5.99.

**2-Aryl-5,6,7,8-tetrahydrobenzo[*b*]thieno[3,2-*e*]-[1,3]oxazin-4(*H*)-one 8 a-d**

**General method:**

Respective benzoyl chloride (0.01mol) was added to a stirred solution of compound **6** (1.97, 0.01 mol) in *N,N*-dimethyl formamide (DMF) (20 ml) in presence of triethylamine (5drops), then the mixture was heated under reflux for 6 hours. The mixture cooled, poured onto ice cold water. The solid was washed with water, and then filtered dried and crystallized from ethanol.

**2-Phenyl-5,6,7,8-tetrahydrobenzo[*b*]thieno[3,2-*e*][1,3]oxazin-4(*H*)-one **8 a****

Compound **8a** was prepared as described in the previous method from **6** and benzoyl chloride. Yield 65 %; m.p. 97-100 °C, IR: 3070 (Ar-H); 2947 (aliphatic C-H); 1675 (C=O); 750 (phenyl); <sup>1</sup>HNMR: 1.76 (br s, 4H, 2CH<sub>2</sub>); 2.73 (s, 2H, CH<sub>2</sub>); 2.88 (s, 2H, CH<sub>2</sub>); 7.39-8.03 (m, 5H, Ar-H). MS m/z: 283 (0.76) (M<sup>+</sup>); 105 (14.38) (C<sub>8</sub>H<sub>9</sub><sup>+</sup>); 91 (10.74) (C<sub>7</sub>H<sub>7</sub><sup>+</sup>); 77 (16.72) (C<sub>6</sub>H<sub>5</sub><sup>+</sup>); 43 (100) (HNCO<sup>+</sup> &/or CH<sub>3</sub>CO<sup>+</sup>). Anal. Calcd for C<sub>16</sub>H<sub>13</sub>NO<sub>2</sub>S; C, 67.82; H, 4.62; N, 4.94, Found: C, 67.98; H, 4.68; N, 4.99.

**2-(4-ChloroPhenyl)-5,6,7,8-tetrahydrobenzo[*b*]thieno[3,2-*e*][1,3]oxazin-4(*H*)-one **8 b****

Compound **8b** was prepared as described in the previous method from **6** and 4-chlorobenzoyl chloride. Yield 80 %; m.p. 197 °C, IR: 3050 (Ar-H); 2930 (aliphatic C-H); 1671 (C=O); 834 (*p*-substituted phenyl); <sup>1</sup>HNMR: 1.90 (s, 4H, 2CH<sub>2</sub>); 2.73 (s, 2H, CH<sub>2</sub>); 2.88 (s, 2H, CH<sub>2</sub>); 7.41-7.94 (m, 4H, Ar-H). MS m/z: 317 (0.50) (M<sup>+</sup>); 156 (29.96) (C<sub>6</sub>H<sub>5</sub>NO<sub>2</sub>S<sup>+</sup>); 139 (100) (C<sub>8</sub>H<sub>10</sub>S<sup>+</sup> &/or C<sub>7</sub>H<sub>6</sub>OS<sup>+</sup>); 111 (50.04) (C<sub>6</sub>H<sub>5</sub>Cl<sup>+</sup>). Anal. Calcd for C<sub>16</sub>H<sub>12</sub>ClNO<sub>2</sub>S; C, 60.47; H, 3.81; N, 4.41; Found: C, 60.58; H, 3.79; N, 4.52.

**2-(4-FlouroPhenyl)-5,6,7,8-tetrahydrobenzo[*b*]thieno[3,2-*e*][1,3]oxazin-4(*H*)-one **8 c****

Compound **8c** was prepared as described in the previous method from **6** and 4-flourobenzoyl chloride. Yield 88 %; m.p.: 158-160 °C, IR: 3074 (Ar-H); 2932, 2860 (aliphatic C-H); 1678 (C=O); 848 (*p*-substituted phenyl); <sup>1</sup>HNMR: 1.74 (s, 4H, 2CH<sub>2</sub>); 2.73 (s, 2H, CH<sub>2</sub>); 2.98 (s, 2H, CH<sub>2</sub>); 6.56-8.05 (m, 4H, Ar-H). MS m/z: 301 (0.01) (M<sup>+</sup>); 123 (100) (C<sub>7</sub>H<sub>7</sub>S<sup>+</sup>); 95 (69.47) (C<sub>6</sub>H<sub>6</sub>F<sup>+</sup>). Anal. Calcd for C<sub>16</sub>H<sub>12</sub>FNO<sub>2</sub>S; C, 63.77; H, 4.01; N, 4.65; Found: C, 63.89; H, 4.09; N, 4.71.

**2-(2,4-DimethoxyPhenyl)-5,6,7,8-tetrahydrobenzo[*b*]thieno[3,2-*e*][1,3]oxazin-4(*H*)-one **8 d****

Compound **8d** was prepared as described in the previous method from **6** and 2,4-dimethoxy-benzoyl chloride. Yield 76 %; m.p.: 122 °C, IR: 3050 (Ar-H); 2963 (aliphatic C-H); 1690 (C=O); 845 (*p*-substituted phenyl). MS m/z: 344 (2.02) (M+1); 165 (100) (C<sub>9</sub>H<sub>9</sub>OS<sup>+</sup>); 153 (52.68) (C<sub>7</sub>H<sub>7</sub>NOS<sup>+</sup>). Anal. Calcd for C<sub>18</sub>H<sub>17</sub>NO<sub>4</sub>S; C, 62.96; H, 4.99; N, 4.08; Found: C, 63.14; H, 4.09; N, 4.71.

**2-(4-Chlorophenyl)-5,6,7,8-tetrahydrobenzo[*b*]thieno[2,3-*d*]pyrimidin-4(*1H*)-one (**9a**)**

A mixture of compound **2** (1.96g, 0.01mol) and *p*-chlorobenzoyl chloride (1.74 g, 0.01mol) in *N,N*-dimethyl formamide (DMF) (10 ml) and trimethylamine (3 drops) was heated under reflux for 6 hours. The reaction mixture was cooled; poured onto ice cold water, the formed solid was filtered, dried and crystallized from ethanol. m.p: 138-140 °C. Yield 71%; IR: 3150 (NH); 3085 (Ar-H); 2930, 2850 (aliphatic C-H); 1657 (C=O); 866 (*p*- substituted phenyl); <sup>1</sup>HNMR: 1.77 (s, 4H, 2CH<sub>2</sub>); 2.73 (s, 2H, CH<sub>2</sub>); 2.88 (s, 2H, CH<sub>2</sub>); 7.43-7.98 (m, 4H, Ar-H); 12.26 (s, 1H, NH exchangeable by D<sub>2</sub>O). Anal. Calcd for C<sub>16</sub>H<sub>13</sub>ClN<sub>2</sub>OS; C, 60.66; H, 4.14; N, 8.84; Found: C, 60.63; H, 4.09; N, 8.81.

**2-(4-Fluorophenyl)-5,6,7,8-tetrahydrobenzo[*b*]thieno[2,3-*d*]pyrimidin-4(*1H*)-one (**9b**)**

A mixture of compound **2** (1.96g, 0.01mol) and *p*-flourobenzoyl chloride (1.57 g, 0.01mol) in *N,N*-dimethyl- formamide (DMF) (10 ml) and trimethylamine (3 drops) was heated under reflux for 6 hours. The reaction mixture was cooled; poured onto ice cold water, the formed solid was filtered, dried and crystallized from ethanol. m.p: 86-87 °C. Yield 71%; IR: 3186 (NH); 3073 (Ar-H); 2930, 2870 (aliphatic C-H); 1660 (C=O); 885 (*p*- substituted phenyl); <sup>1</sup>HNMR: 1.78 (s, 4H, 2CH<sub>2</sub>); 2.73 (s, 2H, CH<sub>2</sub>); 2.89 (s, 2H, CH<sub>2</sub>); 6.96-7.95 (m, 4H, Ar-H); 8.60 (s, 1H, NH exchangeable by D<sub>2</sub>O). MS m/z: 301 (0.06) (M+1), 300 (0.08) (M<sup>+</sup>), 140 (66.19) (C<sub>6</sub>H<sub>8</sub>N<sub>2</sub>S<sup>+</sup>), 123 (100) (C<sub>7</sub>H<sub>7</sub>S<sup>+</sup>), 95 (66.98) (C<sub>6</sub>H<sub>5</sub>F<sup>+</sup>). Anal. Calcd for C<sub>16</sub>H<sub>13</sub>FN<sub>2</sub>OS; C, 63.98; H, 4.36; N, 9.33; Found: C, 64.00; H, 4.30; N, 9.37.

**Docking studies**

All docking studies were performed using "Internal coordinate Mechanics (Molsoft ICM 3.5-0a)" as follow:

**Generation of the structures of ligands and enzymes:**

2qu5<sup>19</sup> were downloaded through the Protein Data Bank PDB/ RCSB site (<http://www.rcsb.org/pdb/home/home.do>) and saved as \*.pdb file.

Molecular modeling of the target compounds were built using ChemDraw Ultra version 9.0.3 and minimized their energy through Chem3D Ultra version 9.0.3/ MOPAC, Jop Type: Minimum RMS Gradient of 0.010 kcal/mol and RMS distance of 0.1 °A, and saved as MDL MolFile (\*.mol).

**Conversion of PDB files (2qu5) into ICM objects:**

This conversion involves addition of hydrogen bonds, assignment of atoms types, and charges from the residue templates.

Click on Mol Mechanics/ Convert/ Protein, and then delete water molecules.

**Docking using Molsoft ICM 3.4-8C program:** <sup>20,21</sup>**Setting up docking project:**

Setting project name:

Clicking on Docking/ Set project name, pressing OK.

Setting up the receptor:

Clicking on Docking/ Receptor Setup, entering the receptor molecule in the receptor molecule data entry box (a\_\*), then clicking on *identify the binding sites* button to identify the potential ligand binding pockets, pressing OK.

After the receptor setup is complete, the program normally displays the receptor with selected binding site residues highlighted in yellow Xstick presentation.

Reviewing and adjusting binding sites:

ICM makes a box around the ligand binding site based on the information entered in the receptor setup section. The position of the box encompasses the residues expected to be involved in ligand binding. Clicking on the menu Docking/ Review/ Adjust ligand/ Box.

**Making receptor maps:**

The step now is to construct energy maps of the environment within the docking box by clicking on menu Docking / Make Receptor Maps and selecting the resolution of the map by entering a value into the grid cell size data entry box which is 0.5.

**Starting docking simulation:**

The interactive docking was used to dock one ligand at a time by clicking on menu Docking/ Interactive docking / Mol Table Ligand, using the drop down arrow to find the table of ligand and / or compounds we wish to dock, and then entering the thoroughness which represents the length of simulation. Generally 1 is reasonable value, selecting Calc. ICM Score, then selecting Display run which display the ligand sampling the energy in the ligand binding project.

**Displaying the result:**

Clicking Docking/ Browse/ Stack Conformations.

**Evaluation of the Antitumor Activity**

**Mammalian cell lines:** The cell lines that used in this study were Human breast cancer cell line (MCF-7 cells) and human hepatocellular cancer cell line (HepG<sub>2</sub> cells). All cell lines were obtained from Vacsera Tissue Culture Unit.

The mammalian cells were propagated in Dulbecco's modified Eagle's medium (DMEM) or RPMI-1640 depending on the type of cell line supplemented with 10% heat-inactivated fetal bovine serum, 1% L-glutamine, HEPES buffer and 50µg/ml gentamycin. All cells were maintained at 37°C in a humidified atmosphere with 5% CO<sub>2</sub> and were subcultured two times a week along experimentation.

**Antitumor activity evaluation using viability assay:**

The extracts or pure compounds were tested against two tumor cell lines i.e., Breast cancer cell line (MCF-7) and liver carcinoma cell line (HepG<sub>2</sub>). All the experiments concerning the cytotoxicity evaluation were performed and analyzed by tissue culture unit at the Regional Center for Mycology and Biotechnology RCMB, Al-Azhar University, Cairo, Egypt. Antitumor viability assay was carried according to the method described by (Saintigny *et al.*, 2004).

**Procedure:**

The tumor cell lines were seeded in 96-well plate in 100µl of growth medium at a cell concentration of  $1 \times 10^4$  cells per well. After 24 h of seeding, the monolayers were then washed with sterile phosphate buffered saline (0.01 M pH 7.2) and simultaneously the cells were treated with 100 µl from different dilutions of the test sample in fresh maintenance medium and incubated at 37°C. Serial two-fold dilutions of the tested compound (50, 25, 12.5, 6.25, 3.125 and 1.56 µg/ml) were added to confluent cell monolayers dispensed into 96-well, flat-bottomed microtiter plates (Falcon, NJ, USA) using a multichannel pipette. The microtiter plates were incubated at 37°C in a humidified incubator with 5% CO<sub>2</sub> for a period of 24 h. Untreated cells were served as controls. Three independent experiments were performed each containing six replicates for each concentration of the tested samples. The cytotoxic effects of the tested compounds were then measured using crystal violet staining viability assay. Briefly, after 24-h of treatment, the medium was removed, 100 µL of 0.5% of crystal violet in 50% methanol was added to each well and incubated for 20 minutes at room temperature and subsequently excess dye was washed out gently by distilled water. The plate was allowed to dry then the viable crystal violet-stained cells were lysed using 33% glacial acetic acid solution. Absorbance at 570 nm was then measured in each well using microplate reader (SunRise, TECAN, Inc, USA). Doxorubicin was used as positive control. The absorbance is proportional to the number of surviving cells in the culture plate. Thus, using this colorimetric procedure, the tested compounds-mediated cell lysis and the cytotoxic effect of doxorubicin (used as a positive control) were measured and compared to the viability of untreated cells.

Because the stock solutions to prepare the different concentrations from the tested compounds were solubilized in DMSO, controls with DMSO alone were performed in parallel for each concentration. <sup>22-26</sup>

**Data analysis**


The percentage cell viability was calculated using the Microsoft Excel®. Percentage cell viability was calculated as follows: according to the following calculation: the percentage of cell viability =  $[1 - (\text{ODt}/\text{ODc})] \times 100\%$ , where ODt is the mean optical density of wells treated with the tested compound and ODc is the mean optical density of untreated cells. The test compounds were compared using the IC<sub>50</sub> value, i.e., the concentration of an individual compound leading to 50% cell death that was estimated from graphical plots of surviving cells vs compound concentrations.

### 3. Results and Discussion

#### Chemistry

Reaction of cyclohexanone in dry benzene, with cyanoacetamide, ammonium acetate and glacial acetic acid<sup>27</sup> afforded 2-cyano-2-cyclohexylideneacetate (**1**). Cyclization of **1** with sulfur applying Gewald's reaction conditions<sup>16, 27</sup> yielded ethyl 2-amino-4,5,6,7-tetrahydro[1]benzothiophene-3-carboxylate (**2**). The main start **2** reacted in three pathways, firstly with cinnamaldehyde in dimethylformamide

DMF in the presence of potassium hydroxide (KOH) and reflux to afford compound **3**.<sup>28</sup> The presence of signals of aromatic protons in the <sup>1</sup>HNMR spectrum and the presence of NH absorption band in the IR spectrum confirmed the formation of compound **3**. The reaction was with 4-(piperazin-1-yl)benzaldehyde and 4-morpholino-benzaldehyde in the same solvent system and stirring at room temperature gave the target compounds **4 a&b** the presence of N=CH and aromatic protons in the <sup>1</sup>HNMR spectrum and the presence of NH<sub>2</sub> absorption bands in the IR spectrum confirmed the formation of open structure **4 a&b**. the compounds number **5 a-g** were synthesized by two ways firstly by refluxing compounds **4 a&b** in the same solvent system for 5 hours or by refluxing the compound **2** with substituted aldehydes in DMF and KOH for 6 hours. The absence of N=CH in addition to the presence of aromatic protons in the <sup>1</sup>HNMR spectrum and the presence of NH absorption bands in the IR spectrum explained the formation of closed **5 a-g**. (**Scheme 1**).



**Scheme 2** starting by diazotization<sup>29</sup> and hydrolysis of **2** brought about 2-hydroxy-4,5,6,7-tetrahydro[1]benzothiophene-3-carboxamide (**6**). The


reaction of compound **2** with chloroacetyl chloride gave 6,7,8,9-Tetrahydrobenzo[*b*]thieno[3,2-*f*][1,4]-oxazepine-2,5-dione (**7**). The presence of O=C-CH<sub>2</sub>-

NH in  $^1\text{H}$ NMR and the absence of broad OH in the IR spectrum confirmed the formation of compound **7**. Compound **6** also reacted with derivatives of benzoyl chloride afforded **8**. These compounds were confirmed by the presence of aromatic protons in the  $^1\text{H}$ NMR spectrum and the absence of OH and  $\text{NH}_2$  in  $^1\text{H}$ NMR spectrum and IR spectrum. Furthermore the compound **2** was reacted with benzoyl chloride derivatives to afford pyrimidine derivative which confirmed by spectral data. (Scheme 2)

#### Biological activity:

Out of fifteen compounds, nine compounds were selected depending on the docking study to be evaluated in the of **MCF-7** and **HepG-2** cells. All tested compound showed the anticancer activity against **HepG-2** cells with  $\text{IC}_{50}$  ranging from **6.04** to **43.5**  $\mu\text{g}$  (Table 3). Compound **9b** elicited the highest antitumor activity against the two cell lines with  $\text{IC}_{50}$  8.6 and 6.04  $\mu\text{g}$  against **MCF-7** and **HepG-2**

respectively. Studying the structure-activity-relationships of oxazine and pyrimidine derivatives (**8c,d** and **5a-f**) revealed that, oxazines showed higher activity than pyrimidines in both cells. Comparing anticancer activities of compounds **5a-f** where the pyrimidine ring is substituted with ortho or para substituted phenyl, compound **5e** with ortho 4-chlorophenylsulfonate group imparts higher activity than **5d** with ortho 4-tolylsulfonate group therefore; the electron withdrawing group in this position is favored for anticancer activity. Introduction of substitutedphenylsulfonate group in para position decrease the anticancer activity. Among compounds with heterocyclic ring **5a-c**, it was obvious that, compound **5a** bearing pipridine ring has slight better activity than **5b,c** having piprazine or morpholine ring .against **MCF-7** while piprazine and morphine increase the activity against **HepG-2** cells.



Scheme 2: Synthesis of compounds 6 -9



Table 1: % Viability of MCF-7 cells

compounds	Viability %					
	1.56 $\mu\text{g}$	3.125 $\mu\text{g}$	6.25 $\mu\text{g}$	12.5 $\mu\text{g}$	25 $\mu\text{g}$	50 $\mu\text{g}$
<b>5a</b>	100	99.17	90.23	90.45	82.76	<b>46.81</b>
<b>5b</b>	100	100	99.64	97.36	89.43	<b>61.48</b>
<b>5c</b>	98.72	96.81	91.46	80.69	72.41	<b>54.78</b>
<b>5d</b>	98.16	95.69	89.47	80.92	65.73	43.59
<b>5e</b>	96.81	90.76	78.34	43.71	29.42	<b>18.97</b>
<b>5f</b>	100	98.46	93.51	82.77	69.23	<b>57.12</b>
<b>8c</b>	99.04	96.87	89.26	76.34	34.65	<b>20.39</b>
<b>8d</b>	91.87	79.61	62.74	54.19	37.36	<b>22.78</b>

<b>9b</b>	94.82	83.78	59.43	36.29	24.78	<b>13.27</b>
-----------	-------	-------	-------	-------	-------	--------------

**Table 2: % Viability of HepG-2 cells**

compounds	Viability %					
	1.56 $\mu$ g	3.125 $\mu$ g	6.25 $\mu$ g	12.5 $\mu$ g	25 $\mu$ g	50 $\mu$ g
<b>5a</b>	100	100	98.18	89.52	81.46	<b>38.97</b>
<b>5b</b>	100	98.74	95.22	87.49	76.27	<b>31.64</b>
<b>5c</b>	100	99.08	96.56	88.73	69.17	<b>34.98</b>
<b>5d</b>	92.40	85.62	79.43	64.29	41.34	<b>23.75</b>
<b>5e</b>	83.64	69.88	57.76	34.19	16.32	<b>10.28</b>
<b>5f</b>	100	100	98.48	91.53	78.42	<b>37.98</b>
<b>8c</b>	95.24	89.26	66.72	42.89	31.05	<b>13.67</b>
<b>8d</b>	92.01	83.68	69.37	48.81	17.08	<b>9.64</b>
<b>9b</b>	92.57	78.34	47.91	23.17	12.94	<b>6.83</b>



**Figure 1: % Viability of MCF-7 cells****Figure 2: % Viability of HepG-2 cells****Table 3: IC<sub>50</sub> of the test set of compounds against human breast cancer cells MCF-7 and human hepatocellular carcinoma cell line HepG2**

Compounds	IC <sub>50</sub> ( $\mu$ g) MCF-7 cells	IC <sub>50</sub> ( $\mu$ g) HepG-2 cells
<b>5a</b>	47.8	43.5
<b>5b</b>	>50	39.7
<b>5c</b>	>50	39
<b>5d</b>	42.8	20.3
<b>5e</b>	11.4	8.31
<b>5f</b>	>50	42.6
<b>8c</b>	20.4	10.6
<b>8d</b>	15.6	12.1
<b>9b</b>	8.8	6.04

**VEGFR-2 (KDR) docking simulation:**

One of the key features of the cocrystal structure of some inhibitors bound KDR is the reorganization of Phe 1047 (of the Asp 1046, Phe 1047, Gly 1048 triad, the “DFG” motif) to induce the inactive “DFG-out” conformation. This conformational change inhibits the ability of the kinase to bind ATP productively and accommodates the binding of an appropriately substituted inhibitor into an extended lipophilic pocket.<sup>19</sup>

KDR requires other binding features including close proximity to the gate keeper site (Val 916) and hinge-region (Cys 919).<sup>30</sup>

**Figure 3: Cytotoxicity of 5e, 8c, 8d & 9b against human breast cancer cells MCF-7****Figure 4: Cytotoxicity of 5e, 8c, 8d & 9b against human hepatocellular carcinoma cell line HepG2**

In our investigation, the 3D-coordinates in X-ray structure of KDR in complex with the ligand, 4-

(2-(4-Chloro-3-(trifluoromethyl)phenyl)amino)-1*H*-benzimidazol-5-yl)oxy)-*N*-methyl-2-pyridinecarboxamide; (PDB code 2qu5)<sup>19</sup> were used as the receptor model in KDR docking simulation.


The docked model of **ligand** revealed ICM score of -98.1 and forms four H bonds with: Cys 919 (2 H bonds), with Glu 885 (strong H bond) and Asp1046 (1 H bond), (**table 4**).

On the other hand, the docking study of our target compounds (table 4) revealed that: All

compounds showed docking scores ranging from -65.37 to -96.55 Kcal/mol., most compounds formed hydrogen bonds with Glu 885 and Asp1046 similar to **ligand**. The most active compound, **9b** formed four hydrogen bonds inside the active site of VEGFR-2 with Asp1046, Glu 885 and Lys 868. (Figure5). Compound **5e** formed six hydrogen bonds with the five amino acids, (**Table 4**).

**Table 4: ICM Scores of the compounds, and hydrogen bonds formed with amino acid residues and their lengths**

Compounds	ICM scores	No. of hydrogen bonds	Involved group of amino acid	Atom of ligand involved	Length of hydrogen bond(A)
<b>Ligand</b>	-98.13	4	C919 hn	n4	2.51
			E885 oe2	h5	1.97
			C919 o	h12	2.05
			D1046 o	h4	2.43
<b>5a</b>	-75.74	2	D1046 hn	n1	<b>2.19</b>
			E885 oe2	h9	<b>2.13</b>
<b>5b</b>	-73.76	2	C919 hn	o1	1.89
			E885 oe2	h18	2.03
<b>5c</b>	-70.57	2	D1046 hn	n1	2.18
			E885 oe2	h9	2.13
<b>5d</b>	-96.55	2	D1046 hn	o3	2.45
			E885 oe2	h9	0.91
<b>5e</b>	-79.65	6	K868 hz1	o3	1.63
			D1046 hn	o2	1.27
			D1046 hn	n1	2.12
			F1047 hn	o4	0.87
			G1048 hn	o4	2.63
			E885 oe2	h9	1.80
<b>5f</b>	-88.49	2	D1046 hn	n1	1.60
			E885 oe2	h9	2.47
<b>8c</b>	-68.48	2	C919 hn	o2	1.97
			C919 hn	n1	2.68
<b>8d</b>	-65.37	2	C919 hn	o2	1.64
			D1046 hn	o3	2.78
<b>9b</b>	-66.69	4	K868 hz1	o1	<b>1.68</b>
			K868 hz2	o1	<b>1.99</b>
			D1046 hn	n1	<b>2.73</b>
			E885 oe1	h5	<b>2.68</b>



**Figure 5: The proposed binding mode of compound 5e in the binding site of KDR, created with Molsoft ICM-Pro software**




Figure 6: The proposed binding mode of compound 8c in the binding site of KDR, created with Molsoft ICM-Pro software




Figure 7: The proposed binding mode of compound 8d in the binding site of KDR, created with Molsoft ICM-Pro software




Figure 8: The proposed binding mode of compound 9b in the binding site of KDR, created with Molsoft ICM-Pro software

#### Corresponding Author:

Dr. Eman Abd EL-maboud Fayed  
Department of Organic Chemistry

Faculty of Pharmacy (Girls), Al-Azhar University,  
Cairo, Egypt

E-mail: [\\*alfayed\\_e@yahoo.com](mailto:*alfayed_e@yahoo.com)

#### References

1. Thakur, C. S.; Jha, B. K.; Dong, B.; Gupta, J. D.; Silverman, K. M.; Mao, H.; Sawai, H.; Nakamura, A. O.; Banerjee, A. K.; Gudkov, A.; Silverman, R. H. *Proc. Natl. Acad. Sci. U.S.A.* (2007), *104* (23), 9585–9590.
2. Silverman, R. *Chem. Abstr.* (2007), *147*, 496301; Patent WO 2007127212, 2007.
3. Kapustina, M. V.; Kharizomenova, I. A.; Shvedov, V. I.; Radkevich, T. P.; Shipilova, L. D. *Pharm. Chem. J.* (1992), *26* (1), 73–75.
4. Hafez, H. N.; El-Gazzar, A. B. *Bioorg. Med. Chem. Lett.* 2008, *18* (19), 5222–5227.
5. Balasubramanian, N.; Meena, K.; Nitin, J.; Avinash, D.; Chandrasekaran, S. *Bioorg. Med. Chem. Lett.* (2006), *16* (18), 4951–4958.
6. Prasad, M. R.; Prashanth, J.; Shilpa, K.; Kishore, D. P. *Chem. Pharm. Bull.* (2007), *55* (4), 557–560.
7. Alagarsamy, V.; Vijayakumar, S.; Solomon, V. R. *Biomed. Pharmacother.* (2007), *61* (5), 285–291.
8. Alagarsamy, V.; Shankar, D.; Solomon, V. R. *ARKIVOC* 2006, *16*, 149–159.
9. Manhas, M. S.; Sharma, S. D.; Amin, S. G. *J. Med. Chem.* (1972), *15*, 106–107.
10. Shishoo, C. J.; Shirsath, V. S.; Rathod, I. S.; Yande, V. D. *Eur. J. Med. Chem.* (2000), *35* (3), 351–358.
11. Wang, Y. D.; Johnson, S.; Powell, D.; McGinnis, J. P.; Miranda, M.; Rabindran, S. K. *Bioorg. Med. Chem. Lett.* (2005), *15* (16), 3763–3766.
12. Jennings, L. D.; Kincaid, S. L.; Wang, Y. D.; Krishnamurthy, G.; Beyer, C. F.; McGinnis, J. P.; Miranda, M.; Discifani, C. M.; Rabindran, S. K. *Bioorg. Med. Chem. Lett.* 2005, *15* (16), 4731–4735.
13. Gotanda, K.; Shinbo, A.; Nakano, Y.; Kobayashi, H.; Okada, M.; Asagarasu, A. *Chem. Abstr.* (2006), *146*, 81884; Patent WO 2006135080, 2006.
14. Gotanda, K.; Shinbo, A.; Nakano, Y.; Kobayashi, H.; Okada, M.; Asagarasu, A. *Chem. Abstr.* (2008), *149*, 734240; Patent WO 2008072778, 2008.
15. Van der Staay, F. J.; Rutten, K.; Barfacker, L.; DeVry, J.; Erb, C.; Heckroth, H.; Karthaus, D.; Tersteegen, A.; Van Kampen, M.; Blokland, A.; Prickaerts, J.; Reymann, K. G.; Schroder, U. H.; Hendrix, M. *Neuropharmacology* (2008), *55*, 908–918.

16. Gewald, k.; Z. Chem. 1962, 2, 305; through chem. Abstr. (1963), 58, 6770.
17. Gewald, K.; Schinke, E.; Böttcher, H.; *Chem Ber.*, (1966), 99, 94.
18. Chusheng, H.; Zhe, Z.; Shuhua, L.; Yulin, L.; *J. Chem. Res (S)*., (1999), 148.
19. Potashman, M. H., Bready, J., Coxon, A., DeMelfi, Jr., T. M., DiPietro, L., Doerr, N., Elbaum, D., Estrada, J., Gallant, P., Germain, J., Gu, Y., Harmange, J., Kaufman, S. A., Kendall, R., Kim, J. L., Kumar, G. N., Long, A. M., Neervannan, S., Patel, V. F., Polverino, A., Rose, P., Plas, S., Whittington, D., Zanon, R., and Zhao H., *J. Med. Chem.*, (2007), 50, 4351-4373.
20. Abagyan, R. A., Totrov M. M., *J. Mol. Biol.*, (1994), 235, 983-1002.
21. Abagyan, R. A., Totrov M. M., and Kuznetsov, D. N. *J. Comp. Chem.*, (1994), 15, 488-506.
22. Saotome K, Morita H, Umeda M., *Toxicol In Vitro*, (1989), 3(4):317-21.
23. Kueng W, Silber E, Eppenberger U., *Anal Biochem*, (1989), 182(1):16-19.
24. Gangadevi, V. and Muthumary, J., *African Journal of Biotechnology*. (2007), 6: 1382-1386.
25. Wilson, A. P. *A Practical Approach*, 3rd ed. (ed. Masters, J. R. W.) Oxford University Press, (2000).
26. Mosmann, T., *J. Immunol. Methods*; (1983), 65: 55-63.
27. Arya, V. P. *Indian J. Chem.*, (1975), 10, 1141.
28. Sergey G. Dzhavakhishvili,, Nikolay Yu. Gorobets,, Svetlana V. Shishkina, Oleg V. Shishkin, Sergey M. Desenko,† and Ulrich M. Groth, *J. Comb. Chem.* (2009), 11, 508–514
29. Stakey, G. W.; Villaescusa, F. W.; Wollner, T. E. *J. Org. Chem.*, (1965), 30, 4074.
30. Dinges, J., Ashworth, K. L., Akritopolou-Zanze, I. Arnold, L. D., Baumeister, S. A. Bousquet, P. F., Cunha, G. A., Davidsen, S. K. Djuric, S. W., Gracias, V. J., Michaelides, M. R., Rafferty, P., Sowin, T. J., Stewart, K. D., Xia Z. and Zhanga H. Q., *Bioorg & Med. Chem. Lett.*, (2006), 16, 4266–4271.

12/21/2014