

Assessment of Genetic diversity in populations of *Parkia biglobosa* in Derived and Guinea Savanna Zones of Nigeria using Nuclear Microsatellite markers (SSR)

¹*Usman, I. A ., ²Lawal, A., ²Onyekwelu, J.C. and ³James, G.O

¹Department of Forest Production and Products, Joseph Sarwuan Tarka University, Makurdi, Nigeria

³Department of Biochemistry, Joseph Sarwuan Tarka University, Makurdi, Nigeria

²Department of Forestry and Wood Technology, Federal University of Technology, Akure, Nigeria

Corresponding email: aukwublie@yahoo.com

Abstract: *Parkia biglobosa* (African locust bean) is a valuable multipurpose tree species widely used by local communities across West Africa for food, medicine, and income generation. Despite its importance, its natural populations are declining due to overexploitation and poor regeneration, putting its genetic resources at risk."This study assessed the genetic diversity of *P. biglobosa* in three populations (Benue, Kogi, and Nasarawa States), which were purposively selected based on the intensity of utilization and marketing of its products located in two savanna ecological zones (derived and guinea) of Nigeria, using nuclear simple sequence repeat (nSSR) markers. Leaf samples were collected, and genomic DNA was extracted from silica gel-dried leaf tissue. Polymerase Chain Reaction (PCR) was performed using five nSSR primers (Parpan3, Parpan4, Parpan9, Parpan13, and Parpan15). The amplified DNA fragments were separated using 2% agarose gel electrophoresis. Fragment sizes were estimated using ladder – DNA sizing marker and analyzed with CERVUS 3.0.7 software. The number of alleles per locus ranged from 2 to 7, observed heterozygosity ranged from 0.000 to 0.333, while expected heterozygosity ranged from 0.533 to 0.923 within populations. Polymorphic Information Content (PIC) values within populations ranged from 0.346 to 0.840, indicating that the markers were highly informative. Across populations and ecological zones, allelic richness ranged from 6.6 to 12.8, observed and expected heterozygosities ranged from 0.0000 to 0.0686 and 0.8339 to 0.9217, respectively. Despite high expectation, heterozygosity, observed heterozygosity was low, most likely due to the presence of null alleles and inbreeding arising from restricted gene flow within fragmented populations. The results show high genetic variation within and between populations, though not fully expressed at the phenotypic level. The SSR markers (Parpan 3, 9 and 13) used proved effective and are recommended for future genetic studies on *P. biglobosa*. However, broader sampling across the species range is needed for more comprehensive conclusions. This study highlights the need for urgent conservation and sustainable management strategies, including habitat protection and plantation development outside the natural range of the species.

[Usman, I. A ., Lawal, A., Onyekwelu, J.C. and James, G.O. **Assessment of Genetic diversity in populations of *Parkia biglobosa* in Derived and Guinea Savanna Zones of Nigeria using Nuclear Microsatellite markers (SSR).** *Nat Sci* 2026,24(1):1-11]. ISSN 1545-0740 (print); ISSN 2375-7167 (online). <http://www.sciencepub.net/nature> 01. doi:[10.7537/marsnsj240126.01](https://doi.org/10.7537/marsnsj240126.01)

Keywords: conservation; genetic diversity; management; *Parkia biglobosa*; PCR

1. Introduction

Parkia biglobosa (African locust bean tree) is one of the indigenous tree species used in agroforestry systems across Sub-Saharan African countries, due to its multiple uses. The species is a leguminous tree that belongs to the Fabaceae family and Mimosoideae subfamily (Amusa *et al.*, 2014; Houndonougbo *et al.*, 2020), which is widely distributed from Senegal in West Africa to Uganda in Central Africa and thrives in derived savanna, guinea savanna, and Sudan savanna climatic zones (Lompo *et al.*, 2018). Onyekwelu *et al.* (2015) listed some indigenous tree species that produce important fruit trees such as *Chrysophyllum albidium*, *Irvingia gabonensis* and *Garcinia kola* and other Non-Timber Forest Products (NTFPs) which are important to the populace.

Parkia biglobosa is an important multipurpose indigenous fruit tree that plays an important role in the

livelihoods of rural communities in Nigeria and beyond. It provides food, nutrition, medicine, income, and other ecosystem services. The seeds are fermented to produce "dawadawa" or "iru", a protein-rich traditional condiment used as a substitute for animal protein (Lamien *et al.*, 2011; Nyadanu *et al.*, 2017). Pods and fruit pulp are rich in proteins, vitamins (A, B, C), carbohydrates, lipids, and minerals (Nyadanu *et al.*, 2017; Agbani *et al.*, 2018), while the leaves, bark, and roots are used in traditional medicine to treat various ailments, including hypertension, malaria, and wounds (Ouedraogo, 1995; Dedehou *et al.*, 2016).

Besides its socio-economic benefits, *P. biglobosa* contributes to ecological functions such as soil fertility improvement through nitrogen fixation (Teklehaimanot, 2004). Despite its nutritional, economic and ecological importance, this species is genetically threatened as a result of overharvesting,

deforestation, bushfires, and unsustainable land-use practices. Natural of *P. biglobosa* is noted to be poor, and its populations are declining due to ageing of the trees and lack of adequate management (Ræbild *et al.*, 2012; Padakale *et al.*, 2015). Many fruit trees are currently being overexploited in their natural habitats due to high demand, continuous harvest, declining tree populations, ageing of existing trees, unsustainable farming systems, high human pressure, absence of livelihood options and poverty. The decline of fruit trees poses threats to food security as well as the environment, wildlife, traditional medicine and human livelihoods (Boboye *et al.*, 2021). The ongoing threats from land-use change, fruit tree overharvesting, and climate variability necessitate urgent investigation and conservation efforts.

Parkia biglobosa has been noted to exhibit moderate to high genetic diversity across its natural range in West and Central Africa (Amusa *et al.*, 2014; Lompo *et al.*, 2016). Preserving the genetic diversity of the species is fundamental for its adaptability, resilience, survival and future use in breeding and afforestation programs, especially in the face of increasing habitat fragmentation, overexploitation, and climate change. Genetic diversity enables a species to withstand environmental changes and stress, making it essential for sustainable forest management and conservation (Oluwajuwon *et al.*, 2022). Lawal *et al.* (2023) noted that genetic diversity connotes the richness of the hereditary information in the gene pool of a given species. However, sustainable utilization and effective conservation require a clear understanding of the genetic structure and variability of tree species for developing conservation strategies.

Molecular markers, especially **simple sequence repeats (SSRs)** have proven to be highly effective in assessing genetic diversity in plant populations. Utilizing microsatellites (SSR) for tree species is optimal because they are codominant, multiallelic, highly polymorphic and may reveal a dominant molecular marker of good potential and have been widely used due to their ability to detect detailed genetic differences (Ahmad *et al.*, 2018). Previous studies have applied SSR and other molecular markers to investigate the genetic diversity of *P. biglobosa* in West Africa, revealing a range of genetic diversity levels across regions (Amusa *et al.*, 2014; Adesoye and Apo, 2015; Lompo *et al.*, 2016; 2018; Popoola *et al.*, 2020). For this study, a nuclear SSR marker was used to assess variability within and between *P. biglobosa* populations because it has been proven effective for *P. biglobosa* populations as reported by Koura *et al.* (2019).

Despite the widespread distribution and utilization of *P. biglobosa* in Nigeria, there is still limited research on its genetic diversity in the country. The present

study was undertaken to evaluate the genetic diversity of *P. biglobosa* populations in the derived and guinea savanna ecosystems of Nigeria, specifically focusing on Benue, Kogi, and Nasarawa States due to the prevalence, utilization and marketing of *P. biglobosa* in them. The findings will support the conservation of *P. biglobosa* genetic resources, inform sustainable management strategies, and guide the selection of superior genotypes for domestication, afforestation, and breeding programs in Nigeria and beyond.

2. Materials and Methods

Study area

The study was carried out in three populations (Benue, Kogi and Nasarawa) of *P. biglobosa* from two ecological zones in Nigeria (Figure 1). Benue State is situated between Longitude 7° 4' and 10° 0' E and Latitude 6° 25' and 8° 8' N. The climate is tropical, with two distinct seasons: rainy season and dry season. The rainy season lasts from April to October, with an average annual rainfall of 1120 to 1500 mm. The dry season starts in November and ends in March (Alapa, 2019). The climate is characterized by high temperatures, with a temperature range of 26 - 37 °C. The relative humidity is dependent on the seasons but varies from 50 to 80%, with the dry season having the lowest and the rainy season having the highest. The vegetation is characterized by open woodland with tall trees that typically have short and wide leaves. Kogi State extends from Latitude 6.33 °N to 8.44 °N and from Longitude 5.40 °E to 7.49 °E. The Niger and Benue rivers, which converge at Lokoja, are the primary sources of drainage for the State. These rivers have vast floodplains. The State has two distinct seasons: the rainy season, which lasts from April to October, and the dry season, which lasts from November to March (Odekunle, 2004). Mean annual rainfall ranges from 800 to 1100 mm (Sani and Haruna, 2010), while mean daily temperature ranges from 24.4 to 33.8 °C. Relative humidity is highest during the rainy season, with a mean value of 82.2 and 61.1% during the dry season. The soil is characterized by the Cretaceous sediments and sandstone (Fatoye, 2018). Nasarawa State is situated in the central part of Nigeria; it is located between latitude 7 ° 45' and 9 ° 2 'N of the equator and between longitudes 7 ° and 9 ° 37'E of the Greenwich meridian. There are two distinct seasons: the rainy and the dry seasons. The rainy season commences at the beginning of May and lasts till October. The dry season is experienced between November and April. Mean annual rainfall varies from 1100 mm to 2000 mm. Daily temperature range between 20 and 34°C. Relative humidity ranges between 65% and 85 % during the rainy season and 25% and 55% during the dry season.




Figure 1: Map of central Nigeria showing the selected states ecological zones of study.

Collection of Plant material

Parkia biglobosa leaf samples from adult trees were collected from derived and guinea savanna ecosystems across Benue, Kogi and Nasarawa States (Figure 1 above). The leaves were cleaned, dried and preserved in silica gel in sealed and well-labelled zip-lock bags and brought to Inqaba Biotech West Africa laboratory in Ibadan, Nigeria and stored at -20°C for DNA Extraction.

Genomic DNA Extraction Protocol

Total genome DNA was extracted from 0.02g powdered silica gel dried *P. biglobosa* leaves put in a 20 ml tube using the plant mini extraction kit protocol according to the manufacturer's instructions at Inqaba Biotech Africa laboratory, Ibadan, Nigeria. The DNA quality and quantity were evaluated using 1% agarose gel electrophoresis in a nanodrop spectrophotometer and stored at -20°C for PCR amplification

PCR amplification and Electrophoresis

Five out of the ten nuclear microsatellite primers (SSR) developed for *Parkia panurensis* (Luettmann *et al.*, 2010) and transferable to *P. biglobosa* population study (Koura *et al.*, 2019) were selected and used for this study. The ten primer bases (F/R) were supplied by Inqaba Biotech West Africa laboratory with the code number: NG2024/55508. The locus name and sequences are presented in Table 1.

The PCR amplification was performed using a volume of 25 μl in a DNA thermal cycler (Eppendorf. master Master cycler Gradient). The reaction mixture contained 3 μl of genomic DNA, 12.5 μl of GoTag green master mix. 2x (Promega), 0.5 μl each of forward and reverse primers and 8.5 μl Nuclease free water. The cycling conditions were: 1

cycle at 94°C for 3 min; followed by 35 cycles at 94°C for 30s, 30s at 50°C annealing temperature, and extension at 68°C for 1 min., with final extension lasting 1min at 68°C. The PCR products were stored at 4 °C until analysis. Amplified PCR products were separated by electrophoresis using 2% agarose gel in 1 X tbe (Triss base EDTA) buffer, stained with 5ul ethidium bromide and visualized under uv light. Fragment size was estimated using a 50-10000bp bench Top PCR marker (ladder-DNA sizing marker).

Table 1: Characteristics of the nuclear microsatellite loci transferable by Koura *et al.* (2019) and used for this study

Primer	Primer sequence	Annealing Temperature(°C)	Repeat motif	Size range
Parpan 3	F:CACGTTAATTCAATCAAAAATGGTG R: TTTTGCCTTTTTTCGGACTTG	56.5	(GT) ₁₅	155-209
Parpan 4	F:TAMRATTGATGGGAGTGGGAAAAAG R: CAGGAGGTGGTCTCTTCAGG	54,0	GT ₁₃ (GA) ₁₅	148-210
Parpan 9	F: FAM-GGGGCTTGTGTCTCTCACTG R: ACTTTGAAGGCACGAGATGG	58.0	(AC) ₁₂	204-262
Parpan 13	F: TAMRA-CCTCCCTCGCTTCACAATC R: CACATGCAAATGAAAATGGTG	58.5	(GT) 17TT(GT) ₈	86-194
Parpan 15	F: HEX-TGGCCTCACTGCATACTGAC R: TGGGATGAACAAAACCTGTGC	55.0	(AC) ₂₄	104-152

Data Analysis

Scoring was done using the gel output files under Photo Light. Each primer pair was considered as ‘locus’ to refer to an nSSR site (defined by the terminal of a PCR primer pair), and the length of variants at a site was treated as an allele. Bright and scorable bands were traced and scored as discrete numbers (absent = 0, present = 1). The Cervus 3.0.7 software (Kalinowski *et al.*, 2007) was used to estimate genetic diversity parameters. The frequencies of the bands were used to calculate the number of alleles (A) per locus, number of individuals at the locus, observed (Hobs) and expected (Hexp) heterozygosity and polymorphic information content (PIC).

3. Results

Genetic diversity within the *P. biglobosa* population in the derived and the guinea savanna ecosystems

The genetic diversity of the *P. biglobosa* population within the derived and guinea savanna ecosystems is presented in Table 2. In the Benue population, the number of alleles (K), which is referred to as allelic richness, ranged from 2 to 7 per locus in the derived savanna and 3 to 6 in the guinea savanna, with a mean of 4.4 alleles. The observed heterozygosity (Hobs) was 0.000 across all loci. The expected heterozygosity ranged from 0.533 to 0.923 and 0.714 to 0.879 with a mean of 0.7694 and 0.8012 in the derived and guinea savanna ecological zones, respectively. The highest genetic diversity in terms of K, HExp in the two ecological zones was observed in parpan 13 (K=7, HExp=0.923), while the lowest value was recorded in parpan 4 (K=2, HExp=0.533). The Polymorphic Information content (PIC) (which is the primer’s discriminating power) value ranged from 0.346 to 0.840 in the derived savanna zone and 0.555 to 0.791 in the guinea savanna. The highest PIC was recorded in Parpan 13 (0.840), while the lowest value was obtained in Parpan 4 (0.346).

In the Kogi population, the allelic richness (K) ranged from 4 to 7 per locus in the derived savanna and 2 to 6 in the guinea savanna, with a mean number of 5.4 and 3.8 alleles, respectively. The observed heterozygosity (HObs) ranged from 0.000 – 0.333, with a mean of 0.117 in the derived savanna and 0.000 in the guinea savanna. The expected heterozygosity (HExp) ranged from 0.821 to 0.900 and 0.667 to 0.879 in the derived and guinea savanna, respectively, with respective means of 0.8719 and 0.7816 for the two ecological zones. The highest genetic diversity in terms of K, HExp, in the two ecological zones of Kogi State was observed in parpan 13 (K=7, HExp=0.900), while the lowest was recorded in parpan 4 (K=2, HExp=0.667). The Polymorphic Information Content (PIC) ranged from 0.671 to 0.825

in the derived savanna zone and 0.375 to 0.791 in the guinea savanna. The highest PIC was recorded in parpan 13 (0.825) while the lowest was in parpan 4 (0.375).

In the Nasarawa population, the **allelic richness (k)** ranged from 3 to 6 and 3 to 7 alleles across all loci, with a mean allelic richness of 4.4 and 4.6 within the derived and guinea savanna, respectively. The **observed heterozygosity (HObs)** was **extremely low (0.000–0.014)** across all loci in the two ecological zones. The **HExp** remained high (0.800–0.909 in derived savanna and 0.714 to 0.890 in guinea savanna across all loci). The highest genetic diversity was observed in parpan 3 (K=7; HExp=0.909) while the lowest was recorded in parpan 15 (K=3; HExp=0.714). The Polymorphic Information content (PIC) values were 0.593–0.810 in the derived savanna and 0.555–0.805 in the guinea savanna in all loci. The highest mean PIC was observed in Parpan 13 (0.810), while the lowest was obtained in Parpan 15 (0.555).

Table 2. Comparison of genetic diversity within the populations of *Parkia biglobosa* in the derived and guinea savanna ecological zones of Benue, Kogi and Nasarawa State

Population	Ecological zones	Derived Savanna					Guinea savanna				
		Locus	K	N	Hexp	Hobs	PIC	K	N	Hexp	HObs
Benue	Parpan 3	4	4	0.857	0.000	0.703	6	7	0.879	0.000	0.791
	Parpan 4	2	3	0.533	0.000	0.346	3	4	0.714	0.000	0.555
	Parpan 9	6	8	0.867	0.000	0.786	5	9	0.837	0.000	0.756
	Parpan 13	7	7	0.923	0.000	0.840	4	6	0.788	0.000	0.671
	Parpan15	3	6	0.667	0.000	0.535	4	6	0.788	0.000	0.671
Kogi	Parpan 3	7	7	0.923	0.000	0.840	5	6	0.848	0.000	0.744
	Parpan 4	4	3	0.867	0.333	0.671	2	2	0.667	0.000	0.375
	Parpan 9	4	4	0.821	0.250	0.667	3	3	0.800	0.000	0.593
	Parpan 13	7	8	0.900	0.000	0.825	6	7	0.879	0.000	0.791
	Parpan 15	5	6	0.848	0.000	0.744	3	4	0.714	0.000	0.555
Nasarawa	Parpan 3	6	7	0.802	0.014	0.715	5	5	0.899	0.000	0.768
	Parpan 4	3	3	0.800	0.000	0.593	3	3	0.800	0.000	0.593
	Parpan 9	4	4	0.857	0.000	0.793	7	7	0.890	0.014	0.805
	Parpan 13	6	6	0.909	0.000	0.810	5	7	0.791	0.000	0.700
	Parpan 15	3	3	0.800	0.000	0.593	3	4	0.714	0.000	0.555

K allelic richness, N =number of individuals typed, Hobs- Observed heterozygosity, HExp- Expected Heterozygosity, PIC =Polymorphic Information Content

Genetic Diversity between *P. biglobosa* populations

The genetic diversity of *P. biglobosa* populations in derived and guinea savanna ecosystems in Nigeria is presented in Table 3. The number of individuals sampled in each ecological zone varied from 34 in the derived savanna to 38 in the Guinea Savanna. The mean Allelic richness (K) was 12.8 in the derived savanna and 11 in the guinea savanna, which were not significantly different. The mean observed heterozygosity was significantly higher in the derived savanna (0.0460) than in the Guinea savanna (0.0106) at 5% probability (Table 3). The mean expected heterozygosity was 0.9217 in the derived savanna and 0.9129 in the guinea savanna, both of which were not significantly different.

The genetic diversity of the *P. biglobosa* population in Benue, Kogi and Nasarawa states is presented in Table 4. The number of individuals sampled from each population was 20 in Nasarawa State, 25 in Kogi State, and 27 in Benue State. Mean Allelic richness was equal in Kogi (7.8) and Nasarawa (7.8) states, both of which were higher than the mean alleleic richness in Benue (6.6), but mean alleleic richness was not significantly different across states. The mean observed heterozygosity was low across the States, with Kogi state (0.0686) having the highest value, followed by Nasarawa (0.0348). The lowest value (0.000) was recorded for Benue state. The mean expected heterozygosity obtained in the Nasarawa population (0.8860) was slightly higher than the mean expected heterozygosity recorded in

P. biglobosa populations in Kogi state (0.8803) and Benue (0.8339). However, the mean expected heterozygosity did not differ significantly across the three states.

Table 3: Comparison of genetic diversity between the *Parkia biglobosa* populations in the derived and guinea savanna ecological zones

Ecological zones	N	Number of loci	K	Hobs	HExp
Derived savanna	34	5	12.8 ^a	0.0460 ^a	0.9217 ^a
Guinea Savanna	38	5	11.0 ^a	0.0106 ^b	0.9129 ^a
p-value	-	-	0.2606	0.02554	0.2427

Note: Columns with different superscripts (alphabets) are statistically different at the 0.05 level. K allelic richness, N =number of individuals typed, Hobs- Observed heterozygosity, HExp- Expected Heterozygosity.

Table 4: Comparison of genetic diversity among different populations of *Parkia biglobosa*

Population	N	K	No. of loci	Hobs	HExp
Benue State	27	6.6 ^a	5	0.0000 ^b	0.8339 ^a
Kogi State	25	7.8 ^a	5	0.0686 ^a	0.8803 ^a
Nasarawa State	20	7.8 ^a	5	0.0348 ^a	0.8860 ^a
p-value	-	0.331	-	2.29e-05	0.265

Note: Columns with different superscripts (alphabets) are statistically different at the 0.05 level.

K= allelic richness, N =number of individual typed, Hobs- Observed heterozygosity, HExp- Expected Heterozygosity,

Discussion

Genetic diversity of *Parkia biglobosa* population

Genetic diversity is fundamental to sustaining the adaptive capacities of species, the loss of which can potentially compromise the survival potential of affected populations or species (Pauls *et al.*, 2013; Wang *et al.*, 2023). Molecular markers play a crucial role in modern plant breeding by allowing breeders to identify and track desirable traits to evaluate genetic diversity (Salgotra and Chauhan, 2023). Understanding genetic diversity is essential for conservation and effective breeding strategies in tree species like *P. biglobosa*. This study revealed valuable insights into the genetic structure of *P. biglobosa* populations across three states (Benue, Kogi, and Nasarawa) within two savanna ecological zones (guinea and derived) in Nigeria.

The allelic richness (number of alleles (K), which reflects genetic variability and long-term potential for adaptability and resilience) was lower in this study compared to previous reports by Lompo *et al.* (2016) (k=5-15) and Popoola *et al.* (2020) (17-50 in *Parkia biglobosa*). However, our results are in line with values presented for tree species like *Vitellaria paradoxa* (Abdulai *et al.*, 2012), k=4.06 in Ghana; *Parkia multifuga* (Phurailatpam *et al.*, 2022), k=5.33 in Manipur; *Khaya grandifolia* (Soares *et al.*, 2020), k=5.9 in Brazil and *Theobroma speciosum* (Dardengo *et al.*, 2018), k=7.88-8.67 in Brazil.

A notable result from this study is the observed low heterozygosity (Ho), particularly in the Benue population, where it was completely zero. This suggests a high level of homozygosity, likely caused by inbreeding, small population sizes, reduced or restricted gene flow, possibly as a result of habitat fragmentation or overexploitation. The lack of observed heterozygosity also suggests that individuals may be more genetically similar than expected under natural conditions. Similar trends have been reported in *P. biglobosa* populations facing anthropogenic pressure (Sina, 2006; Ho=0.26; Popoola *et al.*, 2020, Ho=0.28) and other species such as *Prunus sibirica* (Wang *et al.*, 2023, Ho=0.32) and *Adansonia digitata* (Abate *et al.*, 2025, Ho=0.065-0.077).

Despite observed low heterozygosity, high expected heterozygosity (He) values were recorded within and between populations, indicating a rich underlying gene pool. This pattern, where expected heterozygosity is high but observed heterozygosity is low, may reflect historical genetic diversity that is no longer fully expressed due to population pressures or recent environmental changes and outcrossing mechanisms, which are typical of insect-pollinated, outcrossing tree species like *P. biglobosa* (Lompo *et al.*, 2016). High expected heterozygosity have been recorded in *P. biglobosa* (Akin-Idowu *et al.*, (2018) (0.41-0.93); Lompo *et al.*, 2018 (0.79); and related species like *Fraxinus angustifolia* (0.72) (Temunovic *et al.*, 2013),

Fagus sylvatica (0.407-0.880) (Szasz-Len and Konnert, 2018), *Vitellaria paradoxa* (0.5-0.7) and *Adasonia digitata* (0.6-0.8) (Boboye *et al.*, 2021), *Celtis zenkeri* (0.78) (Olamidayo *et al.*, 2021). The results of Soares *et al.* (2020), $He=0.56$; Sexton *et al.* (2015), $He=0.64$ and Tsipidou *et al.* (2021), $He=0.622$ are within the range of this study. Several studies reported very low expected heterozygosity in *Theobroma speciosum* (Dardengo *et al.*, 2018; $He = 0.21-0.26$), *P. biglobosa*. (Sina, 2006; $He=0.34$), *Cunninghamia lanceolata* (Jin *et al.*, 2023; $He=0.233$), which contradicts our results.

The Polymorphic Information Content (PIC) values recorded in this study indicate that most of the nuclear SSR markers (Parpan 3, 9 and 13) used were highly informative and reliable for detecting genetic variation, as noted by Koura *et al.* (2019). These values are comparable to those reported by Popoola *et al.* (2020) (0.51-0.89) in *P. biglobosa* and higher than PIC values from other studies, like Attikora *et al.* (2024) (0.24) for *Fraxinus angustifolia*, Mohammed *et al.* (2022) (0.3092-0.5658) for *Vitellaria paradoxa* and Denis (2025) (0.45) for *Kigelia africana*.

The expected heterozygosity values in derived and guinea ecological zones and across the three states ranged from 0.8339 to 0.8860, indicating generally high genetic diversity as opined by de Lafontaine *et al.* (2013), who noted that the expected heterozygosity value range of 0.6-0.8 is high. Thus, the overall estimates of genetic diversity of *P. biglobosa* were high with low observed heterozygosity. Areas experiencing high deforestation and human pressure often display reduced observed heterozygosity despite retaining relatively high expected values. High heterozygosity is often linked with greater evolutionary potential, improved adaptability, and reduced risk of inbreeding (Odebunmi *et al.*, 2017; Frankham, 2022). However, the discrepancy between observed and expected heterozygosity in all populations of the species, especially in Benue, may point to an excess of homozygotes, which could reflect inbreeding, small population sizes, or habitat fragmentation (Hartl and Clark, 2007). It also suggests that, while the population harbours genetic potential, actual genetic variation is not fully expressed in the individuals observed. This trend highlights a broader issue of genetic erosion in tree species facing habitat loss, reduced pollination, and human exploitation.

Some previous studies (Adesoye *et al.*, 2013; Adesoye and Apo, 2015; Ouedraogo, 2015) have documented wide genetic variation in *P. biglobosa*, emphasizing the role of environmental factors and genetic makeup, which agrees with this study. However, if this diversity is not maintained through adequate management and conservation, the evolutionary potential of the species could be severely reduced. The

findings of this study contradict studies that reported low genetic diversity in *P. biglobosa* and other tropical species, such as Amusa *et al.* (2014) for *P. biglobosa*; Boboye *et al.* (2018) for *Chrysophyllum albidum*; Akinagbe *et al.* (2019) for *Mansonia altissima*; and Lawal *et al.* (2023) for Khaya species. It was observed in this study that populations of *P. biglobosa* in derived savanna showed slightly higher genetic diversity than those in Guinea savanna. This may be due to better gene flow, less habitat fragmentation, or historical differences in land-use. Katumo *et al.* (2022) noted the role of human activity, pollinator availability, and landscape structure in shaping the genetic diversity of plants.

The overexploitation of *P. biglobosa* could have threatened their genetic diversity and hence, could limit their conservation and evolutionary development of the remaining populations. Cui *et al.* (2022) reported that genetic diversity is a prerequisite for future adaptive change and evolution and has a profound effect for species conservation. Our results highlight the urgent need for conservation actions for *P. biglobosa* in the savanna regions of Nigeria, especially in Benue State, where genetic diversity is at risk. Maintaining high genetic diversity is crucial for the long-term survival of *P. biglobosa*, particularly under the growing threats of climate change, habitat loss, and unsustainable harvesting.

Conclusion

This study revealed high genetic diversity in *Parkia biglobosa* populations across the Guinea and Derived savannas of Benue, Kogi and Nasarawa States, despite low observed heterozygosity, especially in Benue State. The high expected heterozygosity suggests that the species retains considerable genetic potential. However, the discrepancy between observed and expected heterozygosity raises concerns about inbreeding, habitat fragmentation, and genetic erosion. The high polymorphism in most of the markers indicates that the nuclear SSR markers used are effective in detecting subtle genetic differences among populations, offering a powerful tool for monitoring genetic diversity and guiding breeding and afforestation programs. Therefore, maintenance of *P. biglobosa* genetic diversity is essential for its adaptive capacity, long-term survival, and sustainable use, given its economic and ecological value. There is an urgent need for in-situ conservation efforts, implementing restoration programs in more vulnerable populations like Benue state, where observed heterozygosity was completely absent. These identified genetic diversities can be prioritized in breeding programs to select high-performing individuals for seed production and propagation, especially populations from Nasarawa and Kogi states. Protecting the remaining natural habitats and

promoting regeneration and ensuring interconnected conservation corridors for gene flow become essential. Strategies such as promoting natural regeneration, protecting pollinators, and encouraging community-based conservation can help restore genetic stability and connectivity among populations.

Corresponding Author:

Usman, I.A.
Department of Forest Production and Products,
Joseph Sarwuan Tarka University, Makurdi, Nigeria
Telephone: +234 7034305481
E-mail: aukwublie@yahoo.com

References

- Abate, N.N., Degu, M.K and Abebe, T (2025). Inbreeding depression manifested in progeny from fragmented populations of the wind-pollinated Dioecious Conifer *Afrocarpus gracilior* (Pilg) CN page. *Ecol. Evol.* 15(2):e70903. doi: 10.1002/ece3.70903.
- Abdulai, I., Krutousley, K and Finkeldey, R (2012). Morphological and genetic diversity of shea tree (*Vitellaria paradoxa*) in the savanna regions of Ghana. *Genetic Resources and Crop Evolution*, 54(6):1253-1268.
- Adesoye, A. I. and Apo, K. A. (2015). Evaluating genetic biodiversity of *Parkia biglobosa* (Jacq.) R. Br. Ex Don (African locust bean) accessions from Nigeria using seed protein electrophoresis. *Journal of Crop Science and Biotechnology* 18(3):171–180.
- Adesoye, A. I., Ogunremi, C. O and Aina, O. O. (2013). Genetic variation and heritability of seedling traits in Africa locust bean *Parkia biglobosa* (Jacq.) R. Br. ex G. Don. *Legume Research*, 36(2):89-97.
- Agbani, P.O., Kafoutchoni, K.M., Salako, K.V., Gbedomon, R.C., Kegbe, A.M., Karen, H and Sinsin, B. (2018). Traditional ecological knowledge-based assessment of threatened woody species and their potential substitutes in the Atakora mountain chain, a threatened hotspot of biodiversity in Northwestern Benin, West Africa. *Journal of ethnobiology and ethnomedicine*, 14(21), doi:10.1186/s13002-018-0219-6.
- Ahmad, A., Wang, J.D., Pan, Y.B., Sharif, R and Gao, S.J (2018). Development and use of simple sequence repeats (SSRs). Markers for sugarcane. *Breeding and Genetic studies. Agronomy*, 8(11):260.doi:10.3390/agronomy8110260
- Akin-Idowu, P.E., Aduloju, A.O., Akinyoola, O.I., Ibitoye, D.O., Orkpeh, U., Adebo, U.G., Yemisi, O and Olagunju, Y.O (2021). Biodiversity assessment of African locust bean (*Parkia biglobosa*) accessions from Savanna and Forest zones of Nigeria as revealed by seed storage proteins and RAPD markers. *Genetic Resources* 2 (3):36–50.
- Akkinagbe, A., Gailing, O and Lawal, A (2019). Towards conservation of genetic Variation of tropical tree species with differing succession status. The case of *Mansonia altissimo* and *Triplichiton sclerexylon* K. Schum. *Tropical Conservation Science*, 12:1-9.
- Alapa, D.S (2019). Assessment of Environmental Integration Approach in Social Infrastructure Distribution in Otukpo Local Government Area, Benue State, Nigeria. Unpublished M.Sc Dissertation, Department of Geography, Faculty of Environmental Sciences, Nasarawa State University, Keffi.
- Amusa, O., Adesoye, A., Ogunkanmi, O., Omoche, O., Olowe, S. and Akinyosoye, T. O (2014). Genetic Diversity of *Parkia biglobosa* from different agroecological zones of Nigeria using RAPD markers. *International Journal of Biodiversity*, 6pp. <https://doi.org/10.1155/2014/457309>.
- Attikora, A.J.P., Yao, S.D.M., Dago, D.N., Silué, S., De- Clerck, C., Kwibuka Y., Diarrassouba N, Alabi T., Achigan- Dako, E.G and Lassois, L (2024).. Genetic diversity and population structure of superior shea trees (*Vitellaria paradoxa* subsp. *paradoxa*) using SNP markers for the establishment of a core collection in Côte d'Ivoire. *BMC Plant Biol* 24, (913). <https://doi.org/10.1186/s12870-024-05617-0>.
- Boboye, O.M, Oyerinde, O.V and Lawal, A (2021). Genetic diversity of *Adansonia digitata* (L.) and *Vitellaria paradoxa* (c.f. Gaertn) in Nigeria. *Journal of Research in Forestry, Wildlife and Environment*, 13(1):108-122.
- Boboye, O.M., Lawal., A and Oyerinde, O (2018). Assessment of the Genetic Variation in *Chrysophyllum albidum* G. Don Using CP SSR Marker. *Applied Tropical Agriculture*, 23(1):35-40.
- Cui, D., Tang, C., Lu, H., Li, J., Ma, X., Xinxiang, A., Han, B., Yang, Y., Dong, C., Zhang, F., Aai, L and Han, L(2021). Genetic differentiation and restricted gene flow in rice landraces from Yunnan, China: effects of isolation-by-distance and isolation-by-environment. *Rice* 14(1):54. <https://doi.org/10.1186/s12284-021-00497-6>.
- Dardengo, J. de F. E., Rossi, A. A. B. and Varella, T. L. (2018). The effects of fragmentation on the genetic structure of *Theobroma speciosum*

- (Malvaceae) populations in Mato Grosso, Brazil. *Revista De Biología Tropical*, 66(1):218–226.
- de Lafontaine, G., Ducouso, A., Lefevre, S., Magnanou, E and Petit, R. J (2013). Stronger spatial genetic structure in recolonized areas than in refugia in the European beech, *Molecular Ecology*, 22:4397-4412.
- Dedehou, V.F.G.N., Olounladé, P.A., Alowanou, G.G., Azando, E.V.B and Hounzangbé-Adoté, S (2016). A review on medicinal plants of *Parkia biglobosa* (Mimosaceae-Fabaceae) and *Pterocarpus erinaceus* (Leguminosae-Papilionoidea). *JMPS* 4(6): 132-137.
- Denis, M.M (2025). Assessment of the genetic diversity of the traditional tree species *Kigelia africana* (Sausage tree) Using molecular markers for conservation Genomics in Kenya. *European Journal of Medicines plants*, 36(1):45-58.
- Fatoye, F, B (2018). Geology and Mineral resources of Kogi State, Nigeria. *International Journal Multidisciplinary Sciences and Engineering*, 9(7):7-13.
- Frankham, R (2022). Evaluation of proposed genetic goals and Targets for the Conservation on Biological Diversity. *Conservation Genetics*, 23:865-870
- Hartl, D.L. and Clark, A.G. (2007). Principles of Population Genetics. 4th Edition Sinauer Associates, Sunderland, 382, <https://doi.org/10.1093/jhered/esm035>
- Houndonougbo, J. S. H., Kassa, B., Mensah, S., Salako, V. K., Kaka'i, R. G., and Assogbadjo, A. E. (2020). A global systematic review on conservation and domestication of *Parkia biglobosa* (Jacq.) R. Br. ex G. Don, an indigenous fruit tree species in Sub-Sahara African traditional parklands: current knowledge and future directions. *Genetic Resources and Crop Evolution* 67(4):1051–1066.
- Jin, Y, Bian, L., Zhang, X., Zhao, B., Cheng, R., Su, Safe, D., Zheng, X., El-Kassaby, Y and Shi, J(2023). Genetic diversity and structure of the 4th cycle breeding population of Chinese fir (*Cunninghamia lanceolata* (Lamb.) hook). *Plant Sci.*, 14:https://doi.org/10.3389/fpls.2023.1106615.
- Kalinowski, S.T.(2005).HP-RARE 1.0: A computer program for performing rarefaction on measures of allelic richness. *Molecular Ecology Notes*. 5(1): 187-189.
- Katumo, D.M., Liang, H., Ochola, A.C., Lv. M., Wang, O.F and Yan, C.F (2022). Pollinator diversity benefits natural and agricultural ecosystems, environmental health and human welfare. *Plant Diversity*, 44(8):429-435.
- Koura, K., Antoine, A.M., Paulin, S., Hubert A.-S., Clément A., Corneille, A. and Jean C.G.(2019). Transferability of Some Nuclear Microsatellite (SSRs) Markers from Related Species in *Parkia biglobosa*, *Journal of Plant Sciences*. 7(5):125-130.
- Lamien, N., Ekue, M., Ouedraogo, M., Loo, J (2011). *Parkia biglobosa*, Africa locust bean. In Conservation and sustainable use of genetic resources of priority food tree species in Sub-saharan Africa. Rome, Italy: *Biovers. Int.* Leaflet pp1-8.
- Lawal, A., Adekunle, V. A. J. and Oghenekome, U. O. (2023): Molecular Characterization of the Genus *Khaya* in South West Nigeria using CPSSR Markers. *Nigerian Journal of Genetics*, 37(1): 87-98.
- Luettmann, K., Michalczyk, L.M., Mengel. C., Ziegenbagen, B., Heymann, E.W., Saboya, P.P.P and Bialozyt, R (2010). Characterization of nuclear microsatellites loci in the neotropical tree *Parkia panurensis*(Fabaceae). *American Journal of Botany*, 97(5):34-36.
- Lompo, D., Vinceti, B., Konrad, H., Gaisberger, H and Geburek, T (2018). Phylogeography of african locust bean (*Parkia biglobosa*) reveals genetic divergence and spatially structured populations in West and Central Africa. *Journal of Heredity*, 109: 811–814.
- Lompo, D.; Konrad, H.; Duminiel, J.; Gaisberger, H.; Vinceti, B.; Ouedraogo, M.; Geburek, T. (2016). Genetic diversity of *Parkia biglobosa* (African locust bean) and its implications for conservation strategies. Abstract] presented at: Tropentag 2016: Solidarity in a competing world fair use of resources. *Vienna (Austria)*, Sep 18-21. 1 p.
- Mohammed, H.I., Mohammed, Z., Warra , A., Abdulrahman, Y., Sabo, I., Ibrahim, G., U Jajere, U., Victor A and Oyi, R (2022). Physicochemical and Genetic Diversity Studies of *Vitellaria paradoxa* in Northern Nigeria. *Journal of Current Biomedical Research*, 2(1):19-37.
- Nyadanu, D., Adu Amoah, R., Obeng, B., Kwarteng, A.O., Akromah, R., Aboagye, L.M and Adu-Dapaah, H (2017). Ethnobotany and analysis of Food components of African locust bean (*Parkia biglobosa* (Jacq.) Benth) in the transitional zone of Ghana: Implications for Domestication Conservation and Breeding of

- improved varieties. *Genetic Resources Crop Evolution*, 64(6):1231-1240
- Odeunmi, A. A., Salami, A. T. and Ogunwande, I. A. (2017). Impact of anthropogenic activities on the regeneration of *Parkia biglobosa* in derived savanna ecosystems. *Ecology and Conservation Journal*, 5(2):100–108.
- Odekunle, T.O. (2004): Rainfall and length of the growing season in Nigeria, *International Journal of Climatology*, 24: 467-479.
- Olamidayo, F Lawal, A, and Adekunle, V.(2021) Impact of Anthropogenic Activities on Genetic Diversity of *Celtis zenkeri* Engl in South-West Nigeria. *International Journal of Genetics and Genomics*. 9(2):31-35.
- Oluwajuwon T.V., Attafuah, R., Offiah, O.J and Krabel, D (2022).Genetic variation in shea tree (*Vitellaria paradoxa*) in the savanna regions of Ghana. *Genetic Resources and Crop Evolution*, 54(6):1253-126.
- Onyekwelu, J.C., Olusola, J.A., Stimm, B., Mosandi, R and Agbelade, A.D(2015). Farm-level tree growth characteristics, fruit phenotypic variation and market potential assessment of three socio-economically important forest fruit tree species, *Forests, Trees and Livelihoods*, 24(1):27-42.
- Ouedraogo, A. S. (1995). *Parkia biglobosa* (Leguminosae) en Afrique de l'Ouest: Biosystématique et Amélioration. Unpublished Ph.D. dissertation, Wageningen University, Institute for Forestry and Nature Research, Wageningen, Netherlands.
- Ouedraogo, M. (2015). Improving and conserving Sahelian fruit trees: a case study of *Parkia biglobosa*(Jacq.) Benth. (Ph.D. IGN Ph.D Department of Geosciences and Natural Resource Management, Faculty of Science, University of Copenhagen, Frederiksberg. 107p .
- Padakale, E., Atakpama, W., Dourma, K.D., Wala, K., Guelly, K.A and Akpagana, K (2015). Woody Species Diversity and Structure of *Parkia biglobosa* Jacq. Dong Parklands in the Sudanian Zone of Togo(West Africa). *Annual Research and Review in Biology*, 6(2):103-114.
- Pauls, S. U., Nowak, C., Bálint, M. and Pfenninger, M. (2013). The impact of global climate change on genetic diversity within populations and species. *Mol. Ecol.* 22: 925–946.
- Phurailatpam, C.S., Lyngdoh, N., Jaganathan, M., Thunga, P.K., Bhat, P and Ravikanth, G (2022). Microsatellite Analysis of Genetic Diversity and Structure of Farmer Selected Genotypes of Tree Bean (*Parkia timoriana* (DC.) Merr) in Manipur. *Indian Journal of Ecology*, 49(1):31-37.
- Popoola, J.O., Agbola, J.O., Ajiboye, A., Akinola, O and Lewu, F (2020). Assessment of Genetic diversity of Afriocan Locust Bean (*Parkia biglobosa* Jacq) landraces usin g microsatellites. *Journal of Critical Reviews*, 7(11):3196-3205.
- Ræbild, A., Larsen, A.S., Jensen, J.S., Ouedraogo, M Groote, S.D Damme, P.V., Bayala, J., Diallo, B.O., Sanou, H., Kalinganire, A., Kjaer, E.D. (2021). Advances in domestication of indigenous fruit trees in the West African Sahel. *New Forests*. 41(3):297-315.
- Sacande, M., Jøker, D., Dulloo, M. E., and Thomsen, K. A. (2007). Comparative seed morphology and germination of *Parkia biglobosa* (Fabaceae) from different provenances in West Africa. *Seed Science and Technology*, 35, 40–56.
- Salgotra R.K and Chauhan, B.S (2023). Genetic Diversity, Conservation, and Utilization of Plant Genetic Resources *Genes*, 14(1):174 DOI: 10.3390/genes14010174.
- Sani, M. H. and Haruna, U. (2010): Planning model for sustainable vegetable crop production in the eastern part of Kogi State, Nigeria. *Journal of Agronomy*, 9 (1):17-22.
- Sexton, G.J., Frere, C.H., Kalinganire, A., Uwamariya, A, Lowe, A.J, Godwin, I. D., Prentis P.J and Dieters, M.J. (2015). Influence of putative forest refugia and biogeographic barriers on the level and distribution of genetic variation in an African savannah tree, *Khaya senegalensis* (Desr.) A. Juss. *Tree Genet Genomes*, 11:103–103.
- Sina, S (2006). Reproduction et Diversité Génétique chez *Parkia biglobosa* (Jacq.) G. Don. PhD thesis Wageningen University, the Netherlands, 102 p.
- Soares, S. D., Bandeira, L.F., Ribeiro, S.B., de Campos Telles, M.P., da Silva, J.A., Borges, C.T., Coelho, A.S.G and Novaes, E (2020). Genetic diversity in populations of African mahogany (*Khaya grandiofiola* C. DC.) introduced in Brazil. *Genet. Mol. Biol*, 43(2): e20180162. doi: 10.1590/1678-4685-GMB-2018-0162.
- Szasz-Len, A..M. and Konnert, M (2018). Genetic diversity in European beech (*Fagus Sylvatica* L.) Seed Stands in the Romanian Carpathians. *Annals of Forest Research*, 6(1): DOI: <https://doi.org/10.15287/afr.2018.1019>.
- Teklehaimanot, Z. (2004). Exploiting the potential of indigenous agroforestry trees: *Parkia*

- biglobosa* and *Vitellaria paradoxa* in sub-Saharan Africa. *Agroforestry Systems*, 61–62, 207–220.
- Temunovic, M., Frascaria-Lacoste, N., Franjic, J., Satovic, A.Z. and Fernandez-Manjarres, J.F. (2013). Identifying refugia from climate change using coupled ecological and genetic data in a transitional Mediterranean-temperate tree species. *Molecular Ecology*, 22(8):2128-2142.
- Tsipidou, O., Leinemann, L., Korakis, G., Finkeldey, R. G, Gailing, O and Papageorgiou, A.C (2022). Fine scale spatial patterns of Genetic diversity of European Beech (*Fagus sylvatica* L) around a Mountainous Glacial Refugium in the SW Balkans. *Forest*, 12(6):725 10.3390//2060725.
- Wang, X., Wang, L., Sun, Y., Chen, J., Liu, Q and Dong, S (2023). Genetic diversity and conservation of Siberian apricot(*Prunus sibirica*, L) based on microsatellites, markers. *Scientific Reports*, 13(1):11245. doi: 10.1038/s41598-023-37993-2.

10/19/2025