

## The Effect Of Anuloma Viloma Pranayama And Kapalbhathi On Resting Pulse Rate And Stress Of School Going Children In Bhubaneswar

Manoranjan Tripathy

Research Scholar, Department of Clinical Psychology, Dev Sanskriti University, Haridwar, India  
[manoranjanonly@gmail.com](mailto:manoranjanonly@gmail.com)

**Abstract:** To determine the effect of Anuloma-viloma Pranayama and Kapalbhathi on resting pulse rate and stress, one hundred fifty (N=150) randomly selected male student aged 15-17 years to participated in the study of DAV Public School, Bhubaneswar, Odisha, India. They were randomly assigned into three groups: A- Pranayama, B- Kapalbhathi and C- Control. The subjects were subjected to the twelve weeks Pranayama and Kapalbhathi training program. The between group differences was assessed using the post-hoc test. The level of  $p \leq 0.05$  was considered significant. The pulse rate and stress significant decrease in group A and B compared with the control group-c.

[Manoranjan Tripathy. **The Effect Of Anuloma Viloma Pranayama And Kapalbhathi On Resting Pulse Rate And Stress Of School Going Children In Bhubaneswar.** *N Y Sci J* 2019;12(1):74-78]. ISSN 1554-0200 (print); ISSN 2375-723X (online). <http://www.sciencepub.net/newyork>. doi: [10.7537/marsnys120219.09](https://doi.org/10.7537/marsnys120219.09).

**Key Words:** Anuloma-viloma Pranayama, Kapalbhathi, Pulse Rate Stress

### Introduction

In this contemporary fast growing world of science and technology, the human elements are treated as ever before. Its goals are distant and dissatisfying<sup>1</sup>. The mechanism of modern living, the force restriction of physical activity leading to a century life, an increased amount of leisure, time, all these factors have resulted in a tremendous increase of professional and public interest in physical activity and health<sup>2</sup>. People seem to be “turning on” to the idea that one looks and feels better and stays healthier by being physically more active. Today yoga, being a matter of varied interests, has gained worldwide popularity<sup>3</sup>. Recent research trends have shown that it can serve as an applied science in a number of fields such as physical education and sports, (Health and family welfare, psychology and medicine and one of the valuable means for the development of human resources for better performance and productivity). However, it has generally been believed that yoga is a spiritual science having emancipation as its goals and hence cannot be treated only as a therapy<sup>4</sup>. It is now being achieved in all parts of the globe that yoga is not only for better development of mind, socio-control (spiritual or moral) but a therapy. Yoga focuses on harmony between mind and body. Yoga derives its philosophy from Indian metaphysical beliefs<sup>5</sup>. The word yoga comes from the Sanskrit language and means union or merger. The ultimate aim of this philosophy is identical to strike a balance between body, mind and attain self-enlightenment. To achieve this, yoga uses movement, breath, relaxation, posture and meditation in order to create a lively, healthy and balanced approach to life<sup>6</sup>.

### Objective of the Study

1. To find out the effect of Kapalbhathi on resting pulse rate of school going children of Bhubaneswar.
2. To find out the effect of Anuloma-viloma Pranayama on resting pulse rate of school going children of Bhubaneswar.
3. To find out the effect of Kapalbhathi on stress of school going children of Bhubaneswar.
4. To find out the effect of Anuloma-viloma Pranayama on stress of school going children of Bhubaneswar.

### Methodology

#### A. Method

The researcher was followed the experimental research method in the study. There were three groups, one act as a Kapalbhathi group and second act as a Pranayama group and no act control group.

- Group “A” - Kapalpalbhathi
- Group “B” - Anuloma-Viloma Pranayama
- Group “C” - Control

The duration of experimental period was twelve weeks.

The pre-test conduct before the practice.

The post-test conduct after the practice.

#### B. Size of Sample

The size of the sample was 150 school boys.

- 75 students were selected from DAV Public School, Bhubaneswar, Odisha, India to Class 11 and 12 those who are interested in yoga.

- 75 students were selected from DAV Public School, Bhubaneswar, Odisha, India to Class 11 and 12 those who are interested in yoga.

**Variables**

- Resting Pulse Rate
- Stress

**Resting Pulse Rate**

Resting Pulse Rate was recorded as the number of Pulse per minute during the state of rest.

**Table 1: Descriptive statistics of the data measured in the post testing pulse rate**

Groups	Practices	N	Mean	Std. Deviation
Group - A	Kapalbhati	50	68.44	6.73
Group – B	Anuloma-Viloma Pranayama	50	71.38	5.72
Group -C	Control	50	74.20	5.13
Total		150	71.34	6.31

Table 1 indicates the values of descriptive statistics of the experimental Groups Kapalbhati Group, Anuloma-Viloma Pranayama Group & Control Group for physiological variable of pulse rate, which shows that the mean and S.D. values of Pranayama

Group, Kapalbhati Group and the Control Group were found to be 68.44±6.73, 71.38±5.72 and 74.20±5.13 respectively. Total the same was 71.34±6.31.

**Table 2: Descriptive statistics of the data measured in the post-testing after adjustment with the initial difference pulse rate**

Groups	Practices	Mean	Std. Error	95% Confidence Interval	
				Lower Bound	Upper Bound
Group - A	Kapalbhati	69.263 <sup>a</sup>	.313	68.643	69.882
Group – B	Anuloma-Viloma Pranayama	71.646 <sup>a</sup>	.312	71.029	72.263
Group -C	Control	73.111 <sup>a</sup>	.314	72.490	73.732

<sup>a</sup> Covariates appearing in the model were evaluated at the following values: pre pulse rate = 72.67

The mean and standard error of different post-testing Groups after adjustment have been shown in

Table 2. Which were for Kapalbhati Group 69.263, Pranayama Group 71.646 & 0.312 & 0.313 and Control Group 73.111 & 0.314.

**Table 3: ANCOVA table for the post-test data on pulse rate**

Source	Sum of Squares	df	Mean Square	F Sig.	(p-value)
Pre-Pulse rate	4401.134	1	4401.134	903.792	.000
Treatment Group	370.271	2	185.135	38.018	.000
Error	710.966	146	4.870		
Corrected Total	5482.371	149			

Table 3 indicates the values test of difference between the subject effects, which shows that there was a significant difference in pre-test values of physiological variable of pulse rate for the three selected Groups, as the value was found to be 903.792,

which proves to be the base of Analysis of Co-Variance. Also, a significant difference was found between the post-test values of the experimental and Control Group as the value was found to be 38.018, which was significant at 0.05 level.

**Table 4: Post hoc comparison for the group means in post-measurement adjusted with the initial differences pulse rate**

(I) Different Groups	(J) Different Groups	Mean Difference (I-J)	SIG. a (p-value)
Pranayama	Kapalbhati	2.384*	.000
	Control	-1.465*	.001
Kapalbhati	Pranayama	-2.384*	.000
	Control	-3.848*	.000
Control	Pranayama	1.465*	.001
	Kapalbhati	3.848*	.000

Based on estimated marginal means:

a. Adjustment for multiple comparisons: Least Significant Difference (equivalent to no adjustments).


\* The mean difference is significant at the 0.05 level.

Table 4 indicates the values of post hoc test for the selected Groups for physiological variable of pulse rate, which shows that a significant difference was

found between the post-test values of Pranayama Group and the Kapalbhathi Group as the value was found to be 2.384 which was significant at 0.05 level, the post-test values of Pranayama Group and the Control Group as the value was found to be 1.465 which was significant at 0.05 level, Kapalbhathi Group and the Control Group as the value was found to be 3.848 which was significant at 0.05 level.

**Table 5: Comparison among three groups, pre and post-test mean of pulse rate**

Group	Subject	Mean (Pre-Test)	Mean (Post-Test)
Pranayama	50	72.36	71.38
Kapalbhathi	50	71.72	68.44
Control	50	73.92	74.20



**Figure 1: Bar diagram showing the mean value of pulse rate among Pranayama group, Kapalbhathi group and Control group**

**Table 6: Descriptive statistics of the data measured in the post testing stress**

Different Groups	N	Mean	Std. Deviation
Pranayama	50	12.280	3.923
Kapalbhathi	50	13.980	3.772
Control	50	17.320	4.488
Total	150	14.527	4.558

Table 6 indicates the values of descriptive statistics of the experimental Groups (Pranayama Group, Kapalbhathi Group) & Control Group for psychological variable of stress, which shows that the

mean and S.D. values of Pranayama Group, Kapalbhathi Group and the Control Group were found to be 12.280±3.923, 13.980±3.772, and 17.320±4.488 respectively. Total the same was 14.527±4.558.

**Table 7: Descriptive statistics of the data measured in the post-testing after adjustment with the initial difference stress**

Groups	Practices	Mean	Std. Error	95% Confidence Interval	
				Lower Bound	Upper Bound
Group - A	Kapalbhathi	14.425 <sup>a</sup>	.228	13.973	14.876
Group - B	Anuloma-Viloma Pranayama	12.835 <sup>a</sup>	.229	12.382	13.287
Group -C	Control	16.321 <sup>a</sup>	.231	15.865	16.777

<sup>a</sup> Covariates appearing in the model have evaluated at the following values: pre stress = 15.6867.

The mean and standard error of different post-testing Groups after adjustment have been shown in

table 6. Which are for Kapalbhathi Group 14.425, Pranayama Group 12.835 & 0.229 & 0.228 and Control Group 16.321 & 0.231.

**Table 8: ANCOVA table for the post-test data of stress**

Source	Sum of Squares	df	Mean Square	F Sig.	(p-value)
Pre-Stress	2058.638	1	2058.638	792.406	.000
Treatment Group	295.559	2	147.779	56.883	.000
Error	379.302	146	2.598		
Corrected Total	3095.393	149			

Table 8 indicates the values test of difference between the subject effects, which shows that there were a significant difference in pre-test values of psychological variable of stress for the three selected Groups, as the value has found to be 792.406, which

proves to be the base of Analysis of Co-Variance. Also, a significant difference is found between the post-test values of the experimental and Control Group as the value has found to be 56.883, which was significant at 0.05 level.

**Table 9: Post hoc comparison for the group means in post-measurement adjusted with the initial differences stress**

(I) Different Groups	(J) Different Groups	Mean Difference (I-J)	SIG. a (p-value)
Pranayama	Kapalbhathi	-1.590*	.000
	Control	-3.486*	.000
Kapalbhathi	Pranayama	1.590*	.000
	Control	-1.896*	.000
Control	Pranayama	3.486*	.000
	Kapalbhathi	1.896*	.000


Based on estimated marginal means

a. Adjustment for multiple comparisons: Least Significant Difference (equivalent to no adjustments).

\* The mean difference is significant at the 0.05 level.

Table 9 indicates the values of post hoc test for the selected Groups for psychological variable of stress, which shows that a significant difference has

found between the posttest values of Pranayama Group and the Kapalbhathi Group as the value has found to be 1.590\* which was significant at 0.05 level, the post-test values of Pranayama Group and the Control Group as the value has found to be 3.486\* which was significant at 0.05 level, Kapalbhathi Group and the Control Group as the value has found to be 1.896\* which was significant at 0.05 level.



**Figure 2: Bar diagram showing the mean value of stress among Pranayama group, Kapalbhathi group and Control group**

**Table 10. Comparison among three groups, pre and post-test mean of Stress**

Group	Subject	Mean (Pre-Test)	Mean (Post-Test)
Pranayama	50	15.65	12.20
Kapalbhati	50	15.70	14.01
Control	50	16.30	16.70

### Discussion and Findings

We observed no significant difference in the baseline parameters of the three groups and therefore, they can be examined comparable for the study. After 12 weeks of the study period, researcher observed a significant decrease in resting pulse rate in both (Anuloma-viloma Pranayama and Kapalbhathi) groups with no change in control group (Group C). But the Kapalbhathi group is more effective then comparison of Anuloma-viloma Pranayama group.

The researcher observed a significant decrease in stress in both (Anuloma-viloma Pranayama and Kapalbhathi) groups with no change in control group (Group C). But the Anuloma-viloma Pranayama group is more effective than comparison of Kapalbhathi group.

To summarize, our study demonstrates that Anuloma-viloma Pranayama and Kapalbhathi were effective in reducing perceived resting pulse rate and stress. Kapalbhathi was more suitable for subjects with stable resting pulse rate and Pranayama was more suitable for subjects with stable stress. On post-test analysis, sample size (150 students) was found adequate for the present study.

### References

1. Deol Nishan Singh, *Futuristic Trends in Physical Education*, p.147 (2012).

2. Shirotriya, A. K., and B. Basumatary. "Occupational Stress & Frustration Tolerance of Indian Security Forces." *Indian Journal of Psychometry and Education* 41, no. 2 (2010): 214-218.
3. Maurya, Shailesh. "Effect of Asana and Pranayama on Flexibility of University Students." (2011).
4. Tomar, Rakesh, and Neelima Singh. "Effect of Ujjayi Pranayama on Selected Physiological Variables." *Academic Journal Ovidius University Annals, Series Physical Education & Sport/Sci* 11, no. 1 (2011): p140.
5. Raja, Chidambara. "Effect of Yogic Practices on Flexibility Cholesterol and Blood Pressure." *Online International Interdisciplinary Research Journal* 2, no. 6 (2012): 221-225.
6. Upadhyay, D. K., "Effects or Alternate Breathing Exercise on Cardio-Respiratory Functions." *Nepal Med Journal*, 2008, 10:1. PP. 25.
7. Sharma, Vivek Kumar, Madanmohan Trakroo, Velkumary Subramaniam, M. Rajajeyakumar, Anand B. Bhavanani, and Ajit Sahai. "Effect of fast and slow pranayama on perceived stress and cardiovascular parameters in young healthcare students." *International journal of yoga* 6, no. 2 (2013): 104.

2/25/2019