



The Efficacy of Methanol, Dichloromethane and N-Butanol Extracts of *Annona muricata* Leaves on Selected Bacteria And Fungi

Akintelu Sunday Adewale¹ And Folorunso Aderonke Similoluwa^{2*}

¹Department of Pure and Applied Chemistry, Ladoke Akintola University of Technology, PMB 4000 Ogbomosho, Nigeria.

²Department of Chemistry, Louisiana State University, Louisiana, USA
folorunsoaderonkesimi@gmail.com

Abstract: In Nigeria, *Annona muricata* is locally used as source of medicine for curing various infections. This research was aimed at determination of the potency of the leaves against selected bacterial and fungal strains. The antibacterial and antifungal activities of methanol, dichloromethane and n-butanol extracts of *Annona muricata* leaves against selected bacterial and fungi strains were examined via standard methods. Methanol, dichloromethane and n-butanol extracts of *Annona muricata* leaves showed high inhibitory activities between (24-29 mm) against *Bacillus subtilis* (25-29mm) against *Clostridium sporogenes*, (25-29) against *Enterococcus faecalis* (30-31) against *Klebsiella pneumonia* (26-31) against and (26-30) against *Staphylococcus aureus*. The antifungal activities of methanol, dichloromethane and n-butanol extracts of *Annona muricata* leaves showed broad inhibition zones against the growth of *Aspergillus flavus*, *Candida albican*, *Fusarium oxysperium* and *Penicillium camemeri*. This study provide scientific information and justification that support the local use *Annona muricata* leaves as medicinal therapy and equally revealed *Annona muricata* leaves as a major ingredient to bank on for the design of novel antibiotics.

[Akintelu Sunday Adewale And Folorunso Aderonke Similoluwa. **The Efficacy of Methanol, Dichloromethane and N-Butanol Extracts of *Annona muricata* Leaves on Selected Bacteria And Fungi.** *N Y Sci J* 2019;12(10):53-56].
ISSN 1554-0200 (print); ISSN 2375-723X (online). <http://www.sciencepub.net/newyork>. 6.
doi:[10.7537/marsnys121019.06](https://doi.org/10.7537/marsnys121019.06).

Keywords: Antifungal, antibacterial, efficacy and *annona muricata*

1. Introduction

The search for novel antibiotics with efficiency to terminate the growth of microbial infections across the world has led to the increase in the use of plants with good medicinal properties as important treatment alternatives (Abba, et al 2009; Akintelu et al 2018; Akintelu et al 2019). The examination of the antimicrobial efficiency of many plant extract has been embarked on recently to discern new antimicrobial agents with less side effect on human and better efficiency against microbes (Sarita, et al 2019). *Annona muricata* is a plant from Annonaceae family whose medicinal value is commended in Nigeria and many Africa countries. The pharmacological investigations on the extracts of *Annona muricata* plant shows its potency and efficiency to cure ulcer, inflammatory and bacterial (Tai et al 2014; Folorunso et al 2019b). Information from researchers also stated that *Annona muricata* fruit has been used for the treatment of cough, skin rashes and parasites infections (Haidan et al 2016).

Traditional healers claim that herbs made from medicinal plants are more potent and possess less side harm when compared with conventional medicines. In undeveloped nations, people with low revenue from the remote societies use herbs for the cure of disease causes

by microbial infections (Haidan et al 2016). Therefore, the use of herbal drugs could serve as an alternate in treatment of sicknesses instigated by multidrug resistant bacteria antifungal (Ramesh et al 2017).

Many antibiotics have lost their usage in combating microbes due to multidrug-resistant strains and other harms they produce when used (Peter et al 2018; Akintelu and Folorunso 2019). Literatures have shown the microbial resistance to antibiotics such as tetracycline, erythromycin and penicillin (Timothy 2017). Also the resistance developed by some human pathogens against conventional antibiotics that are readily available in the market and pharmaceutical stores calls for adoption of other alternatives from other sources including medicinal plants. This fact then trigger us to embark on the study of the efficacy of *Annona muricata* leaves extract on some microorganism.

Materials and methods

Analytical grade reagents include n-hexane, methanol, dichloromethane, n-butanol, aluminum oxide, silica gel, diethyl ether and methanol and apparatus use are autoclave, incubators, pipette, petri-dish canister, inoculating loop, spatula,

drying oven and weighing balance.

Sample collection and Extraction process

Annona muricata leaves were obtained from Okitipupa town in Ondo state. The leaves were authenticated by a botanist from Obafemi Awolowo University, Ife, Nigeria. The leaves were properly rinsed to remove dust and other impurities and air dried. The dried leaves were then pulverized. About 2 kg of powdered *Annona muricata* leave was extracted with 10 L of methanol for 48 hours with adequate agitation at the intervals of 3 hours. The extracting solvent was decanted and filtered after 48 hours of extraction, the extract was partitioned with methanol, dichloromethane, and n- butanol. Each of the fractions was concentrated on a digital rotary evaporator (Heidolphlaborata 4010) to obtain various extracts of the *Annona muricata* used.

Table 1.0 Microorganism used

Bacteria	Fungi
<i>Bacillus subtilis</i>	<i>Aspergillus flaws</i>
<i>Clostridium sporogenes</i>	<i>Candida albican</i>
<i>Entrococcusfaecalis</i>	<i>Fusariumoxysperium</i>
<i>Klebsiella pneumonia</i>	<i>Penicilliumcamemeri</i>
<i>Pseudomonas aeruginosa</i>	
<i>Staphyloccusaureus</i>	

Determination of the antifungal activities of the extracts of the *annona muricata* leaves.

Clinical isolated fungi obtained were grown on Sabouraud dextrose agar at 25 °C. They were monitored until they sporulated. The fungi spores were gathered

into the broth and properly standardized. 20 mL of Sabouraud dextrose agar was seeded with 200 µL of the standardized broth into Maccartney bottles and was transferred into a sterile petri dish after swirling. Several wells of 6 mm diameter were made on the agar with sterilized cork borer. 1 mg/L of each of the extract was transferred into a well and another well was filled with 1 mg/L of amphotericin B as a positive control. Proper diffusion of compound into the agar was ensured. The incubation of the plates at 25 °C for 96 hours was done. The zones of inhibition around the wells were determined. This experiment was repeated twice.

Determination of the Antibacterial activities of the extracts of the *Annona muricata* leaves.

The test bacteria strains were inoculated into tubes containing peptone water and incubated for 18 hours at a temperature of 37 °C before usage. Each of the cultures was adjusted to standard of 0.5 Mcfarland turbidity. 20 mL of sterile molten Mueller Hinton agar was seeded with 200 µL of standardized cell suspensions into the Maccartney bottles and was gently transferred into the sterile plates. Wells with 6 mm diameter were made on the plates via cork borer. 0.2 mL of 2mg/L of each extract were transferred into each well. About 1 mg/L of the positive control (Streptomycin) were transferred into the remaining wells. Proper diffusion of the cultured plates was monitored and the plates they were incubated at 35 °C for 24 hours. The zones of inhibition were recorded. This procedure was repeated for two more times.

3. Results

Table 2.0 Result of the antifungal activities of the methanol, dichloromethane and n-butanol extract of *annona muricata* leaves

Tested Fungi	Inhibition zone (mm)			
	methanol 2 mg/L	dichloromethane 2 mg/L	n-butanol 2 mg/L	Amphotercin B 1 mg/L
<i>Aspergillus flaws</i>	27	26	0.0	20
<i>Candida albican</i>	26	0.0	25	22
<i>Fusariumoxysperium</i>	25	0.0	26	21
<i>Penicilliumcamemeri</i>	26	0.0	31	23

Table 3.0 Result of the antibacterial activities of the methanol, dichloromethane and n-butanol extract of *annona muricata* leaves

Bacteria isolates	Inhibition zone (mm)			
	methanol 2 mg/L	dichloromethane 2 mg/L	n-butanol 2 mg/L	Steptomycin 1 mg/L
<i>Bacillus subtilis</i>	27	26	25	27
<i>Clostridium sporogenes</i>	29	26	25	24
<i>Entrococcusfaecalis</i>	27	28	25	26
<i>Klebsiella pneumonia</i>	31	0.0	30	27
<i>Pseudomonas aeruginosa</i>	26	28	31	25
<i>Staphyloccusaureus</i>	26	29	30	24

4. Discussions

The result obtained from the antifungal activities of the methanol, dichloromethane and n-butanol extract of *annona muricata* leaves on the selected fungi shows that the methanol extract has highest Inhibition zone against *Aspergillus flaws* (27mm) and lowest Inhibition zone against *Fusarium oxysperium* (25mm), The dichloromethane only has an inhibition zone against *Aspergillus flaws* (27mm) while the n-butanol extract has highest inhibition zone against *Penicillium camemeri* (31mm) and lowest inhibition zone against *Candida albican* (25mm). The extracts had no inhibition against the growth of *Candida albican*, *Fusarium oxysperium*, *Penicillium camemeri* in dichloro methane extract and *Aspergillus flaws* in the n-butanol fraction as shown in table 2.

The methanol, dichloromethane and n-butanol extracts showed higher activity when compared with the control (amphotericin B) judging from their zones of inhibition. The antifungal investigation of *Annona muricata* leaves showed better antifungal potency when compared with *Garcinia kola* seed from the study of (Folorunso et al 2019a)

This findings support the usefulness of *Annona muricata* leaves in folklore remedies for the treatment of diseases caused by these pathogen.

Antimicrobial analysis (Table 3.0) showed that the methanol, dichloromethane and n-butanol extract of *Annona muricata* leave were active against all the clinical isolated bacterial strains except in dichloromethane extract against *Klebsiella pneumonia* where no inhibition zone was observed. The extracts showed inhibition zones comparable to streptomycin used as control in this study. The antibacterial activity of this study was found higher than that obtained from the study of (Yohannes et al 2019) and were in close range with the antibacterial activity obtained from the study of (Oyebamiji et al 2019). The bioactive compounds present in this plant may be responsible for the antibacterial activity of these extracts (Avato et al 2006; Mwitari et al 2013; Alfredo et al 2016) Manikkuwadura et al (2019) also reported that the antibacterial properties of plants are due to the presence of phytochemicals such as carotenoids, flavonoids, lycopene, phenolics and β -carotene present in them.

Conclusion

The extract of *Annona muricata* shows a broad spectrum against the selected fungi and bacterial which shows the potency of *Annona muricata* as a remedy for treating diseases cause by pathogens. The antimicrobial property displayed by these extracts validate the need for further study on isolation and characterization of active phytochemicals that can fight microbial resistance exhibited by majority of existing antibiotics

from the leaves of *Annona muricata*.

Corresponding Author:

Miss Folorunso Aderonke Similoluwa
Department of Chemistry, Louisiana State University,
Louisiana, USA
Email: folorunsoaderonkesimi@gmail.com

Reference

1. Abba, D., Inabo, H. I., Yakubu, S. E., and Olonitola, O. S. (2009). Contamination of herbal medicinal products marketed in Kaduna Metropolis with selected pathogenic bacteria. *African Journal of Traditional, Complementary and Alternative Medicines*, 6, 70-77.
2. Akintelu S.A, Abiola B.E, Ajayi S.O, Olabemiwo O.M (2019) Preliminary Study of the Effects of some Herbal Drugs on the Heamatological and Biochemical Parameter of Winster Albino Rats. *International Journal of Sciences*. 8(10) 26-31 DOI: 10.18483/ijSci.2032;
3. Sarita, M., Shisir, L., and Raj, K.D (2019). In Vitro Antimicrobial Activity of Some Medicinal Plants against Human Pathogenic Bacteria. *Journal of Tropical Medicine*.1-5. doi.org/10.1155/2019/1895340.
4. Tai, S. Kedari, and Ayesha A. K (2014) Guyabano (*Annona Muricata*): A review of its Traditional uses Phytochemistry and Pharmacology. *American Journal of Research Communication* 2(10) 247 - 268.
5. Haidan, Y., Qianqian, M., Li, Y., and Guangchun, P., (2016). The Traditional Medicine and Modern Medicine from Natural Products. *Molecules* 21,1-18.
6. Folorunso A.S, Oyebamiji A.K, Erazua E.A and Akintelu S.A (2019a) The Exploration of Antifungal Activity of *Garcinia kola* Seed as Novel Antifungal Agent. *International Journal of Traditional and Natural Medicines*, 9(1): 41-49
7. Ramesh, S., Mathivanan, N., Klaus, D.F (2017) Plant-derived antimicrobials to fight against multi-drug-resistant human pathogens 3 *Biotech* 7:172.
8. Peter M H., Roderic, E W., David, M L., Cliodna, A M M., David, A.E., Jonathan A.O., Peter AR., (2018). Wilson Treatment of infections caused by multidrug-resistant Gram-negative bacteria: report of the British Society for Antimicrobial Chemotherapy/Healthcare Infection Society/British Infection Association Joint Working Party. *Journal of Antimicrobial Chemotherapy*, (73)3; 2–78, <https://doi.org/10.1093/jac/dky027>.

9. Oyebamiji A.K, Akintelu S.A, Folorunso A.S., Abiola B.E., Ajayi S.O., Abdusalam I O., and Morakinyo A E. (2019), Computational and Experimental Studies on Antimicrobial Activity of the Bark of *Annona muricata* against Some Selected Human Pathogenic Bacteria and Fungi International Journal of Modern Chemistry, 11(1): 9-27
10. Timothy, J.F (2017) Antibiotic resistance in *Staphylococcus aureus* Current status and Future prospects. FEMS Microbiology Reviews, (41) 3; 430–449.
11. Yohannes K.E, Ebrahim A.S, Tekleab T.T, and Gedefaw G.A (2019) Antibacterial Potential of *Aloe weloensis* (Aloeacea) Leaf Latex against Gram-Positive and Gram-Negative Bacteria Strains International Journal of Microbiology. 5328238, 1-4
<https://doi.org/10.1155/2019/5328238>.
12. Folorunso A , Akintelu S. ,Oyebamiji A.K, · Ajayi S Babawale Abiola B, Abdusalam I and Morakinyo A.(2019b) Biosynthesis, characterization and antimicrobial activity of gold nanoparticles from leaf extracts of *Annona muricata* Journal of Nanostructure in Chemistry, 9:111–117. doi.org/10.1007/s40097-019-0301-1
13. Avato, P., Bucci, R., Tava, A., Vitali, C., Rosato, A., Bialy, Z. et al., (2006) ‘Antimicrobial activity of saponins from *Medicago* sp.: Structure-activity relationship’, *Phytotherapy Research* 20(6), 454–457.
14. Akintelu S.A, Abiola B.E, Ajayi S.O and Olabemiwo O.M (2018)Quantification and Preliminary Estimation of Toxic Effects of Polycyclic Aromatic Hydrocarbon in Some Antimalarial Herbal Drugs in Southwest Nigeria *Bulletin of Pharmaceutical Research*, 8(1):152. DOI: 10.21276/bpr.2018.8.1.1
15. Mwitari, P.G., Ayeka, P.A., Ondicho, J., Matu, E.N. & Bii, C.C., 2013, ‘Antimicrobial activity and probable mechanisms of action of medicinal plants of Kenya: *Withaniasomnifera*, *Warbugiaugandensis*, *Prunusafricana* and *Plectrunthusbarbatus*’, *PLoS One* 8(6), e65619.
16. Akintelu SA, Folorunso AS (2019) Characterization and Antimicrobial Investigation of Synthesized Silver Nanoparticles from *Annona muricata* Leaf Extracts. J Nanotechnol Nanomed Nanobiotechnol 6:1-5. DOI: 10.24966/NTMB-2044/100022
17. Alfredo, A., Eduardo, M., Rosa C., Carla D and Maria J.S (2016) Phytochemical Composition and Antibacterial Activity of Hydroalcoholic Extracts of *Pterospartum tridentatum* and *Mentha pulegium* against *Staphylococcus aureus* Isolates 1-11.<http://dx.doi.org/10.1155/2016/520187912>.
18. Manikkuwadura, H.N., Hasanga, R., Ruwani, P.H and Weerasinghe, MD G (2019). Determination of In Vitro Antimicrobial Activity of Five Sri Lankan Medicinal Plants against Selected Human Pathogenic Bacteria. International Journal of Microbiology.1-8
<https://doi.org/10.1155/2019/7431439>.

10/17/2019