



Microbiological pollution and heavy metals in two freshwater turtles from Garmat Ali- canal in Basrah City/ Southern Iraq

Ghazi M. Al-Maliki

Department of Marine Biology, Marine Science Center, University of Basrah, Iraq

Corresponding author E mail: ghazialmaliki@yahoo.com

Abstract: The aim from this study for determine the microbiological pollution and some heavy metals (Fe, Cu, Zn and Pb) levels in two freshwater turtles species (*Mauremys c. caspica* (Gmelin 1774), and *Rafetus (Trionyx) euphraticus* (Daudin 1802)) were collected from Garmat Ali- canal. Samples of two freshwater Turtles were collected during summer season of the year 2017, were examined. Total- plate techniques were used for microbial pollution while heavy metal concentrations in two turtles species were determined using atomic absorption spectrophotometer (AAS). Total bacteria count had the highest number of bacteria in *M. c. caspica* with 29.60×10^5 cfu/ml. the total plate count had the highest bacteria number in the *R. euphraticus* – 19.26×10^5 cfu/ml ranged. bacterial pathogens isolated include: *E. coli*, *Pseudomonas spp.*, *Vibrio spp.*, and *Staphylococcus spp.* Microbial species were characterized based on morphological and biochemical tests.

M. c. caspica Turtle had iron concentration of 21.52 ± 0.031 mg/kg while *R. euphraticus* Turtle had iron concentration of 34.69 ± 0.152 mg/kg. *M. c. caspica* and *R. euphraticus* had copper concentrations of 0.87 ± 0.034 and 0.92 ± 0.057 mg/kg respectively. *M. c. caspica* had a lead concentration of 0.06 ± 0.041 mg/kg while *R. euphraticus* had a lead concentration of 0.09 ± 0.064 mg/kg.

[Ghazi M. Al-Maliki. **Microbiological pollution and heavy metals in two freshwater turtles from Garmat Ali- canal in Basrah City/ Southern Iraq.** *N Y Sci J* 2019;12(11):47-51]. ISSN 1554-0200 (print); ISSN 2375-723X (online). <http://www.sciencepub.net/newyork>. 6. doi: [10.7537/marsnys121119.06](https://doi.org/10.7537/marsnys121119.06).

Key words: Microbiological pollution, heavy metals, *E. coli*, bacterial pathogens, *Mauremys c. caspica* turtles.

Introduction

Three species of freshwater turtles and tortoise are recorded from Iraq: the Mediterranean spur-thighed tortoise, *Testudo graeca terrestris* (Forskall 1775), Eastern Caspian turtle, *Mauremys c. caspica* (Gmelin 1774), and Euphrates soft-shelled turtle *Rafetus (Trionyx) euphraticus* (Daudin 1802), (Al-Barwari and Saeed 2007). Both species, *R. euphraticus* and *M. c. caspica*, are probably threatened in Iraq. The turtles of Iraq have received scant scientific attention since Khalaf (1959); however, little is known about their distribution and status (Pritchard 1967, 1979; Ghaffari 2002).

The Garmat Ali River was subjected to multiple impacts from hydrological and human activities. After inundation of the southern marshes in 2003, the East Hammar marsh was fed primarily from the Euphrates River and entering the Garmat Ali River then the Shatt Al-Arab River that eventually flows into the Arabian Gulf. So, this river affected by the water from the Euphrates and tidal current of the Gulf through the Shatt Al-Arab River. But, the flow of the Euphrates was diverted away from the north East Hammar marsh during the last years, consequently the water level in the marsh dropped sharply, causing the water salinity higher than before, which led to negatively affected on

water quality and quantity of the Garmat Ali River (Al-Tememi, et al., 2015). Moreover, the Shatt Al-Arab River suffered from massive regression in water quality related to the decline in rates of discharge from the Tigris and the Euphrates Rivers (Al-Mahmood, et al., 2015) as a result of several hydrological projects constructed in the riparian countries (Partow, 2001), and the diversion of the Karun River into Iranian terrene (Hameed and Aljorany 2011).

Microbial and heavy metal from polluted industrial sewage may contaminate the turtle feeding grounds. Also other studies in Iraq reported Microbial contamination and heavy metal contamination in fish and other organisms from Garmat Ali river and Shatt Al-Arab river (Abed et al., 2016; Al-kanany et al., 2017; Bannai et al., 2017; Al-Khafaji et al., 2018). The objectives of this study are to determine microbial contamination and qualities in skin, intestine content and muscle tissues of *Glemmys caspica* Turtles in Garmat Ali- River. This study is crucial and may be used as a conservation indicator for the survival of freshwater turtles in this region.

2. Materials And Methods

Collection of samples:

Garmat Ali River, Basrah city, southern of Iraq. it is a waterway between the East Hammar marsh and the Shatt Al-Arab River (Fig. 1). It is about 6 km, 280 m width and the mean depth is 9 m. The river is affected by the tidal current of the Arabian Gulf through the Shatt Al-Arab River. Samples were collected from the one site on along the river, near from Basrah university site in Garmat Ali River during summer 2017. Samples were collected at a depth of 20 – 100 cm. The predominant vegetations on the banks were *Phragmites australis*, and *Typha domingensis*, whereas *Ceratophyllum demersum* and *Potamogeton crispus* were dominant in the deeper areas.

A total of (20) fresh water turtles were collected randomly by hand net captured. Were collected aseptically and immediately transported in a thermal bag to the laboratory and processed within 3hrs of acquisition.

The bacterial counts on the external surfaces were estimated as follows:

Skin surfaces: Sample from some locations of the skin of 20 life turtles was taken by rubbing the sterilized cotton swab over the skin and then inoculated into 9ml of Nutrient broth, MacConkey broth and Selenite F broth which are dispensed in separate tubes. 10 fold serial dilution of the bacterial suspension already inoculated in peptone water was prepared in duplicate and viable aerobic bacterial counts were enumerated using 0.1ml and 1ml inoculums in standard plate count agar as described by Slaby *et al.*, (1981), and then incubated at 37°C for 48 hrs.

Total bacteria counts were determined using pour plate method with MacConkey agar, EMB Agar respectively. Mueller-Hinton Agar for *Pseudomonas spp.* *Salmonella spp.* and *Shigella spp.*, were enumerated using Salmonella Shigella Agar (SSA) and Thiosulphate Citrate Bile Salt Sucrose (TCBS) agar for pathogenic *Vibrio spp.* The plates were incubated at 37 °C for 24h.

Heavy Metal Analysis: The turtles skin samples were homogenized separately in a mortar and weighed

accurately in a porcelain crucible. Before ashing, 1ml of concentrated HNO₃ was added to the samples and allowed to pre-digest overnight in order to reduce losses of volatile metals. The samples were charred on an electric hot plate before ashing in a muffle furnace at 550°C for 4hrs. The white ash was dissolved in 5 ml of 1:1 HCl and a solution made in a 50 ml standard flask [15]. Metal concentrations of the samples were read against appropriate blank and standard solutions using a Perkin-Elmer model 306 Atomic Absorption spectrophotometer (AAS). A blank solution for the biotic samples was made by diluting 1ml concentrated HNO₃ with 5ml 1:1HCl and a 50ml solutions made up with distilled water. the individual metals was expressed on a dry weight basis as mg/kg and the data generated were analyzed statistically.

3. Results

In this study, the total plate count (TPC) of the 20 turtles samples of two Turtles species was analyses in skin had the highest number of bacteria in *M. c. caspica* with 29.60 x 10⁵ cfu/ml. the total plate count had the highest number in the *R. euphraticus* – 19.26 x 10⁵ cfu/ml showed in Table. 1; revealed the isolation of *Pseudomonas spp.* In the skin of *M. c. caspica* having the highest number in (53.73 x 10³cfu/ml), while in the *R. euphraticus* skin had the highest number of *Pseudomonas* isolates. Isolated had the total count of 37.07 x 10³ cfu/ml from the *R. euphraticus* samples. No isolation of *Vibrio spp.* on the *M. c. caspica* skin while *R. euphraticus* skin had a high count of 7.19x 10² cfu/ml (Table 1). *E. coli* isolation showed the highest count in skin of the *M. c. caspica* 85.45x10⁴ while *R. euphraticus* skin (26.18 x 10⁴ cfu/ml). The *Staphylococcus spp.* had a low isolation rate in all samples of two species were analyses as generally compared with other isolated from other organisms except *Vibrio spp.* that had the lowest (Table 1). *Salmonella spp.* and *Shigella spp.* are not isolated in all turtle's samples analyses in this study.

Table 1. The total number of bacteria count cfu/ml in skin of two Turtles species.

Turtles species	TPC Cfu/ml	Pseudomonas spp. Cfu/ml	E. coli Cfu/ml	Staphylococcus spp. Cfu/ml	Vibrio spp Cfu/ml	S/S Cfu/ml
<i>M. c. caspica</i>	29.60x10 ⁵	53.73X10 ³	85.45X10 ⁴	4.46X10 ²	NI	NI
<i>R. euphraticus</i>	19.26x10 ⁵	37.07X10 ³	26.18X10 ⁴	3.15X10 ²	1.19X10 ²	NI

Where: TPC- Total Plate Count, S/S – *Salmonella spp./Shigella spp.*, NI – No isolation

All bacterial isolates including pathogens antagonistic isolates were identified using biochemical and morphological tests. The species of bacteria

isolated were identified and predominant by gram-negative bacteria. Of these, (*Escherichia coli*, *Pseudomonas spp.*, *Vibrio spp.* and *Staphylococcus*

aureus). Were found common in skin of two species Turtles. Table (2). *Escherichia coli* had the highest frequency of 90% while *Streptococcus* spp. and *Vibrio*

spp. had a highest frequency of 10% respectively. Table (3).

Table 2: Morphological and Biochemical characteristics of bacteria species isolated from sampling of the skin of two species Turtles.

bacteria species.	<i>Pseudomonas spp.</i>	<i>Escherichia coli</i>	<i>Vibrio spp.</i>	<i>Staphylococcus spp.</i>
Reaction agent				
Gram staining Shape	G ^{-ve} bacilli	G ^{-ve} bacilli	G ^{-ve} bacilli	G ^{+ve} cocci
Motility	M	M	M	Ma
Indole test	+	-	-	-
Methyl red test	+	-	-	-
Voges proskuer test	-	-	-	-
Citrate utilization test	-	+	-	+
Urease test	-	+	+	+
TSI test		A/A+G	A/A	K/K
H2S	-	-	+	-
Gas	-	+	-	+
Nitrate reduction test	+	-	-	-
Catalase test	+	+	+	+
Oxidase test	+	+	+	+
Carbohydrate test	+	+	-	+
Glucose	+	+	-	+
Maltose	-	-	-	-
Sucrose	-	-	-	-

Note: - Negative, + Positive, A/A -Glucose & lactose and /or sucrose fermentation, K/K -no fermentation, A/A+G- Glucose & lactose and /or sucrose fermentation+ gas produced.

Table 3: Frequency distribution of bacteria isolates from skin samples of two species Turtles

Bacteria species isolate	Percentage frequency (100%)
<i>Escherichia coli</i>	90
<i>Pseudomonas spp.</i>	76
<i>Vibrio spp.</i>	10
<i>Staphylococcus spp.</i>	10

Table 4 shows the levels of Fe, Cu and Pb that were expressed in mg/kg dry weight sample. The data clearly shows variation in the level of heavy metals among the freshwater turtles species sampled.

The analysis of the selected metals in the present study revealed an order of Fe>Cu>Pb in two freshwater turtles species.

M. c. caspica Turtle had iron concentration of 21.52±0.031 mg/kg while *R. euphraticus* Turtle had iron concentration of 34.69±0.152 mg/kg. *M. c. caspica* and *R. euphraticus* had copper concentrations of 0.87±0.034 and 0.92±0.057 mg/kg respectively. *M. c. caspica* had a lead concentration of 0.06±0.041 mg/kg while *R. euphraticus* had a lead concentration of 0.09±0.064 mg/kg.

Table 4: Heavy metals concentration (mg/kg) in two Turtles species

Turtles species	Fe	Cu	Pb
<i>M. c. caspica</i>	21.52±0.031	0.87±0.034	0.06±0.041
<i>R. euphraticus</i>	34.69±0.152	0.92±0.057	0.09±0.064

4. Discussion

Specimens of *M. c. caspica* were collected from the Garmat Ali canal because this station that good conditions in comparison with the other sites in shatt al-arab river.

Garmat Ali station has alluvial soil that is suitable for *M. c. caspica* nesting (Kami et al. 2006) added to calm waters and low salinity rate of their tributaries which created a favorable conditions for turtles (Ghaffari et al. 2008).

Continuous exposure of organisms, such as freshwater turtles, to heavy metals resulted in chronic intoxication which may be fatal (Harper et al,2007; Storelli et al, 2008).

heavy metal deposition from industrial activities, aquaculture activities, treated and untreated municipal wastewater from urban developments may lead to heavy metals accumulation in the feeding habitats. (Al-Musharafi et al, 2014a; 2014b).

Fe is an essential element in human diet and Iron forms part of hemoglobin which allows oxygen to be carried from the lungs to the tissues.

In this study, The copper concentrations were similar to other studies of (Yilmaz, 2009; Rejomon et al. 2010; Olowu et al. 2010).

In this study, Higher concentration of lead is known to inhibit active transport mechanisms involving ATP and may also suppress cellular oxidation-reduction reactions and even inhibit of protein synthesis (Adeyeye et al, 1996). The level of lead from this study, could not be said to pose any health risk since the values were within the FAO permissible limit of 0.3 mg/kg.

In addition to heavy metals, several studies related to pathogenic bacteria were published, in Iraq some studies revealed that sewage contaminated effluents contribute to environmental pollution (Abed et al 2016; Alkanany et al. 2017; Al-Khafaji et al., 2018).

In parts of the world, including Iraq, pathogenic bacteria were isolated from turtles, fish and fresh water habitats.

References

1. Abed HH, Kadhim ZY, Abed SM, Mahdi ZS. 2016, Levels of bacterial and chemical pollutants in Euphrates river in Samawah, Iraq. *World J Exp Biosci* 4: 66-69.
2. Adeyeye, E.T., N.J. Akinyugha, M.E. Fesobi and 27. FEPA, 2003. Guideline and V.O. Tenabe, 1996. Determination of some metals *Clarias gariepinus* (Cuvier and Valenciennes), *Cyprinus carpio* (L.) and *Oreochromis niloticus* (L.) fishes in a polyculture fresh water pond and their environments. *Aquaculture*, 147: 205-214.
3. Al-Barwari SE, Saeed I. 2007. On the Helminth Fauna of Some Iraqi Reptiles. *Türkiye Parazitoloji Dergisi*, 31 (4): 330-336.
4. Al-kanany FNA, Rasheed BA, Khudiar MM, 2017, Determination of Bacterial Contamination and Some Chemical Parameters in Basrah Governorate Rivers, *Journal of Pharmaceutical, Chemical and Biological Sciences*, 5(2): 118-124.
5. Al-Khafaji K.K., Al-Malki G.M., and Karim R.M., 2018, Biochemical composition and heavy metal accumulation in tissues of the blue crab *Portunus pelagicus* (Linnaeus, 1766) from NW of Arabian Gulf, South Iraq, *International Journal of Marine Science*, 8(17): 146-150 (doi: 10.5376/ijms.2018.08.0017).
6. Al-Mahmood, H.K.H, Hassan, W.F., Alhello, A.Z.A., Hammood, A.I. & Muhson, N.K. (2015). Impact of low discharge and drought of the water quality of the Shatt Al Arab and Al-Basrah Rivers (south of Iraq). *J. Int. Acad. Res. Multidisciplinary*. 3(1), pp. 285-296.
7. Al-Musharafi, SK. Mahmoud IY. and Al-Bahry SN. Environmental hazards and pollution from liquid waste lagoons. *IPCBE 69*, pp. 1-5. 2014b.
8. Al-Musharafi, SK. Mahmoud IY. and Al-Bahry SN. Environmental contamination by industrial effluents and sludge relative to heavy metals. *GEP 2*, pp. 14-18. 2014a.
9. Al-Tememi, M.K., Hussein, M.A., Khaleefa, U.Q., Ghaleb, H.B., AL-Mayah, A.M. & A.J. Ruhmah, A.J. (2015). The Salts diffusion between East Hammar marsh area and Shatt Al-Arab River Northern Basra City. *Marsh Bulletin*, 10, pp. 36-45.
10. Ghaffari H. 2002. Investigation of Rafetus euphraticus in Iran. BSc Project, Azad University, Faculty of Technical and Engineering: Tehran, Iran.
11. Hameed, A.H. & Aljorany, Y.S. (2011). Investigation on nutrient behavior along Shatt Al-Arab River, Basrah, Iraq. *J. Appl. Sci. Res.*, 7, pp. 1340-1345.
12. Harper, ER. Leger, ST. Westberg, JA. Mazzaro, L. Schmitt, T. Reidarson, TH. Tucker, M. Cross, DH. and Puschner, B. Tissue heavy metal concentrations of stranded California sea lions (*Zalophus californianus*) in Southern California. *Environ. Pollut.* 147, pp. 677-682. 2007.
13. Kami HG, Hojati V, Rad SP, Sheidaee M. 2006. A biological study of the European Pond Turtle, *Emys orbicularis persica*, and the Caspian Pond Turtle, *Mauremys caspica caspica*, in the

- Golestan and Mazandaran provinces of Iran. *Zoology in the Middle East* 37: 21–28.
14. Khalaf, K.T. 1959. Reptiles of Iraq with some notes on amphibian. AlRabitta press, Baghdad, Iraq. 96pp.
 15. Majid A.A. Bannai, Essa T. Mohammad, Ahmed J. Shamary, Khassn A.Al- Najjar, Fawzi M. Al-Khwaja, Mohammed A. Aldoghachi. A contribution and pathology of E.coli and Pseudomonas aeruginosa bacterial infection on Soft shell turtle Rafetus euphraticus (Gray, 1864) East Hammar marshes, Iraq // *Russian Journal of Parasitology*, 2017, V. 40, Iss. 2, pp.
 16. Okoye, B.C.O., 1991. Heavy metals and organisms in the Lagos Lagoon. *International Journal Environmental Studies*, 37: 285-292.
 17. Olowu, R.A., O.O. Ayejuyo, G.O. Adewuyi, I.A. Adejoro, A.A.B. Denloye, A.O. Babatunde and A.L. Ogundajo, 2010. Determination of heavy metals in fish tissues, water and sediment from Epe and Badagry lagoons, Lagos, Nigeria. *E- J. Chem.*, 71: 215-221.
 18. Partow, H. (2001). *The Mesopotamian Marshlands: Demise of an ecosystem*. Nairobi (Kenya): Division of early warning and assessment, United Nation for Environmental Programs: UNEP publication UNEP/DEWA, 103p.
 19. Pritchard PCH. 1967. *Living Turtles of the World*. Jersey City, NJ: T.F.H. Publications, 288 pp.
 20. Pritchard PCH. 1979. *Encyclopedia of Turtles*. Neptune, NJ: T.F.H. Publications, 895 pp.
 21. Rejomon, G., M. Nair and T. Joseph, 2010, Trace metal Dynamics in fishes from the southwest coast of India. *Microbial Environ. Monit. Assess.*, 167: 234-255.
 22. Slaby BM, Martin RE, Ramsdell GE. Reproducibility of Microbiological counts on frozen Cod: A collaborative study. *J. Food Sci.* 1981;46(3):716-719.
 23. Storelli, MM. Barone, G. Storelli, A. and Marcotrigiano, GO. Total and subcellular distribution of trace elements (Cd, Cu and Zn) in the liver and kidney of green turtles (*Chelonia mydas*) from the Mediterranean Sea. *Chemosphere*. 70, p. 908-913. 2008.
 24. Yilmaz, F., 2009. The comparison of heavy metal concentrations (Cd, Cu, Mn, Pb and Zn) in tissues of three economically important fish (*Anguilla anguilla*, *Mugil cephalus* and *Oreochromis niloticus*) inhabiting Koycegiz lake-Mugla. *Turkish Journal of Science and Technology*, 4: 7-15.

11/24/2019