

In 1991 Fermat's Last Theorem Has Been Proved (I)

Jiang, Chun-Xuan

Institute for Basic Research, Palm Harbor, FL34682-1577, USA

And: P. O. Box 3924, Beijing 100854, China

jiangchunxuan@sohu.com, cxjiang@mail.bcf.net.cn, jcxuan@sina.com, Jiangchunxuan@vip.sohu.com,
jcxxxx@163.com

Abstract: In 1637 Fermat wrote: “It is impossible to separate a cube into two cubes, or a biquadrate into two biquadrates, or in general any power higher than the second into powers of like degree: I have discovered a truly marvelous proof, which this margin is too small to contain.” This means: $x^n + y^n = z^n$ ($n > 2$) has no integer solutions, all different from 0 (i.e., it has only the trivial solution, where one of the integers is equal to 0). It has been called Fermat's last theorem (FLT). It suffices to prove FLT for exponent 4. and every prime exponent P . Fermat proved FLT for exponent 4. Euler proved FLT for exponent 3. On October 25, 1991 we proved Fermat last theorem. [Jiang, Chun-Xuan. **In 1991 Fermat's Last Theorem Has Been Proved (I).** *Researcher* 2016;8(8):15-20]. ISSN 1553-9865 (print); ISSN 2163-8950 (online). <http://www.sciencepub.net/researcher>. 4. doi:[10.7537/marsrj080816.04](https://doi.org/10.7537/marsrj080816.04).

Keywords: 1991; Fermat's Last Theorem; prime exponent P

In this paper using the complex hyperbolic functions we prove FLT for exponents $3P$ and P , where P is an odd prime. The proof of FLT must be direct. But indirect proof of FLT is disbelieving.

In 1974 Jiang found out Euler formula of the cyclotomic real numbers in the cyclotomic fields

$$\exp\left(\sum_{i=1}^{n-1} t_i J^i\right) = \sum_{i=1}^n S_i J^{i-1} \quad (1)$$

where J denotes a n th root of unity, $J^n = 1$, n is an odd number, t_i are the real numbers.

S_i is called th complex hyperbolic functions of order n with $n-1$ variables [1-7].

$$S_i = \frac{1}{n} \left[e^A + 2 \sum_{j=1}^{\frac{n-1}{2}} (-1)^{(i-1)j} e^{B_j} \cos\left(\theta_j + (-1)^j \frac{(i-1)j\pi}{n}\right) \right] \quad (2)$$

2000 mathematics subject classification 11D41.

Key words complex hyperbolic, Euler proof.

where $i=1,2,\dots,n$;

$$A = \sum_{\alpha=1}^{n-1} t_\alpha, \quad B_j = \sum_{\alpha=1}^{n-1} t_\alpha (-1)^{\alpha j} \cos \frac{\alpha j \pi}{n}, \\ \theta_j = (-1)^{j+1} \sum_{\alpha=1}^{n-1} t_\alpha (-1)^{\alpha j} \sin \frac{\alpha j \pi}{n}, \quad A + 2 \sum_{j=1}^{\frac{n-1}{2}} B_j = 0 \quad (3)$$

(2) may be written in the matrix form

$$\begin{bmatrix} S_1 \\ S_2 \\ S_3 \\ \dots \\ S_n \end{bmatrix} = \frac{1}{n} \begin{bmatrix} 1 & 1 & 0 & \dots & 0 \\ 1 & -\cos \frac{\pi}{n} & -\sin \frac{\pi}{n} & \dots & -\sin \frac{(n-1)\pi}{2n} \\ 1 & \cos \frac{2\pi}{n} & \sin \frac{2\pi}{n} & \dots & -\sin \frac{(n-1)\pi}{n} \\ \dots & \dots & \dots & \dots & \dots \\ 1 & \cos \frac{(n-1)\pi}{n} & \sin \frac{(n-1)\pi}{n} & \dots & -\sin \frac{(n-1)^2\pi}{2n} \end{bmatrix} \begin{bmatrix} e^A \\ 2e^{B_1} \cos \theta_1 \\ 2e^{B_1} \sin \theta_1 \\ \dots \\ 2 \exp B_{\frac{n-1}{2}} \sin \theta_{\frac{n-1}{2}} \end{bmatrix} \quad (4)$$

where $(n-1)/2$ is an even number.

From (4) we have its inverse transformation

$$\begin{bmatrix} e^A \\ e^{B_1} \cos \theta_1 \\ e^{B_1} \sin \theta_1 \\ \dots \\ \exp(B_{\frac{n-1}{2}}) \sin(\theta_{\frac{n-1}{2}}) \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 & \dots & 1 \\ 1 & -\cos \frac{\pi}{n} & \cos \frac{2\pi}{n} & \dots & \cos \frac{(n-1)\pi}{n} \\ 0 & -\sin \frac{\pi}{n} & \sin \frac{2\pi}{n} & \dots & \sin \frac{(n-1)\pi}{n} \\ \dots & \dots & \dots & \dots & \dots \\ 0 & -\sin \frac{(n-1)\pi}{2n} & -\sin \frac{(n-1)\pi}{n} & \dots & -\sin \frac{(n-1)^2\pi}{2n} \end{bmatrix} \begin{bmatrix} S_1 \\ S_2 \\ S_3 \\ \dots \\ S_n \end{bmatrix} \quad (5)$$

From (5) we have

$$\begin{aligned} e^A &= \sum_{i=1}^n S_i, \quad e^{B_j} \cos \theta_j = S_1 + \sum_{i=1}^{n-1} S_{1+i} (-1)^{ij} \cos \frac{ij\pi}{n} \\ e^{B_j} \sin \theta_j &= (-1)^{j+1} \sum_{i=1}^{n-1} S_{1+i} (-1)^{ij} \sin \frac{ij\pi}{n}, \end{aligned} \quad (6)$$

In (3) and (6) t_i and S_i have the same formulas. (4) and (5) are the most critical formulas of proofs for FLT. Using (4) and (5) in 1991 Jiang invented that every factor of exponent n has the Fermat equation and proved FLT [1-7] Substituting (4) into (5) we prove (5).

$$\begin{bmatrix} e^A \\ e^{B_1} \cos \theta_1 \\ e^{B_1} \sin \theta_1 \\ \dots \\ \exp(B_{\frac{n-1}{2}}) \sin(\theta_{\frac{n-1}{2}}) \end{bmatrix} = \frac{1}{n} \begin{bmatrix} 1 & 1 & 1 & \dots & 1 \\ 1 & -\cos \frac{\pi}{n} & \cos \frac{2\pi}{n} & \dots & \cos \frac{(n-1)\pi}{n} \\ 0 & -\sin \frac{\pi}{n} & \sin \frac{2\pi}{n} & \dots & \sin \frac{(n-1)\pi}{n} \\ \dots & \dots & \dots & \dots & \dots \\ 0 & -\sin \frac{(n-1)\pi}{2n} & -\sin \frac{(n-1)\pi}{n} & \dots & -\sin \frac{(n-1)^2\pi}{2n} \end{bmatrix} \times$$

$$\begin{aligned}
& \begin{bmatrix} 1 & 1 & 0 & \cdots & 0 \\ 1 & -\cos \frac{\pi}{n} & -\sin \frac{\pi}{n} & \cdots & -\sin \frac{(n-1)\pi}{2n} \\ 1 & \cos \frac{2\pi}{n} & \sin \frac{2\pi}{n} & \cdots & -\sin \frac{(n-1)\pi}{n} \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ 1 & \cos \frac{(n-1)\pi}{n} & \sin \frac{(n-1)\pi}{n} & \cdots & -\sin \frac{(n-1)^2\pi}{2n} \end{bmatrix} \begin{bmatrix} e^A \\ 2e^{B_1} \cos \theta_1 \\ 2e^{B_1} \sin \theta_1 \\ \cdots \\ 2 \exp(B_{\frac{n-1}{2}}) \sin(\theta_{\frac{n-1}{2}}) \end{bmatrix} \\
& = \frac{1}{n} \begin{bmatrix} n & 0 & 0 & \cdots & 0 \\ 0 & \frac{n}{2} & 0 & \cdots & 0 \\ 0 & 0 & \frac{n}{2} & \cdots & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & 0 & \cdots & \frac{n}{2} \end{bmatrix} \begin{bmatrix} e^A \\ 2e^{B_1} \cos \theta_1 \\ 2e^{B_1} \sin \theta_1 \\ \cdots \\ 2 \exp(B_{\frac{n-1}{2}}) \sin(\theta_{\frac{n-1}{2}}) \end{bmatrix} \\
& = \begin{bmatrix} e^A \\ e^{B_1} \cos \theta_1 \\ e^{B_1} \sin \theta_1 \\ \cdots \\ \exp(B_{\frac{n-1}{2}}) \sin(\theta_{\frac{n-1}{2}}) \end{bmatrix}, \tag{7}
\end{aligned}$$

where $1 + \sum_{j=1}^{n-1} (\cos \frac{j\pi}{n})^2 = \frac{n}{2}$, $\sum_{j=1}^{n-1} (\sin \frac{j\pi}{n})^2 = \frac{n}{2}$.

From (3) we have

$$\exp(A + 2 \sum_{j=1}^{\frac{n-1}{2}} B_j) = 1 \tag{8}$$

From (6) we have

$$\exp(A + 2 \sum_{j=1}^{\frac{n-1}{2}} B_j) = \begin{vmatrix} S_1 & S_n & \cdots & S_2 \\ S_2 & S_1 & \cdots & S_3 \\ \cdots & \cdots & \cdots & \cdots \\ S_n & S_{n-1} & \cdots & S_1 \end{vmatrix} = \begin{vmatrix} S_1 & (S_1)_1 & \cdots & (S_1)_{n-1} \\ S_2 & (S_2)_1 & \cdots & (S_2)_{n-1} \\ \cdots & \cdots & \cdots & \cdots \\ S_n & (S_n)_1 & \cdots & (S_n)_{n-1} \end{vmatrix}, \tag{9}$$

$$(S_i)_j = \frac{\partial S_i}{\partial t_j}$$

where [7].

From (8) and (9) we have the circulant determinant

$$\exp(A + 2 \sum_{j=1}^{\frac{n-1}{2}} B_j) = \begin{vmatrix} S_1 & S_n & \cdots & S_2 \\ S_2 & S_1 & \cdots & S_3 \\ \cdots & \cdots & \cdots & \vdots \\ S_n & S_{n-1} & \cdots & S_1 \end{vmatrix} = 1 \quad (10)$$

If $S_i \neq 0$, where $i = 1, 2, \dots, n$, then (10) has infinitely many rational solutions.

Assume $S_1 \neq 0, S_2 \neq 0, S_i = 0$ where $i = 3, 4, \dots, n$. $S_i = 0$ are $n-2$ indeterminate equations with $n-1$ variables. From (6) we have

$$e^A = S_1 + S_2, \quad e^{2B_j} = S_1^2 + S_2^2 + 2S_1S_2(-1)^j \cos \frac{j\pi}{n}. \quad (11)$$

From (10) and (11) we have the Fermat equation

$$\exp(A + 2 \sum_{j=1}^{\frac{n-1}{2}} B_j) = (S_1 + S_2) \prod_{j=1}^{\frac{n-1}{2}} (S_1^2 + S_2^2 + 2S_1S_2(-1)^j \cos \frac{j\pi}{n}) = S_1^n + S_2^n = 1 \quad (12)$$

Example[1]. Let $n = 15$. From (3) we have

$$A = (t_1 + t_{14}) + (t_2 + t_{13}) + (t_3 + t_{12}) + (t_4 + t_{11}) + (t_5 + t_{10}) + (t_6 + t_9) + (t_7 + t_8)$$

$$B_1 = -(t_1 + t_{14}) \cos \frac{\pi}{15} + (t_2 + t_{13}) \cos \frac{2\pi}{15} - (t_3 + t_{12}) \cos \frac{3\pi}{15} + (t_4 + t_{11}) \cos \frac{4\pi}{15}$$

$$-(t_5 + t_{10}) \cos \frac{5\pi}{15} + (t_6 + t_9) \cos \frac{6\pi}{15} - (t_7 + t_8) \cos \frac{7\pi}{15},$$

$$B_2 = (t_1 + t_{14}) \cos \frac{2\pi}{15} + (t_2 + t_{13}) \cos \frac{4\pi}{15} + (t_3 + t_{12}) \cos \frac{6\pi}{15} + (t_4 + t_{11}) \cos \frac{8\pi}{15}$$

$$+ (t_5 + t_{10}) \cos \frac{10\pi}{15} + (t_6 + t_9) \cos \frac{12\pi}{15} + (t_7 + t_8) \cos \frac{14\pi}{15},$$

$$B_3 = -(t_1 + t_{14}) \cos \frac{3\pi}{15} + (t_2 + t_{13}) \cos \frac{6\pi}{15} - (t_3 + t_{12}) \cos \frac{9\pi}{15} + (t_4 + t_{11}) \cos \frac{12\pi}{15}$$

$$-(t_5 + t_{10}) \cos \frac{15\pi}{15} + (t_6 + t_9) \cos \frac{18\pi}{15} - (t_7 + t_8) \cos \frac{21\pi}{15},$$

$$B_4 = (t_1 + t_{14}) \cos \frac{4\pi}{15} + (t_2 + t_{13}) \cos \frac{8\pi}{15} + (t_3 + t_{12}) \cos \frac{12\pi}{15} + (t_4 + t_{11}) \cos \frac{16\pi}{15}$$

$$+ (t_5 + t_{10}) \cos \frac{20\pi}{15} + (t_6 + t_9) \cos \frac{24\pi}{15} + (t_7 + t_8) \cos \frac{28\pi}{15},$$

$$B_5 = -(t_1 + t_{14}) \cos \frac{5\pi}{15} + (t_2 + t_{13}) \cos \frac{10\pi}{15} - (t_3 + t_{12}) \cos \frac{15\pi}{15} + (t_4 + t_{11}) \cos \frac{20\pi}{15}$$

$$-(t_5 + t_{10}) \cos \frac{25\pi}{15} + (t_6 + t_9) \cos \frac{30\pi}{15} - (t_7 + t_8) \cos \frac{35\pi}{15},$$

$$B_6 = (t_1 + t_{14}) \cos \frac{6\pi}{15} + (t_2 + t_{13}) \cos \frac{12\pi}{15} + (t_3 + t_{12}) \cos \frac{18\pi}{15} + (t_4 + t_{11}) \cos \frac{24\pi}{15}$$

$$\begin{aligned}
 & + (t_5 + t_{10}) \cos \frac{30\pi}{15} + (t_6 + t_9) \cos \frac{36\pi}{15} + (t_7 + t_8) \cos \frac{42\pi}{15}, \\
 B_7 = & -(t_1 + t_{14}) \cos \frac{7\pi}{15} + (t_2 + t_{13}) \cos \frac{14\pi}{15} - (t_3 + t_{12}) \cos \frac{21\pi}{15} + (t_4 + t_{11}) \cos \frac{28\pi}{15} \\
 & - (t_5 + t_{10}) \cos \frac{35\pi}{15} + (t_6 + t_9) \cos \frac{42\pi}{15} - (t_7 + t_8) \cos \frac{49\pi}{15}, \\
 A + 2 \sum_{j=1}^7 B_j = & 0, \quad A + 2B_3 + 2B_6 = 5(t_5 + t_{10})
 \end{aligned} \tag{13}$$

From (12) we have the Fermat equation

$$\exp(A + 2 \sum_{j=1}^7 B_j) = S_1^{15} + S_2^{15} = (S_1^5)^3 + (S_2^5)^3 = 1 \tag{14}$$

From (13) we have

$$\exp(A + 2B_3 + 2B_6) = [\exp(t_5 + t_{10})]^5 \tag{15}$$

From (11) we have

$$\exp(A + 2B_3 + 2B_6) = S_1^5 + S_2^5 \tag{16}$$

From (15) and (16) we have the Fermat equation

$$\exp(A + 2B_3 + 2B_6) = S_1^5 + S_2^5 = [\exp(t_5 + t_{10})]^5 \tag{17}$$

Euler proved that (14) has no rational solutions for exponent 3[8]. Therefore we prove that (17) has no rational solutions for exponent 5[1].

Theorem 1. [1-7]. Let $n = 3P$, where $P > 3$ is odd prime. From (12) we have the Fermat's equation

$$\exp(A + 2 \sum_{j=1}^{\frac{3P-1}{2}} B_j) = S_1^{3P} + S_2^{3P} = (S_1^P)^3 + (S_2^P)^3 = 1 \tag{18}$$

From (3) we have

$$\exp(A + 2 \sum_{j=1}^{\frac{P-1}{2}} B_{3j}) = [\exp(t_p + t_{2p})]^p \tag{19}$$

From (11) we have

$$\exp(A + 2 \sum_{j=1}^{\frac{P-1}{2}} B_{3j}) = S_1^P + S_2^P \tag{20}$$

From (19) and (20) we have the Fermat equation

$$\exp(A + 2 \sum_{j=1}^{\frac{P-1}{2}} B_{3j}) = S_1^P + S_2^P = [\exp(t_p + t_{2p})]^p \tag{21}$$

Euler proved that (18) has no rational solutions for exponent 3[8]. Therefore we prove that (21) has no rational solutions for $P > 3$ [1, 3-7].

Note. Wiles had not proved Fermat last theorem[9-11]

References

1. Jiang, C-X, Fermat last theorem had been proved, Potential Science (in Chinese), 2.17-20 (1992), Preprints (in English) December (1991). <http://www.wbabin.net/math/xuan47.pdf>.
2. Jiang, C-X, Fermat last theorem had been proved by Fermat more than 300 years ago, Potential Science (in Chinese), 6.18-20(1992).
3. Jiang, C-X, On the factorization theorem of circulant determinant, Algebras, Groups and Geometries, 11. 371-377(1994), MR. 96a: 11023, <http://www.wbabin.net/math/xuan45.pdf>.
4. Jiang, C-X, Fermat last theorem was proved in 1991, Preprints (1993). In: Fundamental open problems in science at the end of the millennium, T.Gill, K. Liu and E. Trell (eds). Hadronic Press, 1999, P555-558. <http://www.wbabin.net/math/xuan46.pdf>.
5. Jiang, C-X, On the Fermat-Santilli theorem, Algebras, Groups and Geometries, 15. 319-349(1998).
6. Jiang, C-X, Complex hyperbolic functions and Fermat's last theorem, Hadronic Journal Supplement, 15. 341-348(2000).
7. Jiang, C-X, Foundations of Santilli Isonumber Theory with applications to new cryptograms, Fermat's theorem and Goldbach's Conjecture. Inter, Acad. Press. 2002. MR2004c:11001, <http://www.wbabin.net/math/xuan13.pdf>. <http://www.i-b-r.org/docs/jiang.pdf>.
8. Ribenboim, P, Fermat last theorem for amateur, Springer-Verlag, (1999).
9. Wiles, A, Modular elliptic curves and Fermat last theorem, Ann. Math 141(3),443-551(1995).
10. Zhivolov,Y, Fermat last theorem and mistakes of Andrew Wiles.2006. www.baoway.com/bbs/viewthread.php?tid=1816_2&fpage=3.
11. Zhivolov,Y, Fermat last theorem and Kenneth Ribet mistakes.2006.
12. Jiang, Chun-Xuan (蒋春暄). Automorphic Functions And Fermat's Last Theorem (1). *Academ Arena* 2016;8(5):78-84]. ISSN 1553-992X (print); ISSN 2158-771X (online). <http://www.sciencepub.net/academia>. 8. doi:[10.7537/marsaaj080516.08](https://doi.org/10.7537/marsaaj080516.08).
13. Jiang, Chun-Xuan (蒋春暄). Automorphic Functions And Fermat's Last Theorem (3) (Fermat's Proof of FLT). *Academ Arena* 2016;8(5):85-92]. ISSN 1553-992X (print); ISSN 2158-771X (online). <http://www.sciencepub.net/academia>. 9. doi:[10.7537/marsaaj080516.09](https://doi.org/10.7537/marsaaj080516.09).
14. Jiang, Chun-Xuan (蒋春暄). Riemann Paper (1859) Is False. *Academ Arena* 2016;8(6):35-39]. ISSN 1553-992X (print); ISSN 2158-771X (online). <http://www.sciencepub.net/academia>. 7. doi:[10.7537/marsaaj080616.07](https://doi.org/10.7537/marsaaj080616.07).
15. Jiang, Chun-Xuan (蒋春暄). The New Prime theorems (341) - (390) . *Academ Arena* 2016;8(6):61-107]. ISSN 1553-992X (print); ISSN 2158-771X (online). <http://www.sciencepub.net/academia>. 12. doi:[10.7537/marsaaj080616.12](https://doi.org/10.7537/marsaaj080616.12).
16. Ma H. The Nature of Time and Space. Nature and science 2003;1(1):1-11. doi:[10.7537/marsnsj010103.01](https://doi.org/10.7537/marsnsj010103.01). <http://www.sciencepub.net/nature/0101/01-ma.pdf>.
17. Wikipedia. The free encyclopedia. <http://en.wikipedia.org>. 2016.
18. 马宏宝. 宙永恒吗? New York Science Journal. 2008;1(3):66-69. ISSN: 1554-0200. doi:[10.7537/marsnys010308.07](https://doi.org/10.7537/marsnys010308.07). http://www.sciencepub.net/newyork/0103/07_0026_mahongbao_universe.pdf.